

Performance Measures

Identifier	Measure	Measure Description	Measure Steward	Federal Reporting Program	Target Population	Condition	Specification Source	2020 (2019 data measurement year) and Subsequent Years Target for Improvement
PTB \$\$	Initiation of Injectable Progesterone for Preterm Birth Prevention	The percentage of women 15-45 years of age with evidence of a previous preterm singleton birth event (24-36 weeks completed gestation) who received one or more progesterone injections between the 16th and 24th week of gestation for deliveries during the measurement year.	State	None	Children's and Maternal Health	Perinatal and Reproductive Health	Section V	24.08
AWC \$\$	Adolescent Well Care Visit	The percentage of enrolled members 12-21 years of age who had at least one comprehensive well-care visit with a PCP or OB/GYN practitioner during the measurement year.	NCQA	CHIPRA	Children's Health	Utilization	HEDIS	NCQA Quality Compass Medicaid National 50th percentile [All LOBs (Excluding PPOs and EPOs): Average] for the year prior to the measurement year
ADD \$\$	Follow-up Care for Children Prescribed ADHD Medication-Initiation Phase	The percentage of children 6-12 years of age as of the index period start date with a newly prescribed ambulatory prescription dispensed for attention-deficit/hyperactivity disorder (ADHD) medication, who had one follow-up visit with a practitioner with prescribing authority during the 30-day Initiation Phase.	NCQA	CHIPRA, MU2	Children's Health	Behavioral Health	HEDIS	NCQA Quality Compass Medicaid National 50th percentile [All LOBs (Excluding PPOs and EPOs): Average] for the year prior to the measurement year
ADD \$\$	Follow-up Care for Children Prescribed ADHD Medication-Continuation Phase	The percentage of children 6-12 years of age as of the index period start date with a newly prescribed ambulatory prescription dispensed for attention-deficit/hyperactivity disorder (ADHD) medication, who remained on the medication for at least 210 days and who, in addition to the visit in the Initiation Phase, had at least two follow-up visits with a practitioner within 270 days (9 months) after the Initiation Phase ended.	NCQA	CHIPRA, MU2	Children's Health	Behavioral Health	HEDIS	NCQA Quality Compass Medicaid National 50th percentile [All LOBs (Excluding PPOs and EPOs): Average] for the year prior to the measurement year

Attachment E

Identifier	Measure	Measure Description	Measure Steward	Federal Reporting Program	Target Population	Condition	Specification Source	2020 (2019 data measurement year) and Subsequent Years Target for Improvement
AMB-ED \$\$	Ambulatory Care- ED Visits	This measure summarizes utilization of ambulatory care ED Visits per 1,000 member months.	NCQA	CHIPRA	Population Health	Utilization	HEDIS	NCQA Quality Compass Medicaid National 50th percentile [All LOBs (Excluding PPOs and EPOs): Average] for the year prior to the measurement year
PPC \$\$	Prenatal and Postpartum Care - Timeliness of Prenatal Care	The percentage of deliveries of live births on or between November 6 of the year prior to the measurement year and November 5 of the measurement year that received a prenatal care visit as a member of the organization in the first trimester, on the enrollment start date or within 42 days of enrollment in the organization.	NCQA	MEDICAID ADULT	Maternal Health	Perinatal and Reproductive Health	HEDIS	NCQA Quality Compass Medicaid National 50th percentile [All LOBs (Excluding PPOs and EPOs): Average] for the year prior to the measurement year
PPC \$\$	Prenatal and Postpartum Care – Postpartum Care (PPC Numerator 2)	The percentage of deliveries that had a postpartum visit on or between 21 and 56 days after delivery.	NCQA	MEDICAID ADULT	Maternal Health	Perinatal and Reproductive Health	HEDIS	NCQA Quality Compass Medicaid National 50th percentile [All LOBs (Excluding PPOs and EPOs): Average] for the year prior to the measurement year
FUH \$\$	Follow-Up After Hospitalization for Mental Illness - Within 30 days of discharge	The percentage of discharges for members 6 years of age and older who were hospitalized for treatment of selected mental illness diagnoses and who had a follow-up visit with a mental health practitioner within 30 days of discharge.	NCQA	MEDICAID ADULT	Behavioral Health	Behavioral Health	HEDIS	NCQA Quality Compass Medicaid National 50th percentile [All LOBs (Excluding PPOs and EPOs): Average] for the year prior to the measurement year

Attachment E

Identifier	Measure	Measure Description	Measure Steward	Federal Reporting Program	Target Population	Condition	Specification Source	2020 (2019 data measurement year) and Subsequent Years Target for Improvement
CBP \$\$	Controlling High Blood Pressure - Total	The percentage of members 18-85 years of age who had a diagnosis of hypertension (HTN) and whose blood pressure (BP) was adequately controlled (<140/90 mm Hg) during the measurement year.	NCQA	MEDICAID ADULT, CMS MU2, HEALTH HOMES	Chronic Disease	Cardiovascular Care	HEDIS	NCQA Quality Compass Medicaid National 50th percentile [All LOBs (Excluding PPOs and EPOs): Average] for the year prior to the measurement year
CDC \$\$	Comprehensive Diabetes Care - Hemoglobin A1c (HbA1c) testing	The percentage of members 18-75 years of age with diabetes (type 1 and type 2) with a Hemoglobin A1c (HbA1c) test.	NCQA	MEDICAID ADULT	Chronic Disease	Diabetes	HEDIS	NCQA Quality Compass Medicaid National 50th percentile [All LOBs (Excluding PPOs and EPOs): Average] for the year prior to the measurement year
CDC \$\$	Comprehensive Diabetes Care - Eye exam (retinal) performed	The percentage of members 18-75 years of age with diabetes (type 1 and type 2) with an eye exam (retinal) performed.	NCQA	MEDICAID ADULT	Chronic Disease	Diabetes	HEDIS	NCQA Quality Compass Medicaid National 50th percentile [All LOBs (Excluding PPOs and EPOs): Average] for the year prior to the measurement year
CDC \$\$	Comprehensive Diabetes Care - Medical attention for nephropathy	The percentage of members 18-75 years of age with diabetes (type 1 and type 2) with medical attention for nephropathy.	NCQA	CHIPRA	Chronic Disease	Diabetes	HEDIS	NCQA Quality Compass Medicaid National 50th percentile [All LOBs (Excluding PPOs and EPOs): Average] for the year prior to the measurement year

Attachment E

Identifier	Measure	Measure Description	Measure Steward	Federal Reporting Program	Target Population	Condition	Specification Source	2020 (2019 data measurement year) and Subsequent Years Target for Improvement
W15 \$\$	Well-Child Visits in the First 15 Months of Life - Six or more well-child visits.	The percentage of members who turned 15 months old during the measurement year and who had six or more well-child visits with a PCP during their first 15 months of life.	NCQA	CHIPRA	Children's Health	Utilization	HEDIS	NCQA Quality Compass Medicaid National 50th percentile [All LOBs (Excluding PPOs and EPOs): Average] for the year prior to the measurement year
W34 \$\$	Well-Child Visits in the Third, Fourth, Fifth and Sixth Years of Life	The percentage of members 3-6 years of age who had one or more well-child visits with a PCP during the measurement year.	NCQA	CHIPRA	Children's Health	Utilization	HEDIS	NCQA Quality Compass Medicaid National 50th percentile [All LOBs (Excluding PPOs and EPOs): Average] for the year prior to the measurement year
CPA \$\$	CAHPS Health Plan Survey 5.0H, Adult (Rating of Health Plan, 8+9+10)	This measure provides information on the experiences of Medicaid members with the organization and gives a general indication of how well the organization meets members' expectations.	NCQA	MEDICAID ADULT	Adult	Member Satisfaction	HEDIS	NCQA Quality Compass Medicaid National 50th percentile [All LOBs (Excluding PPOs and EPOs): Average] for the year prior to the measurement year
CPC \$\$	CAHPS Health Plan Survey 5.0H, Child (Rating of Health Plan-General Population, 8+9+10)	This measure provides information on parents' experience with their child's Medicaid organization.	NCQA	MEDICAID, CHIPRA	Child	Member Satisfaction	HEDIS	NCQA Quality Compass Medicaid National 50th percentile [All LOBs (Excluding PPOs and EPOs): Average] for the year prior to the measurement year

Identifier	Measure	Measure Description	Measure Steward	Federal Reporting Program	Target Population	Condition	Specification Source
HEDIS Measures							
CIS	Childhood Immunization Status	The percentage of children 2 years of age who had four diphtheria, tetanus and acellular pertussis (DTaP); three polio (IPV); one measles, mumps and rubella (MMR); three haemophilus influenza type B (HIB); three hepatitis B (HepB), one chicken pox (VZV); four pneumococcal conjugate (PCV); one hepatitis A (HepA); two or three rotavirus (RV); and two influenza (flu) vaccines by their second birthday. The measure calculates a rate for each vaccine and nine separate combination rates.	NCOA	CHIPRA, MU2	Children's Health	Prevention	HEDIS
IMA	Immunization Status for Adolescents	Percentage of adolescents that turned 13 years old during the measurement year and had specific vaccines by their 13th birthday. Report all individual vaccine numerators and combinations.	NCOA	CHIPRA	Children's Health	Prevention	HEDIS
WCC	Weight Assessment and Counseling for Nutrition and Physical Activity for Children/Adolescents; Body Mass Index Assessment for Children/Adolescents	Percentage of children ages 3 to 17 that had an outpatient visit with a primary care practitioner (PCP) or obstetrical/gynecological (OB/GYN) practitioner and whose weight is classified based on body mass index percentile for age and gender. The percentage of children ages 3 to 17 that had an outpatient visit with a primary care practitioner (PCP) or obstetrical/gynecological (OB/GYN) practitioner, with evidence of : <ul style="list-style-type: none"> • BMI percentile documentation • Counseling for nutrition • Counseling for physical activity 	NCOA	CHIPRA, MU2	Children's Health	Prevention	HEDIS

Attachment E

Identifier	Measure	Measure Description	Measure Steward	Federal Reporting Program	Target Population	Condition	Specification Source
SAA	Adherence to Antipsychotic Medications for Individuals with Schizophrenia	The measure calculates the percentage of individuals 19 years of age or greater as of the beginning of the measurement year with schizophrenia or schizoaffective disorder who are prescribed an antipsychotic medication, with adherence to the antipsychotic medication [defined as a Proportion of Days Covered (PDC)] of at least 0.8 during the measurement year (12 consecutive months).	NCOA	MEDICAID ADULT	Population Health	Behavioral Health	HEDIS
MPM	Annual Monitoring for Patients on Persistent Medications	The percentage of members 18 years of age and older who received at least 180 treatment days of ambulatory medication therapy for a select therapeutic agent during the measurement year and at least one therapeutic monitoring event for the therapeutic agent in the measurement year. For each product line, report each of the two rates separately and as a total rate.	NCOA	MEDICAID ADULT	Chronic Disease	Prevention	HEDIS
ABA	Adult BMI Assessment	The percentage of members 18-74 years of age who had an outpatient visit and whose body mass index (BMI) was documented during the measurement or the year prior to the measurement year.	NCOA	MEDICAID ADULT, CMS HEALTH HOMES	Population Health	Prevention	HEDIS
AMM	Antidepressant Medication Management	The percentage of members 18 years of age and older with a diagnosis of major depression and were newly treated with antidepressant medication, and who remained on an antidepressant medication treatment. Two rates are reported.	NCOA	MEDICAID ADULT, MU2	Population Health	Behavioral Health	HEDIS

Attachment E

Identifier	Measure	Measure Description	Measure Steward	Federal Reporting Program	Target Population	Condition	Specification Source
CCS	Cervical Cancer Screening	Percentage of women 21–64 years of age who were screened for cervical cancer: <ul style="list-style-type: none"> Women 21-64 who had cervical cytology performed every 3 years Women 30-64 who had cervical cytology/HPV co-testing performed every 5 years 	NCOA	MEDICAID ADULT, MU2	Population Health	Prevention	HEDIS
AMR	Asthma Medication Ratio	The percentage of patients 5–64 years of age who were identified as having persistent asthma and had a ratio of controller medications to total asthma medications of 0.50 or greater during the measurement year.	NCOA	MEDICAID	Population Health	Pulmonary/ Critical Care	HEDIS
FVA	Flu Vaccinations for Adults Ages 18 to 64	The percentage of adults 18 years of age and older who self-report receiving an influenza vaccine within the measurement period.	NCOA	MEDICAID ADULT	Population Health	Prevention	HEDIS/CAHPS
MSC	Medical Assistance With Smoking and Tobacco Use Cessation	Assesses different facets of providing medical assistance with smoking and tobacco use cessation. MCOs will report three components (questions): <ul style="list-style-type: none"> Advising Smokers and Tobacco Users to Quit Discussing Cessation Medications Discussing Cessation Strategies 	NCOA	MEDICAID ADULT	Population Health	Prevention	HEDIS/CAHPS
MMA	Medication Management for People with Asthma	The percentage of patients 5-64 years of age during the measurement year who were identified as having persistent asthma and were dispensed appropriate medications that they remained on during the treatment period. Two rates are reported.	NCOA	CHIPRA	Population Health	Pulmonary/ Critical Care	HEDIS

Attachment E

Identifier	Measure	Measure Description	Measure Steward	Federal Reporting Program	Target Population	Condition	Specification Source
CHL	Chlamydia Screening in Women	The percentage of women 16–24 years of age who were identified as sexually active and who had at least one test for Chlamydia during the measurement year.	NCQA	CHIPRA, MEDICAID ADULT	Population Health, Maternal Health	Perinatal and Reproductive Health, Sexually Transmitted Infectious Diseases	HEDIS
BCS	Breast Cancer Screening	Percentage of women 50-74 years of age who had a mammogram to screen for breast cancer.	NCQA	MEDICAID ADULT, MU2	Senior Care	Prevention	HEDIS
CAP	Child and Adolescents' Access to Primary Care Practitioners	Percentage of children ages 12 months – 19 years who had a visit with a PCP. The MCO reports four separate percentages: <ul style="list-style-type: none"> Children 12-24 months and 25 months – 6 years who had a visit with a PCP in the measurement year Children 7-11 years and adolescents 12-19 years who had a visit with a PCP in the measurement year or the year prior to the measurement year. 	NCQA	CHIPRA	Children's Health	Access/ Availability of Care	HEDIS
COL	Colorectal screening	The percentage of members 50-75 years of age who had appropriate screening for colorectal cancer.	NCQA	MEDICAID ADULT	Population Health	Prevention	HEDIS
SSD	Diabetes screening for people with Schizophrenia or Bipolar who are using Antipsychotic medications	The percentage of members 18-64 years of age with schizophrenia or bipolar disorder, who were dispensed an antipsychotic medication and had a diabetes screening test during the measurement year.	NCQA	MEDICAID ADULT	Population Health	Behavioral Health	HEDIS

Attachment E

Identifier	Measure	Measure Description	Measure Steward	Federal Reporting Program	Target Population	Condition	Specification Source
SPC	Statin Therapy for Patients with Cardiovascular Disease	<ul style="list-style-type: none"> The percentage of males 21-75 years of age and females 40-75 years of age during the measurement year, who were identified as having clinical atherosclerotic cardiovascular disease (ASCVD) and who received statin therapy (were dispensed at least one high or moderate-intensity statin medication during the measurement year.) The percentage of males 21-75 years of age and females 40-75 years of age during the measurement year, who were identified as having clinical atherosclerotic cardiovascular disease (ASCVD) and who had statin adherence of at least 80% (who remained on a high or moderate-intensity statin medication for at least 80% of the treatment period.) 	NCOA	MEDICAID ADULT	Population Health	Cardiovascular Care	HEDIS
CDC	Comprehensive Diabetes Care - HbA1c poor control (>9.0%)	The percentage of members 18-75 years of age with diabetes (type 1 and type 2) with HbA1c poor control (>9.0%).	NCOA	MEDICAID ADULT	Chronic Disease	Diabetes	HEDIS
CDC	Comprehensive Diabetes Care - HbA1c control (<8.0%)	The percentage of members 18-75 years of age with diabetes (type 1 and type 2) with HbA1c control (<8.0%).	NCOA	MEDICAID ADULT	Chronic Disease	Diabetes	HEDIS
CDC	Comprehensive Diabetes Care - BP control (<140/90 mm Hg).	The percentage of members 18-75 years of age with diabetes (type 1 and type 2) with BP control (<140/90 mm Hg).	NCOA	MEDICAID ADULT	Chronic Disease	Diabetes	HEDIS
PCR	Plan All-Cause Readmissions	For members 18-64 years of age, the risk-adjusted rate of acute inpatient stays during the measurement year that were followed by an unplanned acute readmission for any diagnosis within 30 days.	NCOA	MEDICAID ADULT	Population Health	All Cause Readmissions	HEDIS

Identifier	Measure	Measure Description	Measure Steward	Federal Reporting Program	Target Population	Condition	Specification Source
AAP	Adults' Access to Preventive/ Ambulatory Health Services	The percentage of members age 20 years and older who had an ambulatory or preventive care visit during the measurement year. Three age stratifications and a total rate are reported: <ul style="list-style-type: none"> • 20-44 years • 45-64 years • 65 years and older • Total 	NCQA	MEDICAID ADULT	Population Health	Prevention	HEDIS
FUH	Follow-Up After Hospitalization for Mental Illness - Within 7 days of discharge	The percentage of discharges for members 6 years of age and older who were hospitalized for treatment of selected mental illness diagnoses and who had a follow-up visit with a mental health practitioner within 7 days of discharge.	NCQA	CHIPRA	Behavioral Health	Behavioral Health	HEDIS
AMB	Ambulatory Care- Outpatient Visits	This measure summarizes utilization of ambulatory care Outpatient Visits per 1,000 member months.	NCQA	MEDICAID	Population Health	Utilization	HEDIS
PQI Measures							
PQI01	Diabetes Short Term Complications Admission Rate	Number of discharges for diabetes short term complications per 100,000 member months per Medicaid enrollees age 18 and older.	AHRQ	MEDICAID ADULT	Chronic Disease	Diabetes	Section V
PQI05	COPD and Asthma in Older Adults Admission Rate	This measure is used to assess the number of admissions for chronic obstructive pulmonary disease (COPD) per 100,000 population. The number of discharges for chronic obstructive pulmonary disease (COPD) or asthma per 100,000 member months for Medicaid enrollees age 40 and older.	AHRQ	MEDICAID ADULT	Population Health	Pulmonary/ Critical Care	Section V

Identifier	Measure	Measure Description	Measure Steward	Federal Reporting Program	Target Population	Condition	Specification Source
PQI08	Heart Failure Admission Rate	Percent of population with an admissions for heart failure (reported by Recipient Parish). The number of discharges for heart failure per 100,000 member months for Medicaid enrollees age 18 and older (reported by Recipient Parish).	AHRQ	MEDICAID ADULT	Chronic Disease	Cardiovascular Care	Section V
PQI15	Asthma in Younger Adults Admission Rate	Admissions for a principal diagnosis of asthma per 100,000 population, ages 18 to 39 years. Excludes admissions with an indication of cystic fibrosis or anomalies of the respiratory system, obstetric admissions, and transfers from other institutions. Number of discharges for asthma per 100,000 member months for Medicaid enrollees ages 18 to 39.	AHRQ	MEDICAID ADULT	Population Health	Pulmonary/ Critical Care	Section V
Vital Record Measures							
LBW	Percentage of low birth weight births	Percentage of live births that weighted less than 2,500 grams in the state during the reporting period.	CDC	CHIPRA, HRSA	Children's and Maternal Health	Perinatal and Reproductive Health	Section V
NQF (PC-01)	Elective Delivery	This measure assesses patients with elective vaginal deliveries or elective cesarean sections at >= 37 and < 39 weeks of gestation completed	TJC	MEDICAID ADULT, MU2	Maternal Health	Perinatal and Reproductive Health	Section V
CMS Measures							
HIV	HIV Viral Load Suppression	Percentage of patients, regardless of age, with a diagnosis of HIV with a HIV viral load less than 200.	HRSA HIV/AIDS Bureau	MEDICAID ADULT	Chronic Disease	HIV	Section V
CCP-CH	Contraceptive Care-Postpartum (ages 15-20)	The percentage of women ages 15-20 who had a live birth and were provided a most or moderately effective method of contraception within 3 and 60 days of delivery. Four rates are reported.	CMS	CHIPRA	Maternal Health	Perinatal and Reproductive Health	OPA

Attachment E

Identifier	Measure	Measure Description	Measure Steward	Federal Reporting Program	Target Population	Condition	Specification Source
CCP-AD	Contraceptive Care-Postpartum (ages 21-44)	The percentage of women ages 21-44 who had a live birth and were provided a most or moderately effective method of contraception within 3 and 60 days of delivery. Four rates are reported.	CMS	MEDICAID ADULT	Maternal Health	Perinatal and Reproductive Health	OPA
NSV	Cesarean Rate for Low-Risk First Birth Women	The percentage of cesareans in live births at or beyond 37.0 weeks gestation to women that are having their first delivery and are singleton (no twins or beyond) and are vertex presentation (no breech or transverse positions).	TJC	CHIPRA	Children's and Maternal Health	Perinatal and Reproductive Health	Section V