



Infectious Disease Epidemiology Section
Office of Public Health
Louisiana Dept of Health & Hospitals
800-256-2748 (24 hr. number)
www.infectiousdisease.dhh.louisiana.gov

Information on Meningitis Public Information

What is meningitis?

Meningitis is an infection of the fluid of a person's spinal cord and the fluid that surrounds the brain. People sometimes refer to it as spinal meningitis. Meningitis is usually caused by a viral or bacterial infection. Knowing whether meningitis is caused by a virus or bacterium is important because the severity of illness and the treatment differ.

Viral meningitis is generally less severe and resolves without specific treatment, while **bacterial meningitis** can be quite severe and may result in brain damage, hearing loss, or learning disability.

For **bacterial meningitis**, it is also important to know which type of bacteria is causing the meningitis because antibiotics can prevent some types from spreading and infecting other people. Before the 1990s, *Haemophilus influenzae* type b (Hib) was the leading cause of bacterial meningitis, but new vaccines being given to all children as part of their routine immunizations have reduced the occurrence of invasive disease due to *H. influenzae*. Today, *Streptococcus pneumoniae* and *Neisseria meningitidis* are the leading causes of bacterial meningitis.

What are the signs and symptoms of meningitis?

Common symptoms of meningitis in anyone over the age of 2 years are:

- high fever
- headache
- stiff neck

These symptoms can develop over several hours, or they may take 1 to 2 days.

Other symptoms may include:

- nausea, vomiting
- discomfort looking into bright lights
- confusion
- sleepiness

In newborns and small infants, the classic symptoms of fever, headache, and neck stiffness may be absent or difficult to detect and the infant:

- may only appear slow or inactive
- be irritable
- have vomiting
- be feeding poorly.

As the disease progresses, patients of any age may have seizures.

How is meningitis diagnosed?

Early diagnosis and treatment are very important. If symptoms occur, the patient should see a doctor immediately. The diagnosis is usually made by growing bacteria from a sample of spinal fluid. The spinal fluid is obtained by performing a spinal tap, in which a needle is inserted into an area in the lower back where fluid in the spinal canal is readily accessible. Identification of the type of bacteria responsible is important for selection of correct antibiotics.

Is meningitis contagious?

Yes, some forms are bacterial meningitis are contagious. The bacteria are spread through the exchange of respiratory and throat secretions (i.e., coughing, kissing). Fortunately, none of the bacteria that cause meningitis are as contagious as things like the common cold or the flu, and they are not spread by casual contact or by simply breathing the air where a person with meningitis has been.

However, sometimes the bacteria that cause meningitis have spread to other people who have had close or prolonged contact with a patient with meningitis caused by *Neisseria meningitidis* (also called meningococcal meningitis) or Hib. People in the same household or day-care center, or anyone with direct contact with a patient's oral secretions (such as a boyfriend or girlfriend) would be considered at increased risk of acquiring the infection. People who qualify as close contacts of a person with meningitis caused by *N. meningitidis* should receive antibiotics to prevent them from getting the disease. Antibiotics for contacts of a person with Hib meningitis disease are no longer recommended if all contacts 4 years of age or younger are fully vaccinated against Hib disease (see below).

Are there vaccines against meningitis?

Yes, there are vaccines against Hib and against some strains of *N. meningitidis* and many types of *Streptococcus pneumoniae*. The vaccines against Hib are very safe and highly effective.

There is also a vaccine that protects against four strains of *N. meningitidis*, but it is not routinely used in the United States and is not effective in children under 18 months of age. The vaccine against *N. meningitidis* is sometimes used to control outbreaks of some types of meningococcal meningitis in the United States. Meningitis cases should be reported to state or local health departments to assure follow-up of close contacts and recognize outbreaks. Although large epidemics of meningococcal meningitis do not occur in the United States, some countries experience large, periodic epidemics. Overseas travelers should check to see if meningococcal vaccine is recommended for their destination. Travelers should receive the vaccine at least one week before departure, if possible.