

Clinical Policy: Deferoxamine (Desferal)

Reference Number: LA.PHAR.146

Effective Date: 09.15.22

Last Review Date: ~~05.09.24~~ 09.18.24

Line of Business: Medicaid

[Coding Implications](#)

[Revision Log](#)

See [Important Reminder](#) at the end of this policy for important regulatory and legal information.

****Please note: This policy is for medical benefit****

Description

Deferoxamine (Desferal®) is an iron-chelating agent.

FDA Approved Indication(s)

Desferal is indicated:

- As an adjunct to standard measures for the treatment of acute iron intoxication.
- For the treatment of transfusional iron overload in patients with chronic anemia.

Limitation(s) of use: Desferal is not indicated for the treatment of primary hemochromatosis (since phlebotomy is the method of choice for removing excess iron in this disorder).

Policy/Criteria

Provider must submit documentation (such as office chart notes, lab results or other clinical information) supporting that member has met all approval criteria.

It is the policy of Louisiana Healthcare Connections that ~~Desferal~~ deferoxamine is **medically necessary** when the following criteria are met:

I. Initial Approval Criteria

A. Acute Iron Intoxication (must meet all):

1. Diagnosis of acute iron intoxication;
2. If request is for brand Desferal, member must use generic deferoxamine, unless contraindicated or clinically significant adverse effects are experienced;
3. Dose does not exceed 6,000 mg in 24 hours (IM or IV).

Approval duration: 1 month

B. Chronic Iron Overload due to Transfusion-Dependent Anemias

1. Diagnosis of chronic iron overload due to transfusion-dependent anemia (e.g., congenital/acquired anemias including thalassemia, sickle cell anemia, aplastic anemia, myelodysplasia);
2. Transfusion history of ≥ 100 mL/kg of packed red blood cells (e.g., ≥ 20 units of packed red blood cells for a 40 kg person);
3. Serum ferritin level $> 1,000$ mcg/L;
4. If request is for brand Desferal, member must use generic deferoxamine, unless contraindicated or clinically significant adverse effects are experienced;

CLINICAL POLICY

Deferoxamine



5. Therapy does not include concurrent use of other iron chelators, unless member has excess cardiac iron as evidence by cardiac T2* < 20 millisecond or iron-induced cardiomyopathy;
6. Dose does not exceed any of the following (a, b, or c):
 - a. SC: 2,000 mg per day;
 - b. IV: 40 mg/kg per day for children; 60 mg/kg per day for adults;
 - c. IM: 1,000 mg per day.

Approval duration: 6 months

C. Other diagnoses/indications (must meet 1 or 2):

1. If this drug has recently (within the last 6 months) undergone a label change (e.g., newly approved indication, age expansion, new dosing regimen) that is not yet reflected in this policy, refer to LA.PMN.255
2. If the requested use (e.g., diagnosis, age, dosing regimen) is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized) AND criterion 1 above does not apply, refer to the off-label use policy LA.PMN.53

II. Continued Therapy

A. Acute Iron Intoxication

1. Re-authorization is not permitted. Members must meet initial approval criteria for new cases of acute iron intoxication.

Approval duration: Not applicable

B. Chronic Iron Overload due to Transfusion-Dependent Anemias (must meet all):

1. Currently receiving medication via Louisiana Healthcare Connections benefit or member has previously met initial approval criteria;
2. Member is responding positively to therapy as evidenced by a decrease in serum ferritin levels as compared to pretreatment baseline;
3. Current documentation (within the last 30 days) shows a serum ferritin level ≥ 500 mcg/L;
4. If request is for brand Desferal, member must use generic deferoxamine, unless contraindicated or clinically significant adverse effects are experienced;
5. Therapy does not include concurrent use of other iron chelators, unless member has excess cardiac iron as evidence by cardiac T2* < 20 millisecond or iron-induced cardiomyopathy;
6. If request is for a dose increase, new dose does not exceed any of the following (a, b, or c):
 - a. SC: 2,000 mg per day;
 - b. IV: 40 mg/kg per day for children; 60 mg/kg per day for adults;
 - c. IM: 1,000 mg per day.

Approval duration: 12 months

C. Other diagnoses/indications (must meet 1 or 2):

1. If this drug has recently (within the last 6 months) undergone a label change (e.g., newly approved indication, age expansion, new dosing regimen) that is not yet reflected in this policy, refer to LA.PMN.255

CLINICAL POLICY

Deferoxamine



2. If the requested use (e.g., diagnosis, age, dosing regimen) is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized) AND criterion 1 above does not apply, refer to the off-label use policy LA.PMN.53

III. Diagnoses/Indications for which coverage is NOT authorized:

- A. Non-FDA approved indications, which are not addressed in this policy, unless there is sufficient documentation of efficacy and safety according to the off-label use policy LA.PMN.53
- B. Primary hemochromatosis;
- C. Parkinson's disease.

IV. Appendices/General Information

Appendix A: Abbreviation/Acronym Key

FDA: Food and Drug Administration

Appendix B: Therapeutic Alternatives

Not applicable

Appendix C: Contraindications/Boxed Warnings

- Contraindication(s):
 - Known hypersensitivity to the active substance
 - Severe renal disease or anuria, since the drug and the iron chelate are excreted primarily by the kidney
- Boxed warning(s): none reported

Appendix D: General Information

- In FAIRPARK-II, deferiprone, an iron chelator, was associated with worse scores in measures of parkinsonism compared to placebo over a 36-week period in participants with newly diagnosed Parkinson's disease who had never received levodopa.

V. Dosage and Administration

Indication	Dosing Regimen	Maximum Dose
Acute iron intoxication	1,000 mg x 1 dose, then 500 mg Q4-12 hr PRN* <i>*IM route if patient not in shock; IV infusion limited to patients in cardiovascular collapse.</i>	6,000 mg/24 hr
Chronic iron overload	Average daily dose between 20-60 mg/kg SC infusion QD	See dosing regimen
	20-40 mg/kg IV daily (children*) and 40-50 mg/kg IV daily (adults) for 5-7 days per week <i>*Maximum recommended daily dose is 40 mg/kg/day until growth (body weight and linear growth) has ceased.</i>	40 mg/kg/day (children) 60 mg/kg/day (adults)
	500-1,000 mg IM/day	1,000 mg/day

VI. Product Availability

CLINICAL POLICY

Deferoxamine



Single-dose vial of lyophilized deferoxamine mesylate: 500 mg, 2 g

VII. References

1. Desferal Prescribing Information. East Hanover, NJ: Novartis Pharmaceuticals Corporation; September 2022. Available at: https://www.accessdata.fda.gov/drugsatfda_docs/label/2022/016267s0621b1novartis.com/us-en/sites/novartis_us/files/desferal.pdf. Accessed April 14, 2023May 23, 2024.
2. Musallam KM, Angastiniotis M, Eleftheriou A, Porter JB. Cross talk between available guidelines for the management of patients with beta-thalassemia major. *Acta Haematol*. 2013; 130: 64-73. DOI: 10.1159/000345734.
3. Hoffbrand AV, Taher A, Cappellini MD. How I treat transfusional iron overload. *Blood*. November 1, 2012; 120(18): 3657-3669.
2. Taher A, Musallam K, Cappellini MD. Guidelines for the management of non-transfusion dependent thalassaemia (NTDT) 2nd edition. Thalassaemia International Federation. 2018. Available at: <https://thalassaemia.org.cy/publications/tif-publications/guidelines-for-the-clinical-management-of-non-transfusion-dependent-thalassaemias-updated-version/>. Accessed May 22, 2024.
3. Taher A, Musallam K, Cappellini MD. Guidelines for the management of non-transfusion dependent β -thalassaemia 3rd edition. Thalassaemia International Federation. 2023. Available at: <https://thalassaemia.org.cy/publications/tif-publications/guidelines-for-the-management-of-non-transfusion-dependent-%ce%b2-thalassaemia-3rd-edition-2023/>. Accessed May 22, 2024.
4. Amid A, Lal A, Coates TD, Fucharoen S, et al. Guidelines for the management of α -thalassaemia. Thalassaemia International Federation. 2023. Available at: <https://thalassaemia.org.cy/publications/tif-publications/guidelines-for-the-management-of-%ce%b1-thalassaemia/?-thalassaemia%2F>. Accessed May 22, 2024.
- 4.5. Cappellini MD, Farmakis D, Porter J, et al. 2021 Guidelines for the management of transfusion dependent thalassemia (TDT) 4th edition. Thalassaemia International Federation. 2021. Available at: <https://thalassaemia.org.cy/publications/tif-publications/guidelines-for-the-management-of-transfusion-dependent-thalassaemia-4th-edition-2021/>. Accessed May 4, 202223, 2024.
- 5.6. Devos D, Labreuche J, Rascol O, et al. Trial of deferiprone in Parkinson's disease. *N Engl J Med* 2022; 387:2045-2055.
6. Children's Hospital & Research Center Oakland. 2012 Standards of Care Guidelines for Thalassemia. Available at: <https://thalassemia.com/documents/SOCGuidelines2012.pdf>. Accessed May 4, 2023.
7. Sheth S. Strategies for managing transfusional iron overload: conventional treatments and novel strategies. *Curr Opin Hematol*. 2019 May; 26(3): 139-144.

Coding Implications

Codes referenced in this clinical policy are for informational purposes only. Inclusion or exclusion of any codes does not guarantee coverage. Providers should reference the most up-to-date sources of professional coding guidance prior to the submission of claims for reimbursement of covered services.

Formatted: Font: Italic

CLINICAL POLICY
Deferoxamine



HCPCS Codes	Description
J0895	Injection, deferoxamine mesylate, 500 mg

Reviews, Revisions, and Approvals	Date	LDH Approval Date
Converted corporate to local policy. Template changes applied to other diagnoses/indications and continued therapy section.	09.22	09.15.22
Template changes applied to other diagnoses/indications and continued therapy section. Added Parkinson disease to section III with rationale in Appendix D. Added verbiage this policy is for medical benefit only.	06.02.23	10.05.23
Annual review: updated FDA approved indications per prescribing information; per competitor analysis for continuation of therapy in chronic iron overload added requirement that member is responding positively to therapy as evidenced by a decrease in serum ferritin levels as compared to pretreatment baseline; for chronic iron overload added requirement that therapy does not include concurrent use of other iron chelators, unless member has excess cardiac iron as evidence by cardiac T2* < 20 millisecond or iron-induced cardiomyopathy; references reviewed and updated.	05.09.24	<u>07.29.24</u>
<u>In Policy/Criteria, clarified policy is medically necessary for all deferoxamine products not only Desferal; references reviewed and updated.</u>	<u>09.19.24</u>	

Important Reminder

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program approval status; evidence-based guidelines and positions of leading national health professional organizations; views of physicians practicing in relevant clinical areas affected by this clinical policy; and other available clinical information. LHCC makes no representations and accepts no liability with respect to the content of any external information used or relied upon in developing this clinical policy. This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved.

The purpose of this clinical policy is to provide a guide to medical necessity, which is a component of the guidelines used to assist in making coverage decisions and administering benefits. It does not constitute a contract or guarantee regarding payment or results. Coverage decisions and the administration of benefits are subject to all terms, conditions, exclusions and limitations of the coverage documents (e.g., evidence of coverage, certificate of coverage, policy, contract of insurance, etc.), as well as to state and federal requirements and applicable LHCC administrative policies and procedures.

CLINICAL POLICY

Deferoxamine



This clinical policy is effective as of the date determined by LHCC. The date of posting may not be the effective date of this clinical policy. This clinical policy may be subject to applicable legal and regulatory requirements relating to provider notification. If there is a discrepancy between the effective date of this clinical policy and any applicable legal or regulatory requirement, the requirements of law and regulation shall govern. LHCC retains the right to change, amend or withdraw this clinical policy, and additional clinical policies may be developed and adopted as needed, at any time.

This clinical policy does not constitute medical advice, medical treatment or medical care. It is not intended to dictate to providers how to practice medicine. Providers are expected to exercise professional medical judgment in providing the most appropriate care, and are solely responsible for the medical advice and treatment of members. This clinical policy is not intended to recommend treatment for members. Members should consult with their treating physician in connection with diagnosis and treatment decisions.

Providers referred to in this clinical policy are independent contractors who exercise independent judgment and over whom LHCC has no control or right of control. Providers are not agents or employees of LHCC.

This clinical policy is the property of LHCC. Unauthorized copying, use, and distribution of this clinical policy or any information contained herein are strictly prohibited. Providers, members and their representatives are bound to the terms and conditions expressed herein through the terms of their contracts. Where no such contract exists, providers, members and their representatives agree to be bound by such terms and conditions by providing services to members and/or submitting claims for payment for such services.

©2024 Louisiana Healthcare Connections. All rights reserved. All materials are exclusively owned by Louisiana Healthcare Connections and are protected by United States copyright law and international copyright law. No part of this publication may be reproduced, copied, modified, distributed, displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise published without the prior written permission of Louisiana Healthcare Connections. You may not alter or remove any trademark, copyright or other notice contained herein. Louisiana Healthcare Connections is a registered trademark exclusively owned by Louisiana Healthcare Connections.