

Infectious Disease: Respiratory Lab Testing

Reference Number: LA.CP.CG.27

[Coding implications](#)

Date of Last Revision [9/24/2025](#)

[Revision Log](#)

OVERVIEW

Respiratory illnesses cause significant morbidity and mortality within the United States and around the world. Seasonal influenza, respiratory syncytial virus (RSV), and SARS-CoV-2 infect many individuals each year, and while most will recover with no complications, a significant number will be hospitalized or die. Diagnostic testing for upper respiratory tract infections can be very useful for clinicians, as clinical signs and symptoms of these infections can have significant overlap between pathogens. Accurate and rapid testing techniques may aid clinicians, via identification of a specific pathogen, in selecting the best course of treatment for patients. Optimally, treatment is started within 48-72 hours of diagnosis. Testing methods range from culture and microscopy to immunoassays and advanced molecular diagnostic techniques; technology in this space is evolving rapidly and clinical guidelines can lag as a result.

This policy is intended for use in the outpatient setting.

POLICY REFERENCE TABLE

Criteria Sections	Example Tests (Labs)	References
Syndromic/Multiple x Respiratory Panels with 6 or More Targets	Respiratory Pathogen Panel, Quest Diagnostics	3
	ePlex Respiratory Pathogen Panel (GenMark Diagnostics, Inc)	
	Biofire FilmArray Respiratory Panel 2.1 (Biofire Diagnostics)	
	QIAstat-Dx Respiratory SARS-CoV-2 Panel (QIAGEN Sciences)	
	ePlex Respiratory Pathogen Panel 2 (GenMark Diagnostics, Inc)	

CLINICAL POLICY

Infectious Disease: Respiratory Lab Testing

	Respiratory Pathogen with ABR (RPX) (Lab Genomics LLC, Thermofisher Scientific)	
	Respiratory Virus PCR Panel IV (Quest Diagnostics)	
	Respiratory Viral Panel, PCR (Quest Diagnostics)	
SARS-CoV-2, RSV, or Influenza A/B, OR Multiplex Respiratory Viral Panels with 5 or Fewer Targets	Xpert Xpress SARS-CoV-2/Flu/RSV for SARS-CoV-2 and Flu targets only (Cepheid)	3, 6, 7
	Xpert Xpress SARS-CoV-2/Flu/RSV for all targets (Cepheid)	
	Infectious Agent Antigen Detection by Immunoassay	
	Infectious Agent Antigen Detection by Immunoassay, Qualitative or Semiquantitative	
	Infectious Agent Antigen Detection by Immunoassay, Qualitative or Semiquantitative, SARS-CoV-2 and Flu A/B	
	Influenza A and B and RSV RNA, Qualitative, Real-Time RT-PCR (Quest Diagnostics)	

CLINICAL POLICY

Infectious Disease: Respiratory Lab Testing



	SARS-CoV-2 RNA (COVID-19), Qualitative NAAT (Quest Diagnostics)	
	SARS-CoV-2 RNA (COVID-19) and Influenza A and B, Qualitative NAAT (Quest Diagnostics)	
	Infectious Agent Antigen Detection by Nucleic Acid (DNA or RNA) SARS-CoV-2/Flu/RSV Multiplex Amplified Probe Technique	
	Infectious Agent Antigen Detection by Immunoassay with Direct Optical Observation	
Bacterial Respiratory Infection/Pneumonia Panels	Infectious Agent: Chlamydia pneumoniae Detection by Nucleic Acid (DNA or RNA), Direct Probe Technique	3
	Chlamydophila pneumoniae, DNA, Qualitative, Real-Time PCR (Quest Diagnostics)	
	Infectious Agent: Chlamydia pneumoniae Detection by Nucleic Acid (DNA or RNA), Quantification	
	Legionella DNA, Qualitative, Real-Time PCR (Quest Diagnostics)	
	Infectious Agent: Mycoplasma pneumoniae Detection by Nucleic Acid (DNA or RNA), Direct Probe Technique	

CLINICAL POLICY

Infectious Disease: Respiratory Lab Testing

	Mycoplasma pneumoniae, DNA PCR (Labcorp)	
	Infectious Agent: Mycoplasma pneumonia Detection by Nucleic Acid (DNA or RNA), Quantification	
Influenza A and B Antibody Tests	Influenza Type A and Type B Antibody, Serum (Quest Diagnostics)	1
Group A Streptococcus Pharyngitis Tests	Streptococcus Group A Antigen Detection by Immunoassay	2
	Streptococcus Group A Antigen Detection by Nucleic Acid Direct Probe Technique	
	Group A Streptococcus Detection, NAA (Labcorp)	
	Streptococcus Group A Antigen, Adult (Quest Diagnostics)	
Group A Streptococcus Pharyngitis Cultures	Streptococcus Group A Culture (Quest Diagnostics)	2, 4
Group A Streptococcus Antibody Tests	Antistreptolysin O (ASO) Antibodies (Labcorp)	2

CRITERIA

It is the policy of Louisiana Healthcare Connections that the specific tests noted below are **medically necessary** when meeting the related criteria:

RESPIRATORY PATHOGEN PANEL TESTS

Syndromic/Multiplex Respiratory Panels with 6 or More Targets

- I. Syndromic Multiplex Respiratory Panels with ~~6~~six or more targets ~~may be, when performed in the outpatient setting, are~~ considered medically necessary ~~when for~~ member/enrollees meeting any of the following:
 - A. ~~The member/enrollee presents in the outpatient setting with signs or symptoms of an acute respiratory infection, AND~~
 1. ~~The member/enrollee meets at least one of the following criteria:~~
 - a) ~~Immunocompromised, OR~~
 - b) ~~Has severe pneumonia, OR~~
 - c) ~~Has exacerbations of airway disease, AND~~
 - B. ~~Results of the testing will influence the member's/enrollee's clinical management.~~
 1. With serious or critical illness, OR
 2. At imminent risk of becoming seriously or critically ill, OR
 3. With immunodeficiency, AND/OR
 4. With a severe underlying condition.
- II. Current evidence does not support the use of Syndromic Multiplex Respiratory Panels with 6 or more targets for all other indications.

SARS-CoV-2, RSV, or Influenza A/B, OR Multiplex Respiratory Viral Panels with 5 or Fewer Targets

- I. SARS-CoV-2, RSV, or Influenza A/B, **OR** Multiplex Respiratory Viral Panels with 5 or fewer targets, ~~may be~~other than CPT 87631 which is out of the scope of this policy, are considered **medically necessary** when:

CLINICAL POLICY

Infectious Disease: Respiratory Lab Testing

- A. The member/enrollee presents in the outpatient setting with [signs or symptoms of an acute respiratory infection](#), **AND**
 - B. Results of the testing will influence the member's/enrollee's clinical management.
- II. Current evidence does not support the use of SARS-CoV-2, RSV, or Influenza A/B, **OR** Multiplex Respiratory Viral Panels with 5 or fewer targets, for all other indications.

Bacterial Respiratory Infection/Pneumonia Panels

- I. Bacterial Respiratory Infection/Pneumonia Panels ~~may be~~ considered **medically necessary** when:
 - A. The member/enrollee presents in the outpatient setting with [signs or symptoms of an acute respiratory infection](#), **AND**
 - B. The member/enrollee meets any of the following criteria:
 - 1. New or worsening lung infiltrates, **OR**
 - 2. [Moderate to severe upper respiratory illness](#), **OR**
 - 3. Has received empiric antibiotics before obtaining cultures, **OR**
 - 4. Has possible multidrug-resistant bacteria or polymicrobial infection, **AND**
 - C. Results of the testing will influence the member's clinical management.
- II. Current evidence does not support the use of Bacterial Respiratory Infection/Pneumonia Panels for all other indications.

Influenza A and B Antibody Tests

- I. Current evidence does not support the use of Influenza A and B Antibody Tests for the purpose of diagnosing influenza.

Group A Streptococcus Pharyngitis Tests

- I. Group A Streptococcus Pharyngitis Tests ~~may be~~ considered **medically necessary** when:
 - A. The member/enrollee presents in the outpatient setting with at least one of the following:
 - 1. Acute pharyngitis, **OR**

CLINICAL POLICY

Infectious Disease: Respiratory Lab Testing

2. Fever, **OR**
 3. Tonsillopharyngeal inflammation, **OR**
 4. Patchy tonsillopharyngeal exudates, **OR**
 5. Palatal petechiae, **OR**
 6. Anterior cervical lymphadenitis, **OR**
 7. Scarletiform rash, **AND**
- B. The member/enrollee does **NOT** have clinical and epidemiological features that strongly suggest a viral etiology (e.g., cough, rhinorrhea, hoarseness, and oral ulcers), **AND**
- C. Results of the testing will influence the member's/enrollee's clinical management.
- II. Current evidence does not support the use of Group A Streptococcus Pharyngitis Tests for all other indications.

Group A Streptococcus Pharyngitis Cultures

- I. Group A Streptococcus Pharyngitis Culture ~~may be~~ considered **medically necessary** when:
 - A. The member/enrollee is between the ages of 3 years and ~~14~~18 years, **AND**
 - B. The member/enrollee had a negative group A Streptococcus rapid antigen detection test (RADT), **AND**
 - C. The member/enrollee presents in the outpatient setting with at least one of the following:
 1. Acute pharyngitis, **OR**
 2. Fever, **OR**
 3. Tonsillopharyngeal inflammation, **OR**
 4. Patchy tonsillopharyngeal exudates, **OR**
 5. Palatal petechiae, **OR**
 6. Anterior cervical lymphadenitis, **OR**
 7. Scarletiform rash, **AND**

CLINICAL POLICY

Infectious Disease: Respiratory Lab Testing

- D. The member/enrollee does **NOT** have clinical and epidemiological features that strongly suggest a viral etiology (e.g., cough, rhinorrhea, hoarseness, and oral ulcers), **AND**
 - E. Results of the testing will influence the member's/enrollee's clinical management.
- II. Current evidence does not support the use of Group A Streptococcus Pharyngitis Culture for all other indications.

Group A Streptococcus Antibody Tests

- I. Current evidence does not support the use of Group A Streptococcus Antibody Tests for the purpose of evaluating a member/enrollee with acute pharyngitis for a possible group A streptococcus infection.

Code Specific Guidelines

~~I. Coverage of Respiratory Viral Panels (87631, 87632 and 87633)~~

- ~~II. CPT code 87631 is deemed medically necessary in the following instances:~~
- ~~a. Infants receiving monthly RSV prophylaxis with palivizumab because of high-risk conditions such as prematurity, respiratory disease, or cardiac disease.~~
 - ~~b. Long-term care facility residents returning to a facility, or a person of any age returning to a congregate setting.~~

~~PLEASE NOTE: A primary care physician may perform this 3-5 panel test if medically necessary.~~

- ~~III. CPT codes 87632 and 87633 are deemed potentially medically necessary only for:~~
- ~~a. Beneficiaries with serious or critical illness or at imminent risk of becoming seriously or critically ill, immunodeficiency, and/or severe underlying condition contributory to testing using an expanded syndromic panel.~~

- ~~III. Testing is approved for the following places of service (POS):~~
- ~~a. Places of service (POS) 19—off-campus outpatient hospital, 21—inpatient hospital, 22~~
 - ~~b. on-campus outpatient hospital, 23—emergency room.~~

~~PLEASE NOTE: Tests should be ordered as follows (for healthcare POS other than those listed in the above bullet):~~

- ~~IIII. Testing for these services should only occur in accordance with one or more of the following instances:~~
- ~~a. For immune-competent beneficiaries, the test must be ordered by an infectious disease specialist or pulmonologist who is diagnosing and treating the beneficiary.~~

CLINICAL POLICY

Infectious Disease: Respiratory Lab Testing

- ~~b. For immune-compromised beneficiaries, the test must be ordered by a clinician specialist in one of the following: infectious diseases, oncology, transplant (for any panel), or pulmonologist who is diagnosing and treating the beneficiary.~~

~~PLEASE NOTE: Regarding the previous two bullets, an exception may be made within geographic locations where the specialist(s) cannot be reasonably reached by the beneficiary; AND the beneficiary is under the care of one of these providers: infectious diseases, oncology, transplant (for any panel), or pulmonologist; AND the ordering provider is located closer to the beneficiary's place of residence than the nearest specialist.~~

~~This exception is intended for beneficiaries living in rural locations with limited clinical specialist access only.~~

NOTES AND DEFINITIONS

1. **Moderate to severe upper upper respiratory illness** includes one or more clinical findings of lower respiratory illness (e.g., pneumonia, severe cough/bronchitis, shortness of breath, difficulty breathing).
2. **Severe pneumonia** is defined by the Infectious Diseases Society of America/American Thoracic Society Criteria as: the presence of one major criterion or at least three minor criteria.

Minor criteria: respiratory rate ≥ 30 breaths/min, PaO₂/FiO₂ ratio ≤ 250 , multilobar infiltrates, confusion/disorientation, uremia (blood urea nitrogen level ≥ 20 mg/dl), leukopenia (white blood cell count $< 4,000$ cells/ μ l), thrombocytopenia (platelet count $< 100,000$ / μ l), hypothermia (core temperature $< 36^{\circ}\text{C}$), and hypotension requiring aggressive fluid resuscitation.

Major criteria: septic shock with need for vasopressors and respiratory failure requiring mechanical ventilation.

3. **Airway disease** is a nonspecific clinical term for a heterogeneous group of conditions including chronic obstructive pulmonary disease (COPD), emphysema, cystic fibrosis, asthma, and bronchiectasis.
4. **Signs and symptoms of acute respiratory infection** include upper or lower respiratory tract symptoms (cough, runny nose, sore throat, bronchitis, pneumonia, bronchiolitis), with or without fever, influenza-like illness (ILI) (fever and either cough or sore throat), and respiratory distress (difficulty in breathing; often characterized by increased respiratory rate and use of accessory muscles of breathing).

CLINICAL POLICY

Infectious Disease: Respiratory Lab Testing

BACKGROUND AND RATIONALE

Syndromic/Multiplex Respiratory Panels with 6 or More Targets

Infectious Diseases Society of America

The IDSA published clinical and diagnostic recommendations in 2020 regarding molecular testing for acute respiratory tract infections (RTIs). These recommendations state the following:

“Multiplex viral NAAT [nucleic acid amplification tests] (potentially combined with bacterial NAAT) also make clinical sense for immunocompromised and critically ill patients with pneumonia as well as for those with exacerbations of airway disease.” (p. 2748).

SARS-CoV-2, RSV, or Influenza A/B, OR Multiplex Respiratory Viral Panels with 5 or Fewer Targets

Infectious Diseases Society of America

The IDSA published clinical and diagnostic recommendations in 2020 regarding molecular testing for acute respiratory tract infections (RTIs). These recommendations state the following:

“Molecular testing for multiple respiratory viruses simultaneously may also be more cost-effective than traditional antigen- or culture-based methods from a laboratory perspective, especially given certain thresholds of disease prevalence.” (p. 2744)

Centers for Disease Control and Prevention

The CDC states the following on their website discussing RSV: “Healthcare providers should consider RSV in patients with respiratory illness, particularly during the RSV season.”

The CDC states the following on their website discussing COVID-19: “Key times to get tested: if you have symptoms, test immediately.”

Bacterial Respiratory Infection/Pneumonia Panels

Infectious Diseases Society of America

The IDSA published clinical and diagnostic recommendations in 2020 regarding molecular testing for acute respiratory tract infections (RTIs). These recommendations state the following:

“...bacterial NAAT may prove most useful in situations where patients have new or worsening lung infiltrates, are moderately to severely ill, have received empiric antibiotics before obtaining

CLINICAL POLICY

Infectious Disease: Respiratory Lab Testing

cultures, and/or there is concern for multidrug-resistant bacteria or a polymicrobial infection.” (p. 2747)

Influenza A and B Antibody Tests

Infectious Diseases Society of America

The IDSA published clinical practice guidelines in 2018 which addressed testing criteria for seasonal influenza A and B viruses. These guidelines state that serologic testing for the diagnosis of influenza should not be used by clinicians, because the results from a single serum specimen cannot be reliably interpreted. (p. 898)

Group A Streptococcus Pharyngitis Tests

Infectious Diseases Society of America

The IDSA published clinical practice guidelines in 2012 which addressed testing criteria for group A Streptococcal pharyngitis.

“Swabbing the throat and testing for GAS [group A Streptococcus] pharyngitis by rapid antigen detection test (RADT) and/or culture should be performed because the clinical features alone do not reliably discriminate between GAS and viral pharyngitis except when overt viral features like rhinorrhea, cough, oral ulcers, and/or hoarseness are present.” (p. e87)

“Patients with GAS pharyngitis commonly present with sore throat (generally of sudden onset), pain on swallowing, and fever. Headache, nausea, vomiting, and abdominal pain may also be present, especially in children. On examination, patients have tonsillopharyngeal erythema, with or without exudates, often with tender, enlarged anterior cervical lymph nodes (lymphadenitis). Other findings may include a beefy, red, swollen uvula; petechiae on the palate; excoriated nares (especially in infants); and a scarlatiniform rash.” (p. e91)

Group A Streptococcus Pharyngitis Culture

Infectious Diseases Society of America

The IDSA published clinical practice guidelines in 2012 which addressed testing criteria for group A Streptococcal pharyngitis.

“In children and adolescents, negative RADT [rapid antigen detection test] tests should be backed up by a throat culture...Routine use of back-up throat cultures for those with a negative RADT is not necessary for adults in usual circumstances, because of the low incidence of GAS [group A Streptococcus] pharyngitis in adults and because the risk of subsequent acute rheumatic fever is generally exceptionally low in adults with acute pharyngitis.” (p. e87)

CLINICAL POLICY

Infectious Disease: Respiratory Lab Testing

“Swabbing the throat and testing for GAS [group A Streptococcus] pharyngitis by rapid antigen detection test (RADT) and/or culture should be performed because the clinical features alone do not reliably discriminate between GAS and viral pharyngitis except when overt viral features like rhinorrhea, cough, oral ulcers, and/or hoarseness are present.” (p. e87)

“Patients with GAS pharyngitis commonly present with sore throat (generally of sudden onset), pain on swallowing, and fever. Headache, nausea, vomiting, and abdominal pain may also be present, especially in children. On examination, patients have tonsillopharyngeal erythema, with or without exudates, often with tender, enlarged anterior cervical lymph nodes (lymphadenitis). Other findings may include a beefy, red, swollen uvula; petechiae on the palate; excoriated nares (especially in infants); and a scarlatiniform rash.” (p. e91)

American Academy of Family Physicians

The 2024 American Academy of Family Physicians (~~AAFP~~) published guidelines for expert-authored evidence review concerning the diagnosis and treatment/management of streptococcal/streptococcus pharyngitis. This guideline defines states the age range between following:

“Rapid antigen testing may be omitted for patients at low clinical risk, including children younger than 3 and 14-years. (p. as a suggestive criterion 345)”

“The Centers for Disease Control and Prevention and the ~~diagnosis~~ American Academy of Streptococcal infection compared to other ages. Pediatrics recommend obtaining a throat culture for all children and adolescents after a negative result on rapid antigen testing because of the higher risk of complications.” (p. 345 (p. 385))

Group A Streptococcus Antibody Tests

Infectious Diseases Society of America

The IDSA published clinical practice guidelines in 2012 which addressed testing criteria for group A Streptococcal pharyngitis.

Per these guidelines, it is not recommended that individuals undergo anti-streptococcal antibody titers for the purpose of routine diagnosis of acute pharyngitis, as these results indicate a past infection and therefore do not aid in the diagnosis of the present illness. (p. e87)

“Measurement of anti-streptococcal antibody titers is often useful for diagnosis of the nonsuppurative sequelae of GAS pharyngitis, such as acute rheumatic fever and acute glomerulonephritis. However, such testing is not useful in the diagnosis of acute pharyngitis because antibody titers of the 2 most commonly used tests, antistreptolysin O (ASO) and

CLINICAL POLICY

Infectious Disease: Respiratory Lab Testing

antiDNase B, may not reach maximum levels until 3–8 weeks after acute GAS pharyngeal infection and may remain elevated for months even without active GAS infection.” (p. e93-94)

Coding Implications

This clinical policy references Current Procedural Terminology (CPT®). CPT® is a registered trademark of the American Medical Association. All CPT codes and descriptions are copyrighted 2023, American Medical Association. All rights reserved. CPT codes and CPT descriptions are from the current manuals and those included herein are not intended to be all-inclusive and are included for informational purposes only. Codes referenced in this clinical policy are for informational purposes only and may not support medical necessity. Inclusion or exclusion of any codes does not guarantee coverage. Providers should reference the most up-to-date sources of professional coding guidance prior to the submission of claims for reimbursement of covered services.

NOTE: Coverage is subject to each requested code’s inclusion on the corresponding LDH fee schedule. Non-covered codes are denoted (*) and are reviewed for Medical Necessity for members under 21 years of age on a per case basis.

CPT® Code	Description
86060	Antistreptolysin 0; titer
86328	Immunoassay for infectious agent antibody(ies), qualitative or semiquantitative, single-step method (eg, reagent strip); severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) (coronavirus disease [COVID-19])
86408	Neutralizing antibody, severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) (coronavirus disease [COVID-19]); screen
86409	Neutralizing antibody, severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) (coronavirus disease [COVID-19]); titer
86413	Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) (coronavirus disease [COVID-19]) antibody, quantitative
86710	Antibody; influenza virus
86769	Antibody; severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) (coronavirus disease [COVID-19])
87040	Culture, bacterial; blood, aerobic, with isolation and presumptive identification of isolates (includes anaerobic culture, if appropriate)
87070	Culture, bacterial; any other source except urine, blood or stool, aerobic, with isolation and presumptive identification of isolates

Infectious Disease: Respiratory Lab Testing

CPT® Code	Description
87071	Culture, bacterial; quantitative, aerobic with isolation and presumptive identification of isolates, any source except urine, blood or stool
87073	Culture, bacterial; quantitative, anaerobic with isolation and presumptive identification of isolates, any source except urine, blood or stool
87075	Culture, bacterial; any source, except blood, anaerobic with isolation and presumptive identification of isolates
87076	Culture, bacterial; anaerobic isolate, additional methods required for definitive identification, each isolate
87077	Culture, bacterial; aerobic isolate, additional methods required for definitive identification, each isolate
87081	Culture, presumptive, pathogenic organisms, screening only;
87084	Culture, presumptive, pathogenic organisms, screening only; with colony estimation from density chart
87101	Culture, fungi (mold or yeast) isolation, with presumptive identification of isolates; skin, hair, or nail
87102	Culture, fungi (mold or yeast) isolation, with presumptive identification of isolates; other source (except blood)
87103	Culture, fungi (mold or yeast) isolation, with presumptive identification of isolates; blood
87106	Culture, fungi, definitive identification, each organism; yeast
87107	Culture, fungi, definitive identification, each organism; mold
87109	Culture, mycoplasma, any source
87116	Culture, tubercle or other acid-fast bacilli (eg, TB, AFB, mycobacteria) any source, with isolation and presumptive identification of isolates
87118	Culture, mycobacterial, definitive identification, each isolate
87140	Culture, typing; immunofluorescent method, each antiserum
87143	Culture, typing; gas liquid chromatography (GLC) or high pressure liquid chromatography (HPLC) method
87147	Culture, typing; immunologic method, other than immunofluorescence (eg, agglutination grouping), per antiserum

Infectious Disease: Respiratory Lab Testing

CPT® Code	Description
87158	Culture, typing; other methods
87275	Infectious agent antigen detection by immunofluorescent technique; influenza B virus
87276	Infectious agent antigen detection by immunofluorescent technique; influenza A virus
87400	Infectious agent antigen detection by immunoassay technique (eg, enzyme immunoassay [EIA], enzyme-linked immunosorbent assay [ELISA], fluorescence immunoassay [FIA], immunochemiluminometric assay [IMCA]), qualitative or semiquantitative; Influenza, A or B, each
87420	Infectious agent antigen detection by immunoassay technique (eg, enzyme immunoassay [EIA], enzyme-linked immunosorbent assay [ELISA], fluorescence immunoassay [FIA], immunochemiluminometric assay [IMCA]), qualitative or semiquantitative; respiratory syncytial virus
87426	Infectious agent antigen detection by immunoassay technique (eg, enzyme immunoassay [EIA], enzyme-linked immunosorbent assay [ELISA], fluorescence immunoassay [FIA], immunochemiluminometric assay [IMCA]), qualitative or semiquantitative; severe acute respiratory syndrome coronavirus (eg, SARS-CoV, SARS-CoV-2 [COVID-19])
87428	Infectious agent antigen detection by immunoassay technique (eg, enzyme immunoassay [EIA], enzyme-linked immunosorbent assay [ELISA], fluorescence immunoassay [FIA], immunochemiluminometric assay [IMCA]), qualitative or semiquantitative; severe acute respiratory syndrome coronavirus (eg, SARS-CoV, SARS-CoV-2 [COVID-19]) and influenza virus types A and B
87430	Infectious agent antigen detection by immunoassay technique (eg, enzyme immunoassay [EIA], enzyme-linked immunosorbent assay [ELISA], fluorescence immunoassay [FIA], immunochemiluminometric assay [IMCA]), qualitative or semiquantitative; Streptococcus, group A
87480	Infectious agent detection by nucleic acid (DNA or RNA); Candida species, direct probe technique
87481	Infectious agent detection by nucleic acid (DNA or RNA); Candida species, amplified probe technique
87482	Infectious agent detection by nucleic acid (DNA or RNA); Candida species, quantification
87485	Infectious agent detection by nucleic acid (DNA or RNA); Chlamydia

Infectious Disease: Respiratory Lab Testing

CPT® Code	Description
	pneumoniae, direct probe technique
87486	Infectious agent detection by nucleic acid (DNA or RNA); Chlamydia pneumoniae, amplified probe technique
87487	Infectious agent detection by nucleic acid (DNA or RNA); Chlamydia pneumoniae, quantification
87498	Infectious agent detection by nucleic acid (DNA or RNA); enterovirus, amplified probe technique, includes reverse transcription when performed
87500	Infectious agent detection by nucleic acid (DNA or RNA); vancomycin resistance (eg, enterococcus species van A, van B), amplified probe technique
87501	Infectious agent detection by nucleic acid (DNA or RNA); influenza virus, includes reverse transcription, when performed, and amplified probe technique, each type or subtype
87502	Infectious agent detection by nucleic acid (DNA or RNA); influenza virus, for multiple types or sub-types, includes multiplex reverse transcription, when performed, and multiplex amplified probe technique, first 2 types or sub-types
87503	Infectious agent detection by nucleic acid (DNA or RNA); influenza virus, for multiple types or sub-types, includes multiplex reverse transcription, when performed, and multiplex amplified probe technique, each additional influenza virus type or sub-type beyond 2 (List separately in addition to code for primary procedure)
87540	Infectious agent detection by nucleic acid (DNA or RNA); Legionella pneumophila, direct probe technique
87541	Infectious agent detection by nucleic acid (DNA or RNA); Legionella pneumophila, amplified probe technique
87542	Infectious agent detection by nucleic acid (DNA or RNA); Legionella pneumophila, quantification
87550	Infectious agent detection by nucleic acid (DNA or RNA); Mycobacteria species, direct probe technique
87551	Infectious agent detection by nucleic acid (DNA or RNA); Mycobacteria species, amplified probe technique
87552	Infectious agent detection by nucleic acid (DNA or RNA); Mycobacteria species, quantification

Infectious Disease: Respiratory Lab Testing

CPT® Code	Description
87555	Infectious agent detection by nucleic acid (DNA or RNA); Mycobacteria tuberculosis, direct probe technique
87556	Infectious agent detection by nucleic acid (DNA or RNA); Mycobacteria tuberculosis, amplified probe technique
87560	Infectious agent detection by nucleic acid (DNA or RNA); Mycobacteria avium-intracellulare, direct probe technique
87561	Infectious agent detection by nucleic acid (DNA or RNA); Mycobacteria avium-intracellulare, amplified probe technique
87580	Infectious agent detection by nucleic acid (DNA or RNA); Mycoplasma pneumoniae, direct probe technique
87581	Infectious agent detection by nucleic acid (DNA or RNA); Mycoplasma pneumoniae, amplified probe technique
87582	Infectious agent detection by nucleic acid (DNA or RNA); Mycoplasma pneumoniae, quantification
87631¹	Infectious agent detection by nucleic acid (DNA or RNA); respiratory virus (eg, adenovirus, influenza virus, coronavirus, metapneumovirus, parainfluenza virus, respiratory syncytial virus, rhinovirus), includes multiplex reverse transcription, when performed, and multiplex amplified probe technique, multiple types or subtypes, 3-5 targets
87632¹87632*	Infectious agent detection by nucleic acid (DNA or RNA); respiratory virus (eg, adenovirus, influenza virus, coronavirus, metapneumovirus, parainfluenza virus, respiratory syncytial virus, rhinovirus), includes multiplex reverse transcription, when performed, and multiplex amplified probe technique, multiple types or subtypes, 6-11 targets
87633¹87633*	Infectious agent detection by nucleic acid (DNA or RNA); respiratory virus (eg, adenovirus, influenza virus, coronavirus, metapneumovirus, parainfluenza virus, respiratory syncytial virus, rhinovirus), includes multiplex reverse transcription, when performed, and multiplex amplified probe technique, multiple types or subtypes, 12-25 targets
87634	Infectious agent detection by nucleic acid (DNA or RNA); respiratory syncytial virus, amplified probe technique
87635	Infectious agent detection by nucleic acid (DNA or RNA); severe acute

¹ See section Criteria “Coverage Specific guidelines” for information regarding CPT codes 87631, 87632, 87633.

Infectious Disease: Respiratory Lab Testing

CPT® Code	Description
	respiratory syndrome coronavirus 2 (SARS-CoV-2) (coronavirus disease [COVID-19]), amplified probe technique
87636	Infectious agent detection by nucleic acid (DNA or RNA); severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) (coronavirus disease [COVID-19]) and influenza virus types A and B, multiplex amplified probe technique
87637	Infectious agent detection by nucleic acid (DNA or RNA); severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) (coronavirus disease [COVID-19]), influenza virus types A and B, and respiratory syncytial virus, multiplex amplified probe technique
87640	Infectious agent detection by nucleic acid (DNA or RNA); Staphylococcus aureus, amplified probe technique
87641	Infectious agent detection by nucleic acid (DNA or RNA); Staphylococcus aureus, methicillin resistant, amplified probe technique
87650	Infectious agent detection by nucleic acid (DNA or RNA); Streptococcus, group A, direct probe technique
87651	Infectious agent detection by nucleic acid (DNA or RNA); Streptococcus, group A, amplified probe technique
87652	Infectious agent detection by nucleic acid (DNA or RNA); Streptococcus, group A, quantification
87653	Infectious agent detection by nucleic acid (DNA or RNA); Streptococcus, group B, amplified probe technique
87797	Infectious agent detection by nucleic acid (DNA or RNA), not otherwise specified; direct probe technique, each organism
87798	Infectious agent detection by nucleic acid (DNA or RNA), not otherwise specified; amplified probe technique, each organism
87799	Infectious agent detection by nucleic acid (DNA or RNA), not otherwise specified; quantification, each organism
87800	Infectious agent detection by nucleic acid (DNA or RNA), multiple organisms; direct probe(s) technique
87801	Infectious agent detection by nucleic acid (DNA or RNA), multiple organisms; amplified probe(s) technique

Infectious Disease: Respiratory Lab Testing

CPT® Code	Description
87804	Infectious agent antigen detection by immunoassay with direct optical (ie, visual) observation; Influenza
87807	Infectious agent antigen detection by immunoassay with direct optical (ie, visual) observation; respiratory syncytial virus
87811	Infectious agent antigen detection by immunoassay with direct optical (ie, visual) observation; severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) (coronavirus disease [COVID-19])
87880	Infectious agent antigen detection by immunoassay with direct optical (ie, visual) observation; Streptococcus, group A
87913	Infectious agent genotype analysis by nucleic acid (DNA or RNA); severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) (coronavirus disease [COVID-19]), mutation identification in targeted region(s)
0202U*	Infectious disease (bacterial or viral respiratory tract infection), pathogenspecific nucleic acid (DNA or RNA), 22 targets including severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), qualitative RT-PCR, nasopharyngeal swab, each pathogen reported as detected or not detected (For additional PLA code with identical clinical descriptor, see 0223U. See Appendix O or the most current listing on the AMA CPT website to determine appropriate code assignment)
0223U*	Infectious disease (bacterial or viral respiratory tract infection), pathogenspecific nucleic acid (DNA or RNA), 22 targets including severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), qualitative RT-PCR, nasopharyngeal swab, each pathogen reported as detected or not detected (For additional PLA code with identical clinical descriptor, see 0202U. See Appendix O or the most current listing on the AMA CPT website to determine appropriate code assignment)
0225U*	Infectious disease (bacterial or viral respiratory tract infection) pathogen-specific DNA and RNA, 21 targets, including severe acute respiratory syndrome coronavirus 2 (SARSCoV-2), amplified probe technique, including multiplex reverse transcription for RNA targets, each analyte reported as detected or not detected
0240U*	Infectious disease (viral respiratory tract infection), pathogen-specific RNA, 3 targets (severe acute respiratory syndrome coronavirus 2 [SARS-CoV-2], influenza A, influenza B), upper respiratory specimen, each pathogen reported as detected or not detected

Infectious Disease: Respiratory Lab Testing

CPT® Code	Description
0241U*	Infectious disease (viral respiratory tract infection), pathogen-specific RNA, 4 targets (severe acute respiratory syndrome coronavirus 2 [SARS-CoV-2], influenza A, influenza B, respiratory syncytial virus [RSV]), upper respiratory specimen, each pathogen reported as detected or not detected
<u>0528U</u>	<u>Lower respiratory tract infectious agent detection, 18 bacteria, 8 viruses, and 7 antimicrobial-resistance genes, amplified probe technique, including reverse transcription for RNA targets, each analyte reported as detected or not detected with semiquantitative results for 15 bacteria</u>
U0001*	CDC Test
U0002*	Non-CDC Viral identification test, amplified probe
U0003*	High throughput Viral identification test, amplified probe
U0004*	High throughput Viral identification test, other than amplified probe
U0005*	Infectious agent detection by nucleic acid (dna or rna); severe acute respiratory syndrome coronavirus 2 (sars-cov-2) (coronavirus disease [covid-19]), amplified probe technique, cdc or non-cdc, making use of high throughput technologies, completed within 2 calendar days from date of specimen collection (list separately in addition to either hcpcs code u0003 or u0004) as described by cms-2020-01-r2

Infectious Disease: Respiratory Lab Testing

Reviews, Revisions, and Approvals	Revision Date	Approval Date
Converted corporate to local policy.	03/24	5/1/24
Added (*) to codes 0202U, 0223U, 0225U, 0240U and 0241U per LDH’s IB 24-16. Did not send to LDH for review as revisions were per IB 24-16.	07/24	7/10/24
<u>Removed (*) from codes 87631, 87632, and 87633. Added footnote to page 16. Added section “Coverage Specific Guidelines with information regarding coverage of the codes and meeting medical necessity” from IB 24-31.</u>	<u>10/24</u>	<u>1/3/25</u>
<u>Annual review. References and background reviewed and updated. Changed verbiage in policy statements from “may be considered medically necessary” to “are considered medically necessary. Noted that CPT 87631 is out of scope for this policy in criteria for SARS-CoV-2, RSV, or Influenza A/B, or Multiplex Respiratory Viral Panels with 5 or Fewer Targets and removed from the CPT table. In criteria for Syndromic/Multiplex Respiratory Panels with 6 or More Targets, clarified that criteria applies when performed in the outpatient setting and replaced prior criteria with the following options: “With serious or critical illness, OR At imminent risk of becoming seriously or critically ill, OR With immunodeficiency, AND/OR With a severe underlying condition”, per Louisiana Department of Health Informational Bulletin 24-31. Added 0528U as an in-scope code in the CPT table.</u>	<u>2/25</u>	

REFERENCES

1. Uyeki TM, Bernstein HH, Bradley JS, et al. Clinical Practice Guidelines by the Infectious Diseases Society of America: 2018 Update on Diagnosis, Treatment, Chemoprophylaxis, and Institutional Outbreak Management of Seasonal Influenza. *Clin Infect Dis.* 2019;68(6):895-902. doi:10.1093/cid/ciy874
2. Shulman ST, Bisno AL, Clegg HW, et al. Clinical practice guideline for the diagnosis and management of group A streptococcal pharyngitis: 2012 update by the Infectious Diseases Society of America. *Clin Infect Dis.* 2012;55(10):e86-e102. doi:10.1093/cid/cis629
3. Hanson KE, Azar MM, Banerjee R, et al. Molecular Testing for Acute Respiratory Tract Infections: Clinical and Diagnostic Recommendations From the IDSA's Diagnostics Committee. *Clin Infect Dis.* 2020;71(10):2744-2751. doi:10.1093/cid/ciaa508

CLINICAL POLICY

Infectious Disease: Respiratory Lab Testing

4. Choby BA. Diagnosis and treatment of streptococcal pharyngitis. *Am Fam Physician*. 2009;79(5):383-390.
5. CDC. Learn more about the flu season. Centers for Disease Control and Prevention. Published September 20, 2022. <https://www.cdc.gov/flu/about/season/index.html>. Accessed ~~January 4~~November 8, 2024.
6. RSV. Center for Disease Control and Prevention. Published 2019. <https://www.cdc.gov/rsv/clinical/index.html>. Accessed ~~January 4~~November 8, 2024.
7. CDC. COVID-19 and Your Health. Centers for Disease Control and Prevention. Published February 11, 2020. <https://www.cdc.gov/coronavirus/2019-ncov/symptoms-testing/testing.html#when-to-get-tested>. Accessed ~~January 4~~November 8, 2024.
8. Louisiana Department of Health. Informational Bulletin 24-31. Published September 18, 2024.

Important Reminder

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program approval status; evidence-based guidelines and positions of leading national health professional organizations; views of physicians practicing in relevant clinical areas affected by this clinical policy; and other available clinical information. LHCC makes no representations and accepts no liability with respect to the content of any external information used or relied upon in developing this clinical policy. This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved.

The purpose of this clinical policy is to provide a guide to medical necessity, which is a component of the guidelines used to assist in making coverage decisions and administering benefits. It does not constitute a contract or guarantee regarding payment or results. Coverage decisions and the administration of benefits are subject to all terms, conditions, exclusions and limitations of the coverage documents (e.g., evidence of coverage, certificate of coverage, policy, contract of insurance, etc.), as well as to state and federal requirements and applicable LHCC administrative policies and procedures.

This clinical policy is effective as of the date determined by LHCC. The date of posting may not be the effective date of this clinical policy. This clinical policy may be subject to applicable legal and regulatory requirements relating to provider notification. If there is a discrepancy between the effective date of this clinical policy and any applicable legal or regulatory requirement, the requirements of law and regulation shall govern. LHCC retains the right to change, amend or withdraw this clinical policy, and additional clinical policies may be developed and adopted as needed, at any time.

This clinical policy does not constitute medical advice, medical treatment or medical care. It is not intended to dictate to providers how to practice medicine. Providers are expected to exercise professional medical judgment in providing the most appropriate care, and are solely responsible for the medical advice and treatment of members/enrollees. This clinical policy is not intended to

CLINICAL POLICY

Infectious Disease: Respiratory Lab Testing

recommend treatment for members/enrollees. Members/enrollees should consult with their treating physician in connection with diagnosis and treatment decisions.

Providers referred to in this clinical policy are independent contractors who exercise independent judgment and over whom LHCC has no control or right of control. Providers are not agents or employees of LHCC.

This clinical policy is the property of LHCC. Unauthorized copying, use, and distribution of this clinical policy or any information contained herein are strictly prohibited. Providers, members/enrollees and their representatives are bound by the terms and conditions expressed herein through the terms of their contracts. Where no such contract exists, providers, members/enrollees and their representatives agree to be bound by such terms and conditions by providing services to members/enrollees and/or submitting claims for payment for such services.

©2024 Louisiana Healthcare Connections. All rights reserved. All materials are exclusively owned by Louisiana Healthcare Connections and are protected by United States copyright law and international copyright law. No part of this publication may be reproduced, copied, modified, distributed, displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise published without the prior written permission of Louisiana Healthcare Connections. You may not alter or remove any trademark, copyright or other notice contained herein. Louisiana Healthcare Connections is a registered trademarks exclusively owned by Louisiana Healthcare Connections.