

## Clinical Policy: Lutetium Lu 177 vipivotide tetraxetan (Pluvicto)

Reference Number: LA.PHAR.582

Effective Date: ~~09.29.23~~06.20.24

Last Review Date: ~~03.25.24~~11.21.24

Line of Business: Medicaid

[Coding Implications](#)

[Revision Log](#)

See [Important Reminder](#) at the end of this policy for important regulatory and legal information.

\*\*Please note: This policy is for medical benefit\*\*

### Description

Lutetium Lu 177 vipivotide tetraxetan (Pluvicto™) is a radioligand therapeutic agent.

### FDA Approved Indication(s)

Pluvicto is indicated for the treatment of adult patients with prostate-specific membrane antigen (PSMA)-positive metastatic castration-resistant prostate cancer (mCRPC) who have been treated with androgen receptor (AR) pathway inhibition and taxane-based chemotherapy.

### Policy/Criteria

*Provider must submit documentation (such as office chart notes, lab results or other clinical information) supporting that member has met all approval criteria.*

It is the policy of Louisiana Healthcare Connections® that Pluvicto is **medically necessary** when the following criteria are met:

#### I. Initial Approval Criteria

##### A. Metastatic Castration-resistant Prostate Cancer (must meet all):

1. Diagnosis of metastatic CRPC;
2. Documentation of disease progression despite bilateral orchiectomy or other androgen deprivation therapy (ADT) (see *Appendix D*);
3. Documentation of PSMA-positive disease confirmed on a ~~GAGa-68~~-PSMA-11 ~~or~~, F-18 piflufolastat, or F-18 flotufolastat positive emission tomography (PET) or computed tomography (CT) scan;
4. Prescribed by or in consultation with an oncologist or urologist;
5. Age ≥ 18 years;
6. Member will use a gonadotropin-releasing hormone (GnRH) analog concurrently or has had a bilateral orchiectomy;
7. Failure of both of the following, unless contraindicated or clinically significant adverse effects are experienced or all are contraindicated (a and b):
  - a. A taxane-based regimen (e.g. docetaxel, cabazitaxel);\*  
*\*Prior authorization may be required for docetaxel and cabazitaxel*
  - b. Abiraterone (Zytiga®), unless member has previously failed Yonsa® (abiraterone) or Xtandi® (enzalutamide);\*  
*\*Prior authorization may be required for Zytiga, Yonsa, and Xtandi*

## CLINICAL POLICY

### Lutetium Lu 177 vipivotide tetraxetan

8. Pluvicto is not prescribed concurrently with cytotoxic chemotherapy, immunotherapy, radioligand therapy, or investigational therapy;

## CLINICAL POLICY

### Lutetium Lu 177 vipivotide tetraxetan

9. Request meets one of the following (a or b):\*
  - a. Dose does not exceed 7.4 GBq (200 mCi) every 6 weeks for up to 6 doses;
  - b. Dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).

*\*Prescribed regimen must be FDA-approved or recommended by NCCN*

**Approval duration: 6 months (up to a total of 6 doses)**

#### **B. Other diagnoses/indications** (must meet 1 or 2):

1. If this drug has recently (within the last 6 months) undergone a label change (e.g., newly approved indication, age expansion, new dosing regimen) that is not yet reflected in this policy, refer to LA.PMN.255
2. If the requested use (e.g., diagnosis, age, dosing regimen) is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized) AND criterion 1 above does not apply, refer to the off-label use policy for the relevant line of business: LA.PMN.53 for Medicaid.

## II. Continued Therapy

#### **A. Metastatic Castration-resistant Prostate Cancer** (must meet all):

1. Currently receiving medication via Louisiana Healthcare Connections benefit, or documentation supports that member is currently receiving Pluvicto for a covered indication and has received this medication for at least 30 days;
2. Member is responding positively to therapy;
3. Member has not received  $\geq 6$  doses (infusions) of Pluvicto;
4. If request is for a dose increase, request meets one of the following (a or b):\*
  - a. New dose does not exceed 7.4 GBq (200 mCi) every 6 weeks for up to 6 doses;
  - b. New dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).

*\*Prescribed regimen must be FDA-approved or recommended by NCCN*

**Approval duration: 6 months (up to a total of 6 doses)**

#### **B. Other diagnoses/indications** (must meet 1 or 2):

1. If this drug has recently (within the last 6 months) undergone a label change (e.g., newly approved indication, age expansion, new dosing regimen) that is not yet reflected in this policy, refer to LA.PMN.255
2. If the requested use (e.g., diagnosis, age, dosing regimen) is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized) AND criterion 1 above does not apply, refer to the off-label use policy for the relevant line of business: LA.PMN.53 for Medicaid.

**Approval duration: Duration of request or 6 months (whichever is less)**

## CLINICAL POLICY

### Lutetium Lu 177 vipivotide tetraxetan

#### III. Diagnoses/Indications for which coverage is NOT authorized:

- A. Non-FDA approved indications, which are not addressed in this policy, unless there is sufficient documentation of efficacy and safety according to the off label use policies – LA.PMN.53 for Medicaid or evidence of coverage documents.

#### IV. Appendices/General Information

##### *Appendix A: Abbreviation/Acronym Key*

ADT: androgen deprivation therapy

AR: androgen receptor

BSOC: best standard of care

CRPC: castration- resistant prostate cancer

CT: computed tomography

FDA: Food and Drug Administration

GnRH: gonadotropin-releasing hormone

LHRH: luteinizing hormone-releasing hormone

NCCN: National Comprehensive Cancer Network

PET: positive emission tomography

PSMA: prostate- specific membrane antigen

##### *Appendix B: Therapeutic Alternatives*

*This table provides a listing of preferred alternative therapy recommended in the approval criteria. The drugs listed here may not be a formulary agent for all relevant lines of business and may require prior authorization.*

Drug Name	Dosing Regimen	Dose Limit/ Maximum Dose
abiraterone (Zytiga®)	1,000 mg PO QD (given in combination with prednisone)	1,000 mg/day; 2,000 mg/day if taking a strong CYP3A4 inducer
docetaxel	Androgen-deprivation therapy with docetaxel 75 mg/m <sup>2</sup> for 6 cycles	Varies
Jevtana® (cabazitaxel)	20 mg/m <sup>2</sup> IV every 3 weeks	25 mg/m <sup>2</sup> once every 3 weeks

*Therapeutic alternatives are listed as Brand name® (generic) when the drug is available by brand name only and generic (Brand name®) when the drug is available by both brand and generic.*

##### *Appendix C: Contraindications/Boxed Warnings*

None reported

##### *Appendix D: General Information*

- Castration-resistant prostate cancer is prostate cancer that progresses clinically, radiographically, or biochemically despite castrate levels of serum testosterone (< 50 ng/dL).
- Per the NCCN, androgen deprivation therapy (ADT) should be continued in patients with metastatic CRPC while additional therapies, including secondary hormone therapies, chemotherapies, immunotherapies, radiopharmaceuticals, and/or targeted therapies are applied.

## CLINICAL POLICY

### Lutetium Lu 177 vipivotide tetraxetan

- Examples of ADT include:
  - Bilateral orchiectomy (surgical castration)
  - Luteinizing hormone-releasing hormone (LHRH) agonist given with or without an anti-androgen:
    - LHRH agonists: Zoladex® (goserelin), leuprolide (Lupron Depot®, Eligard®), and Trelstar® (triptorelin)
    - Anti-androgens: bicalutamide (Casodex®), flutamide (Eulexin®), nilutamide (Nilandron®), Xtandi® (enzalutamide), Erleada® (apalutamide), Nubeqa® (darolutamide)
  - LHRH ~~antagonist~~antagonists: Firmagon® (degarelix), Orgovyx® (relugolix)

#### V. Dosage and Administration

Indication	Dosing Regimen	Maximum Dose
Metastatic CRPC	7.4 GBq (200 mCi) IV every 6 weeks for up to 6 doses	See dosing regimen

#### VI. Product Availability

Injection, single-dose vial: 1,000 MBq/mL (27 mCi/mL)

#### VII. References

1. Pluvicto Prescribing Information. Millburn, NJ: Novartis AG.; October 2022. Available at [https://www.novartis.com/us-en/sites/novartis\\_us/files/pluvicto.pdf](https://www.novartis.com/us-en/sites/novartis_us/files/pluvicto.pdf). Accessed January ~~6,~~ 202310, 2024.
2. National Comprehensive Cancer Network. Prostate Cancer Version ~~14.~~2023. Available at [https://www.nccn.org/professionals/physician\\_gls/pdf/prostate.pdf](https://www.nccn.org/professionals/physician_gls/pdf/prostate.pdf). Accessed January ~~11,~~ 202322, 2024.
3. ClinicalTrials.gov. Study of 177Lu-PSMA-617 in Metastatic Castrate-Resistant Prostate Cancer (VISION). Available at <https://clinicaltrials.gov/ct2/show/NCT03511664>. Accessed ~~April 15, 2022~~January 22, 2024.
4. IPD Analytics. NOC Code Guide: Pluvicto (lutetium Lu 177 vipivotide tetraxetan) injection, for intravenous use by Advanced Accelerator Applications USA, Inc. Published April 2022.
5. IPD Analytics. New Drug Review: Pluvicto (lutetium Lu 177 vipivotide tetraxetan). Published April 6, 2022.
6. Virgo KS, Rumble B, de Wit R, et al. Initial Management of Non-Castrate Advanced, Recurrent or Metastatic Prostate Cancer. American Society of Clinical Oncology (ASCO). Published ahead of print ~~January 26, 2021~~April 3, 2023, DOI: 10.1200/JCO.20.03256. Available at: <https://www.asco.org/practice-patients/guidelines/genitourinary-cancer#/9521>. Accessed January ~~11,~~ 202322, 2024.
7. Basch E, Loblaw DA, Oliver TK, et al. Systemic Therapy in Men with Metastatic Castration-Resistant Prostate Cancer (CRPC). American Society of Clinical Oncology (ASCO). Published online before print September 8, 2014. Available at: <https://ascopubs.org/doi/full/10.1200/JCO.2013.54.8404>. Accessed January ~~11,~~ 202322, 2024.

## CLINICAL POLICY

### Lutetium Lu 177 vipivotide tetraxetan

8. Garje R, Rumble B, Parikh RA. Systemic Therapy Update on 177Lutetium-PSMA-617 for Metastatic Castration-Resistant Prostate Cancer: ASCO Rapid Recommendation. *Journal of Clinical Oncology* 2022. 40(31): 3664-3666. Available at: <https://www.asco.org/practice-patients/guidelines/genitourinary-cancer#/9496>.
9. Garje R, Rumble B, Parikh RA. Systemic Therapy Update on 177Lutetium-PSMA-617 for Metastatic Castration-Resistant Prostate Cancer: ASCO Guideline Rapid Recommendation Update. *Journal of Clinical Oncology*. November 6, 2023. Available at: <https://ascopubs.org/doi/10.1200/JCO.23.02128>. Accessed January 22, 2024.
- 9.10. Sartor O, de Bono J, Chi KN, et al. Lutetium-177-PSMA-617 for Metastatic Castration-Resistant Prostate Cancer. *N Engl J Med*. 2021 Sep 16; 385(12): 1091-1103.

### Coding Implications

Codes referenced in this clinical policy are for informational purposes only. Inclusion or exclusion of any codes does not guarantee coverage. Providers should reference the most up-to-date sources of professional coding guidance prior to the submission of claims for reimbursement of covered services.

HCPSC Codes	Description
A9607	Lutetium Lu 177 vipivotide tetraxetan, therapeutic, 1 <del>mCi</del> millicurie

Reviews, Revisions, and Approvals	Date	LDH Approval Date
Policy created	05.01.23	08.28.23
Annual review: added clarification to approval duration is for up to a total of 6 doses; revised continued therapy approval duration from 12 to 6 months; for continued therapy added requirement that member has not received $\geq 6$ doses (infusions) of Pluvicto; added piflufolastat F-18 as an additional radioactive diagnostic agent for identification of PSMA-positive disease; updated <i>Appendix D</i> examples of androgen deprivation therapy per NCCN; removed inactive HCPSC code A9699; references reviewed and updated.	03.25.24	<u>06.20.24</u>
<u>Added F-18 flutufolastat as an additional option to confirm PSMA-positive disease; references reviewed and updated.</u>	<u>11.21.24</u>	

### Important Reminder

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program approval status; evidence-based guidelines and positions of leading national health professional organizations; views of physicians practicing in relevant clinical areas affected by this clinical policy; and other available clinical information. LHCC makes no representations and accepts no liability with respect to the content of any external information used or relied upon in developing

## CLINICAL POLICY

### Lutetium Lu 177 vipivotide tetraxetan

this clinical policy. This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved.

The purpose of this clinical policy is to provide a guide to medical necessity, which is a component of the guidelines used to assist in making coverage decisions and administering benefits. It does not constitute a contract or guarantee regarding payment or results. Coverage decisions and the administration of benefits are subject to all terms, conditions, exclusions, and limitations of the coverage documents (e.g., evidence of coverage, certificate of coverage, policy, contract of insurance, etc.), as well as to state and federal requirements and applicable LHCC administrative policies and procedures.

This clinical policy is effective as of the date determined by LHCC. The date of posting may not be the effective date of this clinical policy. This clinical policy may be subject to applicable legal and regulatory requirements relating to provider notification. If there is a discrepancy between the effective date of this clinical policy and any applicable legal or regulatory requirement, the requirements of law and regulation shall govern. LHCC retains the right to change, amend or withdraw this clinical policy, and additional clinical policies may be developed and adopted as needed, at any time.

This clinical policy does not constitute medical advice, medical treatment, or medical care. It is not intended to dictate to providers how to practice medicine. Providers are expected to exercise professional medical judgment in providing the most appropriate care, and are solely responsible for the medical advice and treatment of members. This clinical policy is not intended to recommend treatment for members. Members should consult with their treating physician in connection with diagnosis and treatment decisions.

Providers referred to in this clinical policy are independent contractors who exercise independent judgment and over whom LHCC has no control or right of control. Providers are not agents or employees of LHCC.

This clinical policy is the property of LHCC. Unauthorized copying, use, and distribution of this clinical policy or any information contained herein are strictly prohibited. Providers, members, and their representatives are bound to the terms and conditions expressed herein through the terms of their contracts. Where no such contract exists, providers, members and their representatives agree to be bound by such terms and conditions by providing services to members and/or submitting claims for payment for such services.

©2024 Louisiana Healthcare Connections. All rights reserved. All materials are exclusively owned by Louisiana Healthcare Connections and are protected by United States copyright law and international copyright law. No part of this publication may be reproduced, copied, modified, distributed, displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise published without the prior written permission of Louisiana Healthcare Connections. You may not alter or remove any trademark, copyright or other notice contained

## **CLINICAL POLICY**

### **Lutetium Lu 177 vipivotide tetraxetan**

herein. Louisiana Healthcare Connections is a registered trademark exclusively owned by Louisiana Healthcare Connections.