

Clinical Policy: Melphalan Flufenamide (Pepaxto)

Reference Number: LA.PHAR.535

Effective Date: ~~09.29.23~~05.23.24

Last Review Date: ~~03.25.24~~11.20.24

Line of Business: Medicaid

[Coding Implications](#)

[Revision Log](#)

See [Important Reminder](#) at the end of this policy for important regulatory and legal information.

****Please note: This policy is for medical benefit****

Description

Melphalan flufenamide (Pepaxto®) is an alkylating drug.

FDA Approved Indication(s)*

Pepaxto is indicated in combination with dexamethasone for the treatment of adult patients with relapsed or refractory multiple myeloma who have received at least 4 prior lines of therapy and whose disease is refractory to at least one proteasome inhibitor, one immunomodulatory agent, and one CD38-directed monoclonal antibody.

This indication is approved under accelerated approval based on response rate. Continued approval for this indication may be contingent upon verification and description of clinical benefit in confirmatory trial(s).

Limitation(s) of use: Pepaxto is not indicated and is not recommended for use as a conditioning regimen for transplant outside of controlled clinical trials.

***Oncopptides, the manufacturer of Pepaxto, voluntarily withdrew Pepaxto after data from the confirmatory Phase 3 OCEAN study revealed Pepaxto did not meet the requirements of the FDA Accelerated Approval regulation. The FDA withdrew its approval for the product (see Appendix D).**

Policy/Criteria

Provider must submit documentation (such as office chart notes, lab results or other clinical information) supporting that member has met all approval criteria.

It is the policy of Louisiana Healthcare Connections® that Pepaxto is **medically necessary** when the following criteria are met:

I. Initial Approval Criteria

A. Multiple Myeloma (must meet all):

1. Provider attestation of acknowledgement of FDA's request for withdrawal of product due to reduced overall survival and failure to demonstrate superior progression-free survival (PFS) compared to Pomalyst® in combination with dexamethasone;
2. Diagnosis of multiple myeloma;
3. Prescribed by or in consultation with an oncologist or hematologist;
4. Age ≥ 18 years;
5. Pepaxto is prescribed in combination with dexamethasone;

CLINICAL POLICY

Melphalan Flufenamide

6. Member has received ≥ 4 prior lines of therapy (*see Appendix B for examples*) that include all of the following (a, b, and c):
 - a. One proteasome inhibitor (e.g., bortezomib, Kyprolis®, Ninlaro®);
 - b. One immunomodulatory agent (e.g., ~~Revlimid®~~, lenalidomide, Pomalyst®, Thalomid®);
 - c. One anti-CD38 antibody (e.g., Darzalex®/Darzalex FasproTM ~~5~~₂, Sarclisa®);
**Prior authorization may be required*
7. Request meets one of the following (a or b):*
 - a. Dose does not exceed 40 mg every 4 weeks;
 - b. Dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).
**Prescribed regimen must be FDA-approved or recommended by NCCN*

Approval duration: 6 months

B. Other diagnoses/indications (must meet 1 or 2):

1. If this drug has recently (within the last 6 months) undergone a label change (e.g., newly approved indication, age expansion, new dosing regimen) that is not yet reflected in this policy, refer to LA.PMN.255
2. If the requested use (e.g., diagnosis, age, dosing regimen) is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized) AND criterion 1 above does not apply, refer to the off-label use policy for the relevant line of business: LA.PMN.53 for Medicaid.

II. Continued Therapy

A. Multiple Myeloma (must meet all):

1. Provider attestation of acknowledgement of FDA's request for withdrawal of product due to reduced overall survival and failure to demonstrate superior PFS compared to Pomalyst in combination with dexamethasone;
2. Currently receiving medication via Louisiana Healthcare Connections benefit, or documentation supports that member is currently receiving Pepaxto for a covered indication and has received this medication for at least 30 days;
3. Member is responding positively to therapy;
4. Pepaxto is prescribed in combination with dexamethasone;
5. If request is for a dose increase, request meets one of the following (a or b):*
 - a. New dose does not exceed 40 mg every 4 weeks;
 - b. New dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).
**Prescribed regimen must be FDA-approved or recommended by NCCN*

Approval duration: 12 months

B. Other diagnoses/indications (must meet 1 or 2):

1. If this drug has recently (within the last 6 months) undergone a label change (e.g., newly approved indication, age expansion, new dosing regimen) that is not yet reflected in this policy, refer to LA.PMN.255

CLINICAL POLICY

Melphalan Flufenamide

2. If the requested use (e.g., diagnosis, age, dosing regimen) is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized) AND criterion 1 above does not apply, refer to the off-label use policy for the relevant line of business: LA.PMN.53 for Medicaid.

III. Diagnoses/Indications for which coverage is NOT authorized:

- A. Non-FDA approved indications, which are not addressed in this policy, unless there is sufficient documentation of efficacy and safety according to the off label use policies – LA.PMN.53 for Medicaid or evidence of coverage documents.

IV. Appendices/General Information

Appendix A: Abbreviation/Acronym Key

CI: confidence interval

FDA: Food and Drug Administration

HR: hazard ratio

PFS: progression free survival

Appendix B: Therapeutic Alternatives

This table provides a listing of preferred alternative therapy recommended in the approval criteria. The drugs listed here may not be a formulary agent for all relevant lines of business and may require prior authorization.

Drug Name	Dosing Regimen	Dose Limit/ Maximum Dose
bortezomib/ Revlimid [®] -(lenalidomide)/dexamethasone	Varies	Varies
bortezomib/cyclophosphamide/dexamethasone	Varies	Varies
bortezomib/doxorubicin (or liposomal doxorubicin)/dexamethasone	Varies	Varies
Kyprolis [®] (carfilzomib)- Revlimid [®] -(lenalidomide)/dexamethasone	Varies	Varies
Kyprolis [®] (carfilzomib)/cyclophosphamide/dexamethasone	Varies	Varies
Kyprolis [®] (carfilzomib – weekly or twice weekly)/dexamethasone	Varies	Varies
Ninlaro [®] (ixazomib)/ Revlimid [®] -(lenalidomide)/dexamethasone	Varies	Varies
Ninlaro [®] (ixazomib)/dexamethasone	Varies	Varies
Ninlaro [®] (ixazomib)/pomalidomide/dexamethasone	Varies	Varies
bortezomib/dexamethasone	Varies	Varies
bortezomib/Thalomid [®] (thalidomide)/dexamethasone	Varies	Varies
cyclophosphamide/ Revlimid [®] -(lenalidomide)/dexamethasone	Varies	Varies
Revlimid [®] -(lenalidomide)/dexamethasone	Varies	Varies
VTD-PACE (dexamethasone/Thalomid [®] (thalidomide)/cisplatin/doxorubicin/cyclophosphamide/etoposide/bortezomib)	Varies	Varies

CLINICAL POLICY

Melphalan Flufenamide

Drug Name	Dosing Regimen	Dose Limit/ Maximum Dose
Revlimid® (lenalidomide) /low-dose dexamethasone	Varies	Varies
Darzalex® (daratumumab) or Darzalex Faspro™ (daratumumab/hyaluronidase-fihj)/bortezomib/melphan/prednisone	Varies	Varies
Darzalex® (daratumumab) or Darzalex Faspro™ (daratumumab/hyaluronidase-fihj)/bortezomib/dexamethasone	Varies	Varies
Darzalex® (daratumumab) or Darzalex Faspro™ (daratumumab/hyaluronidase-fihj)/Revlimid® (lenalidomide)/dexamethasone	Varies	Varies
Darzalex® (daratumumab) or Darzalex Faspro™ (daratumumab/hyaluronidase-fihj)	Varies	Varies
Darzalex® (daratumumab) or Darzalex Faspro™ (daratumumab/hyaluronidase-fihj)/pomalidomide/dexamethasone	Varies	Varies
Empliciti® (elotuzumab)/ Revlimid® (lenalidomide) /dexamethasone	Varies	Varies
Empliciti® (elotuzumab)/bortezomib/dexamethasone	Varies	Varies
Empliciti® (elotuzumab)/pomalidomide/dexamethasone	Varies	Varies
bendamustine/bortezomib/dexamethasone	Varies	Varies
bendamustine/ Revlimid® (lenalidomide) /dexamethasone	Varies	Varies
panobinostat/bortezomib/dexamethasone	Varies	Varies
panobinostat/Kyprolis® (carfilzomib)	Varies	Varies
panobinostat/ Revlimid® (lenalidomide) /dexamethasone	Varies	Varies
pomalidomide/cyclophosphamide/dexamethasone	Varies	Varies
pomalidomide/dexamethasone	Varies	Varies
pomalidomide/bortezomib/dexamethasone	Varies	Varies
pomalidomide/Kyprolis® (carfilzomib)/dexamethasone	Varies	Varies
Sarclisa® (isatuximab-irfc)/pomalidomide/dexamethasone	Varies	Varies

Therapeutic alternatives are listed as Brand name® (generic) when the drug is available by brand name only and generic (Brand name®) when the drug is available by both brand and generic.

Appendix C: Contraindications/Boxed Warnings

- Contraindication(s): history of serious hypersensitivity reaction to melphalan flufenamide or melphalan
- Boxed warning(s): none reported

CLINICAL POLICY

Melphalan Flufenamide

Appendix D: Withdrawal from Market

- Oncoceptes, the manufacturer of Pepaxto, voluntarily withdrew Pepaxto after data from the confirmatory Phase 3 OCEAN study revealed Pepaxto did not meet the requirements of the FDA Accelerated Approval regulation. The FDA Oncologic Drugs Advisory Committee review of the OCEAN study concluded the following:
 - The median overall survival was 19.7 months in the Pepaxto arm, compared to 25 months in the Pomalyst arm, HR 1.104 (95% CI 0.846, 1.441), indicating a safety concern.
 - The PFS results showed no statistical difference, with a HR 0.82 (95% CI 0.654, 1.027), indicating a lack of confirmed clinical benefit.
- Previously at the FDA's request, Oncoceptes stopped marketing Pepaxto in the US on October 22, 2021, and Pepaxto is currently not commercially available in the US but was available via the Individual Patient Expanded Access Investigational Drug Application (IND) process if deemed appropriate by the treating physician. At this same time Oncoceptes indicated that they planned to voluntarily withdraw Pepaxto, but later rescinded the withdrawal request and submitted additional analyses of the OCEAN study. This led to the September 2022 Oncologic Drugs Advisory Committee review that voted 14 to 2 that Pepaxto's benefit/risk profile was unfavorable.
- The Multiple Myeloma Research Foundation suggests those currently on Pepaxto therapy should contact their treating physician to see if remaining on therapy is appropriate.

V. Dosage and Administration

Indication	Dosing Regimen	Maximum Dose
Multiple myeloma	40 mg IV infusion on Day 1 of each 28-day treatment cycle, in combination with dexamethasone.	40 mg/dose

VI. Product Availability

Lyophilized powder in a single-dose vial for reconstitution and dilution for injection: 20 mg

VII. References

1. Pepaxto Prescribing Information. Waltham, MA: Oncoceptes Inc.; February 2021. Available at: https://www.accessdata.fda.gov/drugsatfda_docs/label/2021/214383s000lbl.pdf. Accessed ~~January 3, 2023~~ February 2, 2024.
2. National Comprehensive Cancer Network. Multiple Myeloma Version ~~3.2023~~ 1.2024. Available at: https://www.nccn.org/professionals/physician_gls/pdf/myeloma.pdf. Accessed ~~January 17, 2023~~ February 2, 2024.
3. National Comprehensive Cancer Network Drugs and Biologics Compendium. Available at: http://www.nccn.org/professionals/drug_compendium. Accessed ~~January 17, 2023~~ February 2, 2024.
4. Press release: Oncoceptes provides update on Pepaxto US marketing authorization. December 7, 2022. Available at: <https://www.oncoceptes.com/en/media/press->

CLINICAL POLICY

Melphalan Flufenamide

releases/oncopeptides-provides-update-on-pepaxto-us-marketing-authorization. Accessed ~~December 15, 2022~~ February 2, 2024.

5. FDA Oncologic Drugs Advisory Committee: FDA Presentations, NDA214383 – Pepaxto. September 23, 2022. Available at: <https://www.fda.gov/media/161761/download>. Accessed ~~December 15, 2022~~ February 2, 2024.

Coding Implications

Codes referenced in this clinical policy are for informational purposes only. Inclusion or exclusion of any codes does not guarantee coverage. Providers should reference the most up-to-date sources of professional coding guidance prior to the submission of claims for reimbursement of covered services.

HCPCS Codes	Description
J9247	Injection, melphalan flufenamide, 1 mg
<u>J9248</u>	<u>Injection, melphalan (hepzato), 1 mg</u>
<u>J9249</u>	<u>Injection, melphalan (apotex), 1 mg</u>

Reviews, Revisions, and Approvals	Date	LDH Approval Date
Policy created.	05.01.23	08.28.23
Annual review: no significant changes; references reviewed and updated.	03.25.24	<u>05.23.24</u>
<u>No significant changes; references reviewed and updated. Added HCPCS codes [J9248, J9249].</u>	<u>11.20.24</u>	

Important Reminder

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program approval status; evidence-based guidelines and positions of leading national health professional organizations; views of physicians practicing in relevant clinical areas affected by this clinical policy; and other available clinical information. LHCC makes no representations and accepts no liability with respect to the content of any external information used or relied upon in developing this clinical policy. This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved.

The purpose of this clinical policy is to provide a guide to medical necessity, which is a component of the guidelines used to assist in making coverage decisions and administering benefits. It does not constitute a contract or guarantee regarding payment or results. Coverage decisions and the administration of benefits are subject to all terms, conditions, exclusions, and limitations of the coverage documents (e.g., evidence of coverage, certificate of coverage, policy,

CLINICAL POLICY

Melphalan Flufenamide

contract of insurance, etc.), as well as to state and federal requirements and applicable LHCC administrative policies and procedures.

This clinical policy is effective as of the date determined by LHCC. The date of posting may not be the effective date of this clinical policy. This clinical policy may be subject to applicable legal and regulatory requirements relating to provider notification. If there is a discrepancy between the effective date of this clinical policy and any applicable legal or regulatory requirement, the requirements of law and regulation shall govern. LHCC retains the right to change, amend or withdraw this clinical policy, and additional clinical policies may be developed and adopted as needed, at any time.

This clinical policy does not constitute medical advice, medical treatment, or medical care. It is not intended to dictate to providers how to practice medicine. Providers are expected to exercise professional medical judgment in providing the most appropriate care, and are solely responsible for the medical advice and treatment of members. This clinical policy is not intended to recommend treatment for members. Members should consult with their treating physician in connection with diagnosis and treatment decisions.

Providers referred to in this clinical policy are independent contractors who exercise independent judgment and over whom LHCC has no control or right of control. Providers are not agents or employees of LHCC.

This clinical policy is the property of LHCC. Unauthorized copying, use, and distribution of this clinical policy or any information contained herein are strictly prohibited. Providers, members, and their representatives are bound to the terms and conditions expressed herein through the terms of their contracts. Where no such contract exists, providers, members and their representatives agree to be bound by such terms and conditions by providing services to members and/or submitting claims for payment for such services.

©2024 Louisiana Healthcare Connections. All rights reserved. All materials are exclusively owned by Louisiana Healthcare Connections and are protected by United States copyright law and international copyright law. No part of this publication may be reproduced, copied, modified, distributed, displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise published without the prior written permission of Louisiana Healthcare Connections. You may not alter or remove any trademark, copyright or other notice contained herein. Louisiana Healthcare Connections is a registered trademark exclusively owned by Louisiana Healthcare Connections.