

Clinical Policy: Sofosbuvir/Velpatasvir (Epclusa)

Reference Number: LA.PHAR.268

Effective Date: 07/16

Last Review Date: 05/18

Line of Business: Medicaid

Revision Log

See Important Reminder at the end of this policy for important regulatory and legal information.

Description

Sofosbuvir/velpatasvir (Epclusa®) is a fixed-dose combination of sofosbuvir, a hepatitis C virus (HCV) nucleotide analog NS5B polymerase inhibitor, and velpatasvir, an HCV NS5A inhibitor.

FDA-Approved Indication

Epclusa is indicated for the treatment of adult patients with chronic HCV genotype 1, 2, 3, 4, 5, or 6 infection:

- without cirrhosis or with compensated cirrhosis
- with decompensated cirrhosis for use in combination with ribavirin

Policy/Criteria

Provider must submit documentation (including office chart notes and lab results) supporting that member has met all approval criteria

It is the policy of health plans affiliated with Centene Corporation® that Epclusa is **medically necessary** when the following criteria are met:

I. Initial Approval Criteria**A. Chronic Hepatitis C Infection** (must meet all):

1. Diagnosis of chronic HCV infection as evidenced by multiple detectable HCV RNA (ribonucleic acid) levels in the last 6 months;
2. Confirmed HCV genotype is 1, 2, 3, 4, 5 or 6;
3. Age \geq 18 years;
4. Life expectancy \geq 12 months with HCV treatment;
5. Prescriber and patient must attest that the patient is not actively participating in illicit IV drug use and/or alcohol use;
6. Advanced liver disease defined as a or b:
 - a. Advanced fibrosis indicated by i, ii, or iii:
 - i. Liver biopsy showing a METAVIR score of F3 or equivalent (Knodell, Scheuer, Batts-Ludwig – F3; Ishak – F4/5);
 - ii. One serologic test showing an equivalent score to METAVIR F3 per Appendix C; or
 - iii. One radiologic test showing an equivalent score to METAVIR F3 per Appendix C;
 - b. Cirrhosis indicated by i, ii, iii, iv or v:
 - i. Hepatocellular carcinoma (HCC) - and the HCC is amenable to resection, ablation or transplant;

- ii. Liver biopsy showing a METAVIR score of F4 or equivalent (Knodell, Scheuer, Batts-Ludwig – F4; Ishak - F5/6);
 - iii. One serologic test showing an equivalent score to METAVIR F4 per Appendix C
 - iv. One radiologic test showing an equivalent score to METAVIR F4 per Appendix C or
 - v. other radiologic test showing evidence of cirrhosis (e.g., portal hypertension);
 - c. If member is HIV/HCV co-infected, there shall be no METAVIR score requirements.
- 7. Prescribed regimen is consistent with an FDA or AASLD-IDSA recommended regimen (*see Section V Dosage and Administration for reference*);
 - 8. If member is without cirrhosis or with compensated cirrhosis (Child-Pugh A): contraindication or intolerance to Mavyret;
 - 9. Member agrees to participate in a medication adherence program meeting both of the following components:
 - a. Medication adherence monitored by pharmacy claims data or member report;
 - b. Member's risk for non-adherence identified by adherence program or member/prescribing physician follow-up at least every 4 weeks;
 - 10. Member is hepatitis B virus (HBV) negative, or if positive, documentation that concurrent HBV infection is being treated (e.g., tenofovir alafenamide, adefovir, entecavir), unless contraindicated or clinically significant adverse effects are experienced (*see Appendix B*);
 - 11. Dose does not exceed sofosbuvir/velpatasvir 400 mg/100 mg (1 tablet) per day.
- Approval duration: up to 24 weeks***
(*Approved duration should be consistent with a regimen in Section V Dosage and Administration)

B. Other diagnoses/indications

- 1. Refer to CP.PHAR.57 if diagnosis is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized).

II. Continued Therapy

A. Chronic Hepatitis C Infection (must meet all):

- 1. Currently receiving medication via Centene benefit, or documentation supports that member is currently receiving Eplusa for chronic hepatitis C virus infection and has received this medication for at least 30 days;
- 2. Member is responding positively to therapy (e.g. decreased HCV RNA level, no unacceptable toxicity);
- 3. Dose does not exceed sofosbuvir/velpatasvir 400mg/100mg (1 tablet) per day.

Approval duration: up to a total of 24 weeks*

(*Approved duration should be consistent with a regimen in Section V Dosage and Administration)

B. Other diagnoses/indications

- 1. Refer to CP.PHAR.57 if diagnosis is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized).

III. Diagnoses/Indications for which coverage is NOT authorized:

- A. Non-FDA approved indications, which are not addressed in this policy, unless there is sufficient documentation of efficacy and safety according to the off label use policy – CP.PHAR.57 or evidence of coverage documents.

IV. Appendices/General Information

Appendix A: Abbreviation/Acronym Key

ALT: alanine aminotransferase

APRI: AST to platelet ratio

AASLD: American Association for the Study of Liver Diseases

FDA: Food and Drug Administration

FIB-4: Fibrosis-4 index

HBeAg: hepatitis B virus envelope antigen

HBV: hepatitis B virus

HCC: hepatocellular carcinoma

HCV: hepatitis C virus

IDSA: Infectious Diseases Society of America

MRE: magnetic resonance elastography

NS3/4A, NS5A/B: nonstructural protein

Peg-IFN: pegylated interferon

PI: protease inhibitor

RBV: ribavirin

RNA: ribonucleic acid

Appendix B: General Information

- Hepatitis B Reactivation is a black box warning for all direct-acting antiviral drugs for the treatment of HCV. The provider must provide either:
 - Documentation of absence of concurrent HBV infection as evidenced by laboratory values showing absence of hepatitis B virus envelope antigen (HBeAg) and HBV DNA;
 - Documentation that HBV co-infected patient may not be candidates for therapy as evidenced by one of the following:
 - Absence of HBeAg, HBV DNA less than 2,000 international units/mL, and alanine aminotransferase (ALT) level within 1 to 2 times the upper limit of normal;
 - HBeAg-positive and HBV DNA greater than 1,000,000 international units/mL and ALT level within 1 to 2 times the upper limit of normal;
 - Documentation that concurrent HBV infection is being treated (e.g., tenofovir alafenamide, adefovir, entecavir), unless contraindicated or clinically significant adverse effects are experienced
- The 2016 AASLD/IDSA treatment guideline for HBV consider ALT levels <30 U/L for men and <19 U/L for women as upper limits of normal.
- The 2016 AASLD/IDSA treatment guideline for HBV recommend adults with compensated cirrhosis, even with low levels of viremia (<2,000 IU/mL) be treated with antiviral therapy to reduce the risk of decompensation, regardless of ALT level. The recommendation extends to adults with decompensated cirrhosis be treated with antiviral therapy indefinitely regardless of HBV DNA level, HBeAg status, or ALT level to decrease the risk of worsening liver-related complications.

Appendix C: Approximate Scoring Equivalencies using METAVIR F3/F4 as reference

CLINICAL POLICY

Sofosbuvir/Velpatasvir

Fibrosis/ Cirrhosis	Serologic Tests*				Radiologic Tests†		Liver Biopsy‡	
	Fibro Test	FIBRO Spect II	APRI	FI B- 4	FibroScan (kPa)	MRE (kPa)	METAVIR	Ishak
Advanced fibrosis	≥0.59	≥42	>1.5	>3 .25	≥9.5	≥4.11	F3	F4-5
Cirrhosis	≥0.75	≥42	>1.5	>3 .25	≥12.0	≥4.71	F4	F5-6

***Serologic tests:**

FibroTest (available through Quest as FibroTest or LabCorp as FibroSure)

FIBROSpect II (available through Prometheus Laboratory)

APRI (AST to platelet ratio index)

FIB-4 (Fibrosis-4 index: includes age, AST level, platelet count)

†Radiologic tests:

FibroScan (transient elastography)

MRE (magnetic resonance elastography)

‡Liver biopsy (histologic scoring systems):

METAVIR F3/F4 is equivalent to Knodell, Scheuer, and Batts-Ludwig F3/F4 and Ishak F4-5/F5-6

METAVIR fibrosis stages: F0 = no fibrosis; F1 = portal fibrosis without septa; F2 = few septa;

F3 = numerous septa without cirrhosis; F4 = cirrhosis

Appendix D: Direct-Acting Antivirals for Treatment of HCV Infection

Brand Name	Drug Class				
	NS5A Inhibitor	Nucleotide Analog NS5B Polymerase Inhibitor	Non-Nucleoside NS5B Palm Polymerase Inhibitor	NS3/4A Protease Inhibitor (PI)	CYP3A Inhibitor
Daklinza	Daclatasvir				
Epclusa*	Velpatasvir	Sofosbuvir			
Harvoni*	Ledipasvir	Sofosbuvir			
Olysio				Simeprevir	
Sovaldi		Sofosbuvir			
Technivie*	Ombitasvir			Paritaprevir	Ritonavir
Viekira XR/PAK*	Ombitasvir		Dasabuvir	Paritaprevir	Ritonavir
Zepatier*	Elbasvir			Grazoprevir	

*Combination drugs

V. Dosage and Administration

Indication	Dosing Regimen	Maximum Dose	Reference
Genotype 1-6 CHC: Without cirrhosis or with compensated	One tablet PO QD for 12 weeks (GT 2 or 3 with compensated cirrhosis for Peg-IFN/RBV or	One tablet (sofosbuvir 400mg /velpatasvir 100mg) per day	1) FDA- approved labeling

CLINICAL POLICY
Sofosbuvir/Velpatasvir

cirrhosis, treatment-naïve or treatment- experienced patient	sofosbuvir-based treatment-experienced patient may use: one tablet PO QD with weight-based ribavirin for 12 weeks) †		2) AASLD- IDSA (updated 04/17)
Genotype 1-6 CHC : With decompensated cirrhosis treatment-naïve or treatment- experienced patient	One tablet PO QD plus weight-based RBV for 12 weeks (GT 1, 4, 5, or 6 with decompensated cirrhosis and RBV ineligible may use: one tablet PO QD for 24 weeks) †	One tablet (sofosbuvir 400mg /velpatasvir 100mg) per day	1) FDA- approved labeling 2) AASLD- IDSA (updated 04/17)

**AASLD/IDSA treatment guidelines for chronic hepatitis C infection are updated at irregular intervals; refer to the most updated AASLD/IDSA guideline for most accurate treatment regimen.*

VI. Product Availability

Tablet: sofosbuvir 400 mg / velpatasvir 100 mg

VII. References

1. Epclusa Prescribing Information. Foster City, CA: Gilead Sciences, Inc.; February 2017. Available at http://www.gilead.com/~media/files/pdfs/medicines/liver-disease/epclusa/epclusa_pi.pdf?la=en. Accessed May 10, 2017.
2. American Association for the Study of Liver Diseases/ Infectious Disease Society of America (AASLD-IDSA). Recommendations for testing, managing, and treating hepatitis C. Updated April 2017. <http://www.hcvguidelines.org>. Accessed May 10, 2017.
3. Bonder A, Afdhal N. Utilization of FibroScan in clinical practice. Curr Gastroenterol Rep. 2014; 16(372): 1-7. DOI 10.1007/s11894-014-0372-6.
4. Halfon P, Bourliere M, Deydier R, et al. Independent prospective multicenter validation of biochemical markers (Fibrotest–Actitest) for the prediction of liver fibrosis and activity in patients with chronic hepatitis C: The Fibropaca study. Am J Gastroenterol. 2006; 101: 547-555. DOI: 10.1111/j.1572-0241.2006.04111.x
5. Hepatitis C Virus (HCV) FibroSure. Laboratory Corporation of America Holdings and Lexi-Comp, Inc. Available at <https://www.labcorp.com>. 2016. Accessed July 15, 2016.
6. Hepatitis C Virus (HCV) FibroTest-ActiTest Panel. Nichols Institute/Quest Diagnostics. Available at http://education.questdiagnostics.com/physician_landing_page. 2016. Accessed July 15, 2016.
7. Hepatitis C Virus (HCV) FIBROSpect II. Prometheus Therapeutics and Diagnostics. Available at http://www.prometheuslabs.com/Resources/Fibrospect/Fibrospect_II_Product_Detail_Sheet_FIB16005_04-16.pdf. April 2016. Accessed July 15, 2016.

CLINICAL POLICY
Sofosbuvir/Velpatasvir

8. Hsieh YY, Tung SY, Lee K, et al. Routine blood tests to predict liver fibrosis in chronic hepatitis C. *World J Gastroenterol*. February 28, 2012; 18(8): 746-53. doi: 10.3748/wjg.v18.i8.746.
9. Heimbach J, Kulik LM, Finn R, et al. AASLD Guidelines for the Treatment of Hepatocellular Carcinoma. *Hepatology* 2017.
10. Terrault NA, Bzowej NH, Chang KM, et al. American Association for the Study of Liver Diseases (AASLD) Guidelines for Treatment of Chronic Hepatitis B. *Hepatology* 2016; 62(1): 261-283.

Reviews, Revisions, and Approvals	Date	Approval Date
New policy created, split from CP.PHAR.17 Hepatitis C Therapies policy. HCV RNA levels over six-month period added to confirm infection is chronic. Life expectancy “≥12 months if HCC and awaiting transplant” is modified to indicate “≥12 months with HCV therapy.” Methods to diagnose fibrosis/cirrhosis are modified to require a liver biopsy or a combination of one serologic and one radiologic test. Serologic and radiologic tests are updated and correlated with METAVIR per Appendix C. Dosing regimens are presented in Appendix. Criteria is added requiring a verification of HCV RNA status at 4 weeks (and again at 6 weeks if present at 4) accordingly, the initial approval period is shortened to 8 weeks.	07/16	07/16
Edited policy so congruent with the other HCV policies as follows: Testing criteria reorganized by cirrhosis status consistent with the regimen tables; HCC population broadened to incorporate those amenable to curative measures (resection, ablation, transplant). Fibrosure test that meets F3 requirement changed to ≥ 0.59 . Criteria added excluding post-liver transplantation unless regimens specifically designate. Preferencing language edited for clarity. Removed creatinine clearance restriction. Under continuing approval, presence of HCV RNA is edited to remove specific timing of testing. Appendix B edited for clarity; Appendix C added. Appendix D – genotype “1” is footnoted to clarify possible subtypes. “Includes HCC” is removed from the decompensated cirrhosis. “Daily” is removed from the “recommended regimen” column; presentation of other data is abbreviated/short-handed.	08/16	09/16

CLINICAL POLICY
Sofosbuvir/Velpatasvir

Reviews, Revisions, and Approvals	Date	Approval Date
Policy converted to new template. Added requirement for prevention of HBV reactivation. Consolidated appendix D and E into dosing and administration in section V, added maximum dose requirement, initial approval duration expanded to full 12 weeks, limited continued therapy approval duration to 12 weeks, deleted viral load and adherence requirement in continued therapy, added documentation of positive response to therapy and continuity of care, and removed CIs in section II, added reference column in section V. Added preferencing information requiring Mavyret for FDA-approved indications. Safety criteria was applied according to the safety guidance discussed at CPAC and endorsed by Centene Medical Affairs. Exception made to require Hep B screening for all patients prior to treatment to ensure that proper risk reduction measures are taken.	08/17	09/17
Due to State requirements, removed prescriber restrictions regarding who can prescribe Hepatitis C DAA agents; Removed the abstinence period of 6 months and added documentation required that member is not actively participating in alcohol and/or illicit IV drugs use,; Added statement for NO Fibrosis score required for HIV/HCV co-infected members	4/18	4/18
Changed language to one serologic test OR one radiologic test from one serologic test and one radiologic test	5/18	5/18

Important Reminder

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program approval status; evidence-based guidelines and positions of leading national health professional organizations; views of physicians practicing in relevant clinical areas affected by this clinical policy; and other available clinical information. The Health Plan makes no representations and accepts no liability with respect to the content of any external information used or relied upon in developing this clinical policy. This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved. "Health Plan" means a health plan that has adopted this clinical policy and that is operated or administered, in whole or in part, by Centene Management Company, LLC, or any of such health plan's affiliates, as applicable.

CLINICAL POLICY**Sofosbuvir/Velpatasvir**

The purpose of this clinical policy is to provide a guide to medical necessity, which is a component of the guidelines used to assist in making coverage decisions and administering benefits. It does not constitute a contract or guarantee regarding payment or results. Coverage decisions and the administration of benefits are subject to all terms, conditions, exclusions and limitations of the coverage documents (e.g., evidence of coverage, certificate of coverage, policy, contract of insurance, etc.), as well as to state and federal requirements and applicable Health Plan-level administrative policies and procedures.

This clinical policy is effective as of the date determined by the Health Plan. The date of posting may not be the effective date of this clinical policy. This clinical policy may be subject to applicable legal and regulatory requirements relating to provider notification. If there is a discrepancy between the effective date of this clinical policy and any applicable legal or regulatory requirement, the requirements of law and regulation shall govern. The Health Plan retains the right to change, amend or withdraw this clinical policy, and additional clinical policies may be developed and adopted as needed, at any time.

This clinical policy does not constitute medical advice, medical treatment or medical care. It is not intended to dictate to providers how to practice medicine. Providers are expected to exercise professional medical judgment in providing the most appropriate care, and are solely responsible for the medical advice and treatment of members. This clinical policy is not intended to recommend treatment for members. Members should consult with their treating physician in connection with diagnosis and treatment decisions.

Providers referred to in this clinical policy are independent contractors who exercise independent judgment and over whom the Health Plan has no control or right of control. Providers are not agents or employees of the Health Plan.

This clinical policy is the property of the Health Plan. Unauthorized copying, use, and distribution of this clinical policy or any information contained herein are strictly prohibited. Providers, members and their representatives are bound to the terms and conditions expressed herein through the terms of their contracts. Where no such contract exists, providers, members and their representatives agree to be bound by such terms and conditions by providing services to members and/or submitting claims for payment for such services.

Note: For Medicaid members, when state Medicaid coverage provisions conflict with the coverage provisions in this clinical policy, state Medicaid coverage provisions take precedence. Please refer to the state Medicaid manual for any coverage provisions pertaining to this clinical policy.

©2016 Centene Corporation. All rights reserved. All materials are exclusively owned by Centene Corporation and are protected by United States copyright law and international copyright law. No part of this publication may be reproduced, copied, modified, distributed, displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise published without the prior written permission of Centene Corporation. You may not alter or

CLINICAL POLICY
Sofosbuvir/Velpatasvir



remove any trademark, copyright or other notice contained herein. Centene® and Centene Corporation® are registered trademarks exclusively owned by Centene Corporation.