

Appendix J - Performance Measure Reporting

Identifier	Measure	Measure Description	Measure Steward	Federal Reporting Program	Target Population	Condition	Data
TBD \$\$	Antenatal Progesterone	TBD	DHH/UIM	None	Children's and Maternal Health	Perinatal and Reproductive Health	HP
NQF #0471 (CSEC) \$\$	Cesarean Rate for Low-Risk First Birth Women	The percentage of cesareans in live births at or beyond 37.0 weeks gestation to women that are having their first delivery and are singleton (no twins or beyond) and are vertex presentation (no breech or transverse positions).	TJC	CHIPRA	Children's and Maternal Health	Perinatal and Reproductive Health	HP
(BHRA)	Behavioral Health Risk Assessment (for Pregnant Women)	Percentage of women, regardless of age, that gave birth during a 12-month period seen at least once for prenatal care who received a behavioral health screening risk assessment that includes the following screenings at the first prenatal visit: depression, alcohol use, tobacco use, drug use, and intimate partner violence.	AMA-PCPI	CHIPRA	Children's and Maternal Health	Perinatal and Reproductive Health	State
NQF #1391 (FPC)	Frequency of Ongoing Prenatal Care	Percentage of Medicaid deliveries between November 6 of the year prior to the measurement year and November 5 of the measurement year that received the following percentages of expected prenatal visits: <21, 21-40, 41-60, 61-80, > or = 80.	NCQA	CHIPRA	Children's and Maternal Health	Perinatal and Reproductive Health	HP
NQF #1382 (LBW)	Percentage of low birth weight births	Percentage of live births that weighted less than 2,500 grams in the state during the reporting period.	CDC	CHIPRA, HRSA	Children's and Maternal Health	Perinatal and Reproductive Health	OPH
NQF #1517 (PPC)	Timeliness of Prenatal Care	The percentage of deliveries of live births between November 6 of the year prior to the measurement year and November 5 of the measurement year. The percentage of deliveries that received a prenatal care visit as a patient of the organization in the first trimester or within 42 days of enrollment in the organization.	NCQA	CHIPRA	Children's and Maternal Health	Perinatal and Reproductive Health	HP
(AWC) \$\$	Adolescent Well Care Visit	The percentage of enrolled members 12-21 years of age who had at least one comprehensive well-care visit with a PCP or an OB/GYN practitioner during the measurement Year	NCQA	CHIPRA	Children's Health	Utilization	HP
(CAP)	Child and Adolescents' Access to Primary Care Practitioners	Percentage of children ages 3 to 17 that had an outpatient visit with a primary care practitioner (PCP) or obstetrical/gynecological (OB/GYN) practitioner and whose weight is classified based on body mass index percentile for age and gender.	NCQA	CHIPRA	Children's Health	Access/Availability of Care	HP
NQF #0038 (CIS)	Childhood Immunization Status	Percentage of children that turned 2 years old during the measurement year and had specific vaccines by their second birthday.	NCQA	CHIPRA, MU2	Children's Health	Prevention	HP
NQF #1448 (DEV)	Developmental Screening in the First Three Years of Life	The percentage of children screened for risk of developmental, behavioral and social delays using a standardized screening tool in the first three years of life. This is a measure of screening in the first three years of life that includes three, age-specific indicators assessing whether children are screened by 12 months of age, by 24 months of age and by 36 months of age.	NCQA	CHIPRA	Children's Health	Prevention	State
NQF # 0108 \$\$	Follow-up Care for Children Prescribed ADHD Medication	The percentage of children newly prescribed attention-deficit/hyperactivity disorder (ADHD) medication who had at least three follow-up care visits within a 10-month period, one of which was within 30 days of when the first ADHD medication was dispensed.	NCQA	CHIPRA, MU2	Children's Health	Behavioral Health	HP

NQF #1959 (HPV)	Human Papillomavirus (HPV) Vaccine for Female Adolescents	Percentage of female adolescents that turned 13 years old during the measurement year and had three doses of the human papillomavirus (HPV) vaccine by their 13th birthday.	NCQA	CHIPRA	Children's Health	Prevention	HP
NQF #1407 (IMA)	Immunization Status for Adolescents	Percentage of adolescents that turned 13 years old during the measurement year and had specific vaccines by their 13th birthday.	NCQA	CHIPRA	Children's Health	Prevention	HP
NQF #0024 (WCC)	Weight Assessment and Counseling for Nutrition and Physical Activity for Children/Adolescents: Body Mass Index Assessment for Children/Adolescents	Percentage of children ages 3 to 17 that had an outpatient visit with a primary care practitioner (PCP) or obstetrical/gynecological (OB/GYN) practitioner and whose weight is classified based on body mass index percentile for age and gender.	NCQA	CHIPRA, MU2	Children's Health	Prevention	HP
NQF #1392 (W15)	Well-Child Visits in the First Fifteen Months of Life	Percentage of patients who turned 15 months old during the measurement year and who had the following number of well-child visits with a PCP during their first 15 months of life. Seven rates are reported.	NCQA	CHIPRA	Children's Health	Utilization	HP
NQF #1516 (W34)	Well-Child Visits in the Third, Fourth, Fifth and Sixth Years of Life	Percentage of patients 3–6 years of age who received one or more well-child visits with a PCP during the measurement year.	NCQA	CHIPRA	Children's Health	Utilization	HP
NQF #0021 (MPM)	Annual Monitoring for Patients on Persistent Medications	The percentage of members 18 years of age and older who received at least 180 treatment days of ambulatory medication therapy for a select therapeutic agent during the measurement year and at least one therapeutic monitoring event for the therapeutic agent in the measurement year. For each product line, report each of the four rates separately and as a total rate.	NCQA	MEDICAID ADULT	Chronic Disease	Prevention	HP
NQF #0057 (HA1C)	Comprehensive Diabetes Care: Hemoglobin A1c testing	The percentage of members 18-75 years of age with diabetes (type 1 and type 2) who received an HbA1c test during the measurement year.	NCQA	MEDICAID ADULT	Chronic Disease	Diabetes	HP
NQF #0063 (LDL)	Comprehensive Diabetes Care: LDL-C Screening	The percentage of members 18-75 years of age with diabetes (type 1 and type 2) who received an LDL-C test during the measurement year.	NCQA	MEDICAID ADULT, MU2	Chronic Disease	Diabetes	HP
NQF #0018 (CBP)	Controlling High Blood Pressure	The percentage of patients 18 to 85 years of age who had a diagnosis of hypertension (HTN) and whose blood pressure (BP) was adequately controlled (<140/90) during the measurement year.	NCQA	MEDICAID ADULT, MU2, CMS HEALTH HOMES	Chronic Disease	Cardiovascular Care	HP
NQF #0277 (PQ108)	Heart Failure Admission Rate	Percent of population with an admissions for heart failure (reported by Recipient Parish)	AHRQ	MEDICAID ADULT	Chronic Disease	Cardiovascular Care	State
NQF #2082 (HIV) \$\$	HIV Viral Load Suppression	Percentage of patients, regardless of age, with a diagnosis of HIV with a HIV viral load less than 200	HRSA HIV/AIDS Bureau	MEDICAID ADULT	Chronic Disease	HIV	HP
NQF #0272 (PQ1 1) \$\$	Diabetes Short Term Complications Rate	Number of discharges for diabetes short term complications per 100,000 Medicaid enrollees age 18 and older.	AHRQ	MEDICAID ADULT	Chronic Disease	Diabetes	State
NQF #0476 (PC-03)	Antenatal Steroids	This measure assesses patients at risk of preterm delivery at >=24 and <32 weeks gestation receiving antenatal steroids prior to delivering preterm newborns.	TJC	MEDICAID ADULT	Maternal Health	Perinatal and Reproductive Health	HP
NQF #0469 (PC-01)	Elective Delivery	This measure assesses patients with elective vaginal deliveries or elective cesarean sections at >= 37 and < 39 weeks of gestation completed	TJC	MEDICAID ADULT, MU2	Maternal Health	Perinatal and Reproductive Health	HP
NQF # 1517 (PPC) \$\$	Postpartum Care (PPC Submeasure)	The percentage of deliveries that had a postpartum visit on or between 21 and 56 days after delivery.	NCQA	MEDICAID ADULT	Maternal Health	Perinatal and Reproductive Health	HP

NQF #1879 (SAA)	Adherence to Antipsychotic Medications for Individuals with Schizophrenia	The measure calculates the percentage of individuals 18 years of age or greater as of the beginning of the measurement period with schizophrenia or schizoaffective disorder who are prescribed an antipsychotic medication, with adherence to the antipsychotic medication [defined as a Proportion of Days Covered (PDC)] of at least 0.8 during the measurement period (12 consecutive months).	CMS	MEDICAID ADULT	Population Health	Behavioral Health	HP
(ABA)	Adult BMI Assessment	The percentage of members 18-74 years of age who had an outpatient visit and whose body mass index (BMI) was documented during the measurement or the year prior to the measurement year.	NCQA	MEDICAID ADULT, CMS HEALTH HOMES	Population Health	Prevention	HP
(AMB) \$\$	Ambulatory Care	Utilization of ambulatory care. Outpatient and ED Visits per 1000 member months	NCQA	CHIPRA	Population Health	Utilization	State
NQF #0105 (AMM)	Antidepressant Medication Management	The percentage of members 18 years of age and older with a diagnosis of major depression and were newly treated with antidepressant medication, and who remained on an antidepressant medication treatment. Two rates are reported.	NCQA	MEDICAID ADULT, MU2	Population Health	Behavioral Health	HP
NQF #0283 (PQI15)	Asthma in Younger Adults Admission Rate	Admissions for a principal diagnosis of asthma per 100,000 population, ages 18 to 39 years. Excludes admissions with an indication of cystic fibrosis or anomalies of the respiratory system, obstetric admissions, and transfers from other institutions.	AHRQ	MEDICAID ADULT	Population Health	Pulmonary/Critical Care	State
NQF #1800 (AMR)	Asthma Medication Ratio	The percentage of patients 5–64 years of age who were identified as having persistent asthma and had a ratio of controller medications to total asthma medications of 0.50 or greater during the measurement year.	NCQA	None - recommended by IPRO	Population Health	Pulmonary/Critical Care	HP
NQF #0648 (CTR)	Care Transition - Transition Record Transmitted to Health Care Professional	Percentage of patients, regardless of age, discharged from an inpatient facility (eg, hospital inpatient or observation, skilled nursing facility, or rehabilitation facility) to home or any other site of care for whom a transition record was transmitted to the facility or primary physician or other health care professional designated for follow-up care within 24 hours of discharge	AMA-PCPI	MEDICAID ADULT, CMS HEALTH HOMES	Population Health	Care Coordination	HP
NQF #0032 (CCS)	Cervical Cancer Screening	Percentage of women 21–64 years of age who were screened for cervical cancer	NCQA	MEDICAID ADULT, MU2	Population Health	Prevention	HP
NQF #0275 (PQI05)	COPD and Asthma in Younger Adults Admission Rate	This measure is used to assess the number of admissions for chronic obstructive pulmonary disease (COPD) per 100,000 population.	AHRQ	MEDICAID ADULT	Population Health	Pulmonary/Critical Care	State
NQF #0039 (FVA)	Flu Vaccinations for Adults Ages 18 to 64	The percentage of adults 18 years of age and older who self-report receiving an influenza vaccine within the measurement period.	NCQA	MEDICAID ADULT	Population Health	Prevention	HP Survey
NQF #0576 (FUH)	Follow up After Hospitalization for Mental Illness	The percentage of discharges for patients 6 years of age and older who were hospitalized for treatment of selected mental illness diagnoses and who had an outpatient visit, an intensive outpatient encounter or partial hospitalization with a mental health practitioner. Two rates are reported	NCQA	CHIPRA, MEDICAID ADULT, CMS HEALTH HOMES	Population Health	Behavioral Health	HP

NQF #0004 (IET)	Initiation and Engagement of Alcohol and Other Drug Dependence Treatment	The percentage of adolescent and adult patients with a new episode of alcohol or other drug (AOD) dependence who received the following. Initiation of AOD Treatment. Engagement of AOD Treatment. The percentage of patients who initiated treatment and who had two or more additional services with a diagnosis of AOD within 30 days of the initiation visit. The percentage of patients who initiate treatment through an inpatient AOD admission, outpatient visit, intensive outpatient encounter or partial hospitalization within 14 days of the diagnosis.	NCQA	MEDICAID ADULT, MU2, CMS HEALTH HOMES	Population Health	Behavioral Health	HP
NQF #0027 (MSC)	Medical Assistance With Smoking and Tobacco Use Cessation	Assesses different facets of providing medical assistance with smoking and tobacco use cessation	NCQA	MEDICAID ADULT	Population Health	Prevention	HP Survey
NQF #1799 (MMA)	Medication Management for People with Asthma	The percentage of patients 5-64 years of age during the measurement year who were identified as having persistent asthma and were dispensed appropriate medications that they remained on during the treatment period. Two rates are reported.	NCQA	CHIPRA	Population Health	Pulmonary/Critical Care	HP
(PCR)	Plan All-Cause Readmission Rate	For members 18 years of age and older, the number of acute inpatient stays during the measurement year that were followed by an acute readmission for any diagnosis within 30 days and the predicted probability of an acute readmission.	NCQA	MEDICAID ADULT, CMS HEALTH HOMES	Population Health	Utilization	HP
NQF #0418 (CDF)	Screening for Clinical Depression and Follow-Up Plan	Percentage of patients aged 12 years and older screened for clinical depression using an age appropriate standardized tool AND follow-up plan documented	CMS	MEDICAID ADULT, MU2, CMS HEALTH HOMES	Population health	Prevention	State
(CAT)	Call Answer Timeliness	Percentage of calls received by the organization's Member Services call centers (during operating hours) during the performance measurement year that were answered by a live voice within 30 seconds.	NCQA	None	Population Health	Access/Availability of Care	HP
(ACCA)	Ambulatory Care-Sensitive Condition Admission	This measure is used to assess the age-standardized acute care hospitalization rate for conditions where appropriate ambulatory care prevents or reduces the need for admission to the hospital per 100,000 population under age 75 years.	CIHI	CMS HEALTH HOMES	Population Health	Access/Availability of Care	HP
NQF #0033 (CHL)	Chlamydia Screening in Women	The percentage of women 16-24 years of age who were identified as sexually active and who had at least one test for chlamydia during the measurement year.	NCQA	CHIPRA, MEDICAID ADULT	Population Health, Maternal Health	Perinatal and Reproductive Health, Sexually Transmitted Infectious Diseases	HP
NQF #0031 (BCS)	Breast Cancer Screening	Percentage of women 40-69 years of age who had a mammogram to screen for breast cancer	NCQA	MEDICAID ADULT, MU2	Senior Care	Prevention	HP

Administrative Measures

Measure	Minimal Performance Standard
% of PCP Practices that provide verified 24/7 phone access with ability to speak with a PCP Practice clinician (MD, DO, NP, PA, RN, LPN) within 30 minutes of member contact.	≥95%
% of regular and expedited service authorization request processed in timeframes in the contract.	≥95%
Rejected claims returned to provider with reason code within 15 days of receipt of claims submission.	≥99%
% of Call Center calls answered within 30 seconds.	≤95%
Call Center average speed of answer.	30 Seconds
Call Center call abandonment rate.	≤5%
% of grievances and request for appeals received by the MCO including grievances received via telephone and resolved within the timeframe of the contract.	≥95%
% of clean claims paid for each provider type with 15 business days.	≥90%
% of clean claims paid for each provider type within 30 calendar days.	≥99%