

## AmeriHealth Caritas Louisiana

National Imaging Associates, Inc.*	
Clinical guidelines <b>LOWER EXTREMITY CT</b> (Foot, Ankle, Knee, Leg or Hip CT)	Original Date: September 1997
CPT Codes: 73700, 73701, 73702	Last Revised Date: <del>May-May</del> 2021 <sup>10</sup>
Guideline Number: NIA_CG_057-2	Implementation Date: January 2021 <sup>11</sup>

**INDICATIONS FOR LOWER EXTREMITY CT (FOOT, ANKLE, KNEE, LEG or HIP):**  
(Plain radiographs must precede CT evaluation)

Some indications are for MRI, CT, or MR or CT Arthrogram. More than one should not be approved at the same time.

If a CT Arthrogram fits approvable criteria below, approve as CT

**Joint-specific provocative orthopedic examination** when MRI is contraindicated or cannot be performed ([see Table 1](#)).<sup>1</sup>

**Note:** With a positive orthopedic sign, an initial x-ray is always preferred. However, it is not required to approve advanced imaging.

● Ankle

●

○ Unstable syndesmotic injury (high ankle injury)

- With inconclusive stress x-rays (a standing CT is preferred)
- Can have positive fibular translation, squeeze or cotton test, but imaging may be needed to confirm diagnosis
- Positive drawer sign ≥ 4 days after injury (Vuurberg, 2018)

- Knee ([Bennett, 2012](#); Doral, 2018; [Fox, 2018](#); Katz, 2013; Mohankumar, 2014; Slaughter, 2014; Smith, 2015; [TuiteTaljanovic, 2019](#)<sup>14</sup>)

○ Any positive test listed

- McMurray's
- Thessaly
- Apley's
- Lachman's
- Anterior or Posterior Drawer sign

\* National Imaging Associates, Inc. (NIA) is a subsidiary of Magellan Healthcare, Inc.

- Varus or valgus stress
- Acute mechanical locking of the knee not due to guarding (Hussin, 2014)
- Hip
  - Anterior Impingement sign (labral tear)-(Hananouchi, 2012; Naraghi, 2015; [Ross, 2018](#)~~Ward, 2013~~)
- Posterior Impingement sign (labral tear)-(Groh, 2009)

**Joint or muscle pain without positive findings on an orthopedic exam as listed above, after x-ray completed and an MRI is contraindicated or cannot be performed - (does not apply to young children)**

(Katz, 2013; Mordecai, 2014) ~~(does not apply to young children)~~

- Persistent joint or musculotendinous pain unresponsive to conservative treatment\*, within the last 6 months which includes active medical therapy (physical therapy, chiropractic treatments, and/or physician-supervised exercise\*\*) of at least four (4) weeks, **OR**
- With progression or worsening of symptoms during the course of conservative treatment
- Persistent hip mechanical symptoms including clicking, locking, catching, giving way or hip instability with a clinical suspicion of labral tear, with or without clinical findings suggestive of impingement-(Groh, 2009; Mintz, 2017)

**Ankle instability and suspected anterior talofibular ligament rupture (anterior and posterior drawer tests) as a result of a sprain requires initial active conservative therapy (above) and x-ray**

**Painful acquired or congenital flatfoot deformity in an adult, after x-ray completed and MRI is contraindicated**

- After failure of active conservative therapy listed above (Abousayed, 2017; Thorpe, 2012)

### **Extremity Mass**

- Mass or lesion after non-diagnostic x-ray or ultrasound (Murphey, 2018) and MRI cannot be performed. CT is better than MRI to evaluate mass calcification or bone involvement and may complement or replace MRI (Subhawong, 2010)
  - Baker's cyst should be initially evaluated with ultrasound
  - If superficial, then ultrasound is the initial study
  - If deep, then x-ray is the initial study
  - If there is a contraindication to MRI

### **Known Cancer of the Extremity**

([Bestic, 2019](#); Fitzgerald, 2015; Holzapfel, 2015; Kircher, 2012; ~~Morrison, 2013~~; [NCCN, 2019](#))

- Cancer staging
- Cancer Restaging
- Signs or symptoms of recurrence

### **Infection of Bone or Joint**

(Dodwell, 2013; Glaudemans, 2019)

**Note:** MRI and nuclear medicine studies are recommended for acute infection as they are more sensitive in detecting early changes of osteomyelitis (Mandell, 2017). CT is better at demonstrating findings of chronic osteomyelitis (sequestra, involucrum, cloaca, sinus tracts) as well as detecting soft tissue gas and foreign bodies (Fayad, 2007).

- Abnormal x-ray or ultrasound
- Negative x-ray but with a clinical suspicion of infection
  - Signs and symptoms of joint or bone infection include:
    - Pain and swelling
    - Decrease range of motion
    - Fevers
  - Laboratory findings of infection include:
    - Elevated ESR or CRP
    - Elevated white blood cell count
    - Positive joint aspiration
- Ulcer (diabetic, pressure, ischemic, traumatic) with signs of infection (redness, warm, swelling, pain, discharge which may range from white to serosanguineous) that is not improving despite treatment and bone or deep infection is suspected
  - Increased suspicion if size or temperature increases, bone is exposed/positive probe-to-bone test, new areas of breakdown, new smell (Bowers, 2020)
- Neuropathic foot with friable or discolored granulation tissue, foul odor, non-purulent discharge, and delayed wound healing (Pitocco, 2019)
- ~~A suspicion of osteomyelitis of the foot in patients with diabetes (Walker, 2019)~~

**Osteonecrosis (Avascular necrosis (AVN), Legg-Calve-Perthes Disease)** when MRI is contraindicated or cannot be performed  
(Felten, 2019; Murphey, 2014; 2016)

- Abnormal x-ray
- Normal or indeterminate x-rays but symptomatic and high risk (e.g., glucocorticosteroid use, renal transplant recipient, glycogen storage disease, alcohol abuse (Fukushima, 2010), sickle cell anemia (Wali, 2011))

**For evaluation of known or suspected autoimmune disease (e.g., rheumatoid arthritis) and MRI is contraindicated:**

-(Colebatch, 2013)

- Further evaluation of an abnormality or non-diagnostic findings on prior imaging
- Initial imaging of a single joint for diagnosis or response to therapy after plain films and appropriate lab tests (e.g., RF, ANA, CRP, ESR) (e.g., RF, ANA, CRP, ESR) (Colebatch, 2013).
- To determine change in treatment or when diagnosis is uncertain prior to start of treatment
- Follow-up to determine treatment efficacy of early rheumatoid arthritis
- Follow-up to determine treatment efficacy of advanced rheumatoid arthritis if x-ray and ultrasound are equivocal or noncontributory

- 

### Crystalline Arthropathy

- Dual-energy CT can be used to characterize crystal deposition disease, such as gout versus CPPD (Chou, 2017)

### Trauma

#### Bone Fracture

- Suspected stress or insufficiency fracture with a negative initial x-ray (Berncardino, 2017; Patel, 2011; Sadineni, 2015):
  - If hips and MRI cannot be done
  - If other parts of the extremities and repeat x-rays in 10-14 days are negative or non-diagnostic
  - If at high risk for a complete fracture with conservative therapy (e.g., navicular bone) ~~and~~ MRI cannot be ~~done~~ performed (Kellar, 2020)
- Suspected acute hip fracture with initial x-rays negative or non-diagnostic (Gill, 2013; Ross, 2019)
- Intra articular fractures that may require surgery (i.e., depressed tibial plateau fracture) (Prat-Fabregat, 2017)
- Nonunion or delayed union as demonstrated by no healing between two sets of x-rays. If a fracture has not healed by 4-6 months, there is delayed union. Incomplete healing by 6-8 months is nonunion (Morshed, 2014; Salih, 2015)

#### Tendon or Muscle Rupture after X-Ray and MRI is contraindicated or cannot be

#### ~~done~~performed

-(Garra, 2012; Peck [J](#), 2017; Wilkins, 2012)

- Clinical suspicion based on mechanism of injury and physical findings

**Suspected ACL Rupture** - Acute knee injury with physical exam limited by pain and swelling with x-ray completed (Wheless, 2018) if MRI is contraindicated (Taljanovic, 2019)

- Inability to perform because of pain and swelling should be considered a red flag
- Suspicion should be based on mechanism of injury, i.e., twisting, blunt force
- Normal x-ray:
  - Extreme pain, inability to stand, audible pop at time of injury, very swollen joint, leg numbness
- Abnormal x-ray:
  - Large joint effusion on x-ray knee effusion (Cecava, 2018)

**Osteochondral Lesions** (defects, fractures, osteochondritis dissecans) and x-ray done (if MRI contraindicated or cannot be done)

(Mintz 2017; Smith, 2012; ~~Tuite, 2014~~; [Taljanovic, 2019](#); Van Bergen, 2016<sup>5</sup>; Van Dijk, 2010)

- Clinical suspicion based on mechanism of injury and physical findings

## Foreign Body

(Laya, 2017)

- Indeterminate x-ray and ultrasound

## Loose bodies or synovial chondromatosis seen on x-ray or ultrasound

- In the setting of joint pain (Rajani, 2016)

**Peripheral Nerve Entrapment (e.g., tarsal tunnel, Morton's neuroma-etc) when and** MRI is contraindicated, including any of the following:

(Domkundwar, 2017; Dong 2012; Donovan, 2010; Tos, 2015)

- Abnormal Electromyogram or Nerve conduction study
- Abnormal x-ray or ultrasound
- Clinical suspicion and failed 4 weeks conservative treatment including at least 2 of the following (active treatment with physical therapy is not required):
  - Activity modification
  - Rest, ice, or heat
  - Splinting or orthotics
  - Medication

## Pediatrics:

**Note:** Leg length discrepancy – the literature indicates that standing plain film x-rays are preferred, but there are some advantages to using a CT scanogram instead and may be preferred

(Guggenberger, 2014; Sabharwal, 2008)

- Osteoid Osteoma after an x-ray is done (Iyer, 2012)
- Painful flatfoot (Pes planus) deformity with suspected tarsal coalition, not responsive to active conservative care (Bouchard, 2014)
  - When MRI cannot be ~~done~~**performed**; **OR**
  - Extra-articular coalition is suspected (bony bridges around the joints); **OR**
  - When needed for surgical planning (Glaser, 2016)
- Slipped Capital Femoral Epiphysis and Chronic Recurrent Multifocal Osteomyelitis – MRI is the appropriate modality, rather than CT

## Pre-operative/procedural evaluation:

- Pre-operative evaluation for a planned surgery or procedure if the imaging provides diagnostic information that is not available on prior studies (See exclusions provider should be referred to the health plan for nondiagnostic surgical planning studies\*)

•

\*CT or MRI requests are not approvable for the following total knee arthroplasty (TKA) procedures: Procedures utilizing computer-navigated or patient-specific or gender-specific instrumentation (Johnson, 2011)

Bicompartmental arthroplasty (investigational at this time) (Dudhniwala, 2016)

~~Note: These surgical procedures are not considered a covered service and are not reimbursable based on lack of current scientific evidence for clinically important improvement, safety or efficacy; or based on scientific evidence of increased risk of serious complications.~~

**Post-operative/procedural evaluation:**

- When imaging, physical, or laboratory findings indicate joint infection, delayed or non-healing, or other surgical/procedural complications
- Joint prosthesis loosening or dysfunction, x-rays non-diagnostic (Fritz, 2014, 2015)
- Trendelenburg sign or other indication of muscle or nerve damage after recent hip surgery



**Table 1: Positive Orthopedic Joint Tests, Lower Extremity**

---

**ANKLE**

~~Posterior draw~~

Fibular translation

Squeeze

Cotton

Thompson

Thumb squeeze test

Mulder click

**HIP**

~~Trendelenburg~~

**KNEE**

Anterior draw

Pivot Shift Test

Lachman

Posterior tibial Sag

Posterior Draw

McMurray's Test

Thessaly

Valgus stress

Varus stress

Ege

---

**BACKGROUND:**

Plain radiographs are typically used as the first-line modality for assessment of lower extremity conditions. Computed tomography (CT) is used for evaluation of tumors, metastatic lesions, infection, fractures, and other problems. Magnetic resonance imaging (MRI) is the first-line choice for imaging of many conditions, but CT may be used in these cases if MRI is contraindicated or unable to be performed.

**OVERVIEW:**

**\*Conservative Therapy:** (musculoskeletal) should include a multimodality approach consisting of a combination of active and inactive components. Inactive components such as rest, ice, heat, modified activities, medical devices (such as crutches, immobilizer, metal braces, orthotics, rigid stabilizer, or splints, ~~etc~~etc. and not to include neoprene sleeves), medications, injections (bursal, and/or joint, not including trigger point), and diathermy, can be utilized. Active modalities may consist of physical therapy, a physician-supervised home exercise program\*\*, and/or chiropractic care.



**\*\*Home Exercise Program (HEP)** – the following two elements are required to meet guidelines for completion of conservative therapy:

- Information provided on exercise prescription/plan AND
- Follow up with member with information provided regarding completion of HEP (after suitable 4-week period), or inability to complete HEP due to physical reason- i.e., increased pain, inability to physically perform exercises. (Patient inconvenience or noncompliance without explanation does not constitute “inability to complete” HEP).

**Joint Implants and Hardware** Dual-energy CT may be useful for metal artifact reduction if [available, but available but](#) is also imperfect as the correction is based on a projected approximation of x-ray absorption, and it does not correct for scatter (Boas, 2012). Dual-energy CT can be used to characterize crystal deposition disease, such as gout versus CPPD ([calcium pyrophosphate deposition](#)) (Chou, 2017).

**CT and Ankle Fractures** – One of the most frequently injured areas of the skeleton is the ankle. These injuries may include ligament sprains, as well as fractures. A suspected fracture is first imaged with conventional radiographs in anteroposterior, internal oblique and lateral projections. CT is used in patients with complex ankle and foot fractures after radiography.

**CT and Hip Trauma** – Computed tomography is primarily used to evaluate acute trauma, e.g., acetabular fracture or hip dislocation. It can detect intraarticular fragments and associated articular surface fractures, and it is useful in surgical planning.

**CT and Knee Fractures** – CT is used after plain films to evaluate fractures to the tibial plateau. These fractures occur just below the knee joint, involving the cartilage surface of the knee. Soft tissue injuries are usually associated with the fractures. The meniscus is a stabilizer of the knee, and it is very important to detect meniscal injury in patients with tibial plateau fractures. CT of the knee with two-dimensional reconstruction in the sagittal and coronal planes may be performed for evaluation of injuries with multiple fragments and comminuted fractures. Spiral CT has an advantage of rapid acquisition and reconstruction times and may improve the quality of images of bone. Soft tissue injuries are better demonstrated with MRI.

**CT and Knee Infections** – CT is used to depict early infection which may be evidenced by increased intraosseous density or the appearance of fragments of necrotic bone separated from living bone by soft tissue or fluid density. Contrast-enhanced CT may help in the visualization of abscesses and necrotic tissue.

**CT and Knee Tumors** – CT complements arthrography in diagnosing necrotic malignant soft-tissue tumors and other cysts and masses in the knee. Meniscal and ganglion cysts are palpable masses around the knee. CT is useful in evaluations of the vascular nature of lesions.

**CT and Legg-Calve-Perthes Disease (LPD)** – This childhood condition is associated with an insufficient blood supply to the femoral head which is then at risk for osteonecrosis. Clinical

signs of LPD include a limp with groin, thigh, or knee pain. Flexion and adduction contractures may develop as the disease progresses and eventually movement may only occur in the flexion-extension plane. This condition is staged based on plain radiographic findings. CT scans are used in the evaluation of LPD and can demonstrate changes in the bone trabecular pattern. They also allow diagnosis of bone collapse and sclerosis early in the disease where plain radiography is not as sensitive.

**CT and Osteolysis** – Since computed tomography scans show both the extent and the location of lytic lesions, they are useful to guide treatment decisions, as well as to assist in planning for surgical intervention when needed, in patients with suspected osteolysis after Total Hip Arthroplasty (THA).

**CT and Tarsal Coalition** – This is a congenital condition in which two or more bones in the mid-foot or hind-foot are joined. It usually presents during late childhood or late adolescence and is associated with repetitive ankle sprains. Mild pain, deep in the subtalar joint and limited range of motion are clinical symptoms. Tarsal coalition is detectable on oblique radiographs, but these are not routinely obtained at many institutions. Clinical diagnosis is not simple; it requires the expertise of skilled examiners. CT is valuable in diagnosing tarsal coalition because it allows differentiation of osseous from non-osseous coalitions and ~~also~~ depicts the extent of joint involvement as well as degenerative changes. It may also detect the overgrowth of the medial aspect of the talus that may be associated with talocalcaneal coalitions.

**American Academy of Pediatrics “Choosing Wisely” Guidelines** advise against ordering advanced imaging studies (MRI or CT) for most musculoskeletal conditions in a child until all appropriate clinical, laboratory and plain radiographic examinations have been completed. “History, physical examination, and appropriate radiographs remain the primary diagnostic modalities in pediatric orthopedics, as they are both diagnostic and prognostic for the great majority of pediatric musculoskeletal conditions. Examples of such conditions would include, but not be limited to, the work up of injury or pain (spine, knees and ankles), possible infection, and deformity. MRI examinations and other advanced imaging studies frequently require sedation in the young child (5 years old or less), and may not result in appropriate interpretation if clinical correlations cannot be made. Many conditions require specific MRI sequences or protocols best ordered by the specialist who will be treating the patient...if you believe findings warrant additional advanced imaging, discuss with the consulting orthopedic surgeon to make sure the optimal studies are ordered [\(AAP, 2018\).](#)”

## POLICY HISTORY

Date	Summary
May 2021	<ul style="list-style-type: none"> <li>• <u>Added unstable syndesmotic injury</u></li> <li>• <u>Removed ankle instability</u></li> <li>— <u>Added the following: navicular bone to high risk stress fracture, e</u></li> <li>— <u>Added information about suspected bone infection in the setting of ulcers and neuropathy and</u></li> </ul>

	<ul style="list-style-type: none"> <li>• <u>Additional information about following treatment for rheumatoid arthritis</u></li> <li>• <u>Clarified that pre-operative imaging is for diagnostic purposes only: “Pre-operative evaluation for <b>a planned surgery or procedure</b> if the imaging provides diagnostic information that is not available on prior studies (provider should be referred to the health plan for nondiagnostic surgical planning studies)” AND</u></li> <li>• <u>Removed *CT or MRI requests are not approvable for the following total knee arthroplasty (TKA) procedures:</u> <ul style="list-style-type: none"> <li>○ <u>Procedures utilizing computer-navigated or patient-specific or gender-specific instrumentation (Johnson, 2011)</u></li> <li>○ <u>Bicompartmental arthroplasty (investigational at this time) (Dudhniwala, 2016)</u></li> <li>○ <u>Note: Robot-assisted TKA (Makoplasty) (Banerjee, 2015; Nair, 2014)</u></li> </ul> </li> </ul> <p><u>These surgical procedures are not considered a covered service and are not reimbursable based on lack of current scientific evidence for clinically important improvement, safety or efficacy; or based on scientific evidence of increased risk of serious complications.</u></p> <p><u>Included early complications of hip surgery to the post operative evaluation list</u></p> <hr/> <ul style="list-style-type: none"> <li>•</li> </ul>
May 2020	<ul style="list-style-type: none"> <li>• Expanded orthopedic signs listing and moved to the top</li> <li>• Added note: With a positive orthopedic sign, an initial x-ray is always preferred. However, it is not required to approve advanced imaging.</li> <li>• Added labral tear/posterior impingement to approvable list</li> <li>• Added flatfoot deformity</li> <li>• Expanded section about initial work-up of a mass</li> <li>• Added the National Comprehensive Care Network as a reference for imaging guidance</li> <li>• Expanded the section on osteomyelitis</li> <li>• Added section on crystalline arthropathy</li> <li>• Revised the section on non or delayed union</li> <li>• Added a section on loose bodies and synovial chondromatosis</li> <li>• Added a pediatric section</li> <li>• Removed Makoplasty from not approvable list</li> </ul>

	<ul style="list-style-type: none"> <li>• Added a section about joint implants and hardware to the background section</li> <li>• Updated references</li> </ul>
May 2019	<ul style="list-style-type: none"> <li>• Reformatting in parallel with the new LE MRI. Updated references</li> <li>• Added indication: peripheral nerve entrapment</li> <li>• Criteria for approval of existing indications specified within the parameters of the current evidence base</li> <li>• Added initial statement about approvals: ‘Some indications are for MRI, CT, or MR or CT Arthrogram. More than one should not be approved at the same time’.</li> <li>• Added Extremity mass indications including peripheral lymphadenopathy; and mass with increased risk for malignancy</li> <li>• Modified Known Cancer indication to be more broad – ‘cancer staging, cancer restaging, signs or symptoms of recurrence’</li> <li>• Expanded section for infection of bone or joint to include list of signs or symptoms and laboratory findings (elevated ESR or CRP, elevated white blood cell count, positive joint aspiration)</li> </ul>

## REFERENCES

[Abousayed M, Alley M, Shakked R, et al. Adult-Acquired flatfoot deformity: Etiology, diagnosis and management. \*JBJS Reviews\*. 2017 Aug; 5\(8\):e7.](#)

American Academy of Pediatrics (AAP). Section on Orthopaedics and the Pediatric Orthopaedic Society of North America. Choosing Wisely®. <http://www.choosingwisely.org/clinician-lists/aap-posna-mri-or-ct-for-musculoskeletal-conditions-in-children/>. Released February 12, 2018.

[Banerjee S, Cherian JJ, Elmallah RK, et al. Robotic-assisted knee arthroplasty. \*Expert Rev Med Devices\*. 2015; 12\(6\):727-735.](#)

Beaman FD, Von Herrmann PF, Kransdorf MJ, et al. American College of Radiology ACR Appropriateness Criteria® - Suspected osteomyelitis, septic arthritis, or soft tissue infection (excluding spine and diabetic foot). *J Am Coll Radiol*. 2017; 14:S326-S337. <https://acsearch.acr.org/docs/3094201/Narrative/>.

Bencardino JT, Stone TJ, et al. ACR Appropriateness Criteria Stress (Fatigue/Insufficiency) Fracture, Including Sacrum, Excluding Other Vertebrae. *J Am Coll Radiol*. 2017 May;14(5S):S293-S306.

Bennett DL, Daffner RH, Weissman BN, et al. American College of Radiology ACR Appropriateness Criteria® - Non-traumatic Knee Pain. <https://acsearch.acr.org/docs/69432/Narrative/>. Published 2012.

[Bestic JM, Wessell DE, Beaman FD, et al. American College of Radiology ACR Appropriateness Criteria® - Primary Bone Tumors. https://acsearch.acr.org/docs/69421/Narrative/. Revised 2019.](https://acsearch.acr.org/docs/69421/Narrative/)

Boas FE, Fleischman D. CT artifacts: Causes and reduction techniques. *Imaging Med.* 2012; 4(2): 229-240.

Bouchard M, Mosca VS. Flatfoot deformity in children and adolescents: surgical indications and management. *J Am Acad Orthop Surg.* 2014; 22(10):623.

**Bowers S, Franco E. Chronic wounds: Evaluation and management. *Am Fam Physician.* 2020 Feb; 101(3):159-66.**

[Cecava ND, Dieckman S, Banks KP, et al. Traumatic knee injury: correlation of radiographic effusion size with the presence of internal derangement on magnetic resonance imaging. \*Emerg Radiol.\* 2018 Oct; 25\(5\):479-87.](#)

~~[Bowers S, Franco E. Chronic Wounds: Evaluation and Management. \*Am Fam Physician.\* 2020](#)~~

Chou H, Chin TY, et al. Dual-energy CT in gout – A review of current concepts and applications. *J Med Radiat Sci.* 2017 Mar; 64(1): 41–51.

Colebatch AN, Edwards CJ, Østergaard M, et al. EULAR recommendations for the use of imaging of the joints in the clinical management of rheumatoid arthritis. *Ann Rheum Dis.* 2013; 72:804-814. <http://ard.bmj.com/content/72/6/804>.

Dodwell ER. Osteomyelitis and septic arthritis in children: Current concepts. *Curr Opin Pediatrics.* 2013 Feb; 5(1):58–63.

Domkundwar S, Autkar G, Khadilkar SV, et al. Ultrasound and EMG–NCV study (electromyography and nerve conduction velocity) correlation in diagnosis of nerve pathologies. *J Ultrasound.* 2017 Jun; 20(2):111-122.

Dommett RM, Redaniel MT, Stevens MC, et al. Features of cancer in teenagers and young adults in primary care: a population-based nested case-control study. *Br J Cancer.* 2013; 108(11):2329. Epub 2013 Apr 25.

Dong Q, Jacobsen JA, Jamadar DA, et al. Entrapment neuropathies in the upper and lower limbs: Anatomy and MRI features. *Radiol Res Practice.* 2012; 2012:230679.

Donovan A, Rosenberg ZS, Cavalcanti CF. MR imaging of entrapment neuropathies of the lower extremity. *Radiographics.* 2010; 30(4).

Doral MN, Bilge O, Huri G, et al. Modern treatment of meniscal tears. *EFORT Open Rev.* 2018; 3(5):260-268.

~~Dudhniwala AG, Rath NK, Joshy S, et al. Early failure with the Journey Deuce bicompartamental knee arthroplasty. *Eur J Orthop Surg Traumatol*. 2016; 5:1-5.~~

Fayad LM, Carrino JA, Fishman EK. Musculoskeletal infection: Role of CT in the emergency department. *Radiographics*. 2007; 27:1723-36.

Felten R, Pemin P, Caillard S, et al. Avascular osteonecrosis in kidney transplant recipients: Risk factors in a recent cohort study and evaluation of the role of secondary hyperparathyroidism. *PLOS ONE*. February 22, 2019.

Fitzgerald JJ, Roberts CC, Daffner RH, et al. American College of Radiology ACR Appropriateness Criteria® - Follow-up of Malignant or Aggressive Musculoskeletal Tumors. <https://acsearch.acr.org/docs/69428/Narrative/>. Published 2015.

Fox MG, Chang, EY, Amini B, et al. American College of Radiology ACR Appropriateness Criteria® - Chronic Knee Pain. <https://acsearch.acr.org/docs/69432/Narrative/>. Revised 2018.

Fox MG, Graham JA, Skelton BW, et al. Prospective evaluation of agreement and accuracy in the diagnosis of meniscal tears: MR Arthrography a short time after injection versus CT arthrography after a moderate delay. *Am J Roentgenol*. 2016; 207:142-149. 10.2214/AJR.15.14517.

Fritz J, Lurie B, Miller TT, et al. MR imaging of hip arthroplasty implants. *Radiographics*. 2014; 34(4):E106-E132. <http://pubs.rsna.org/doi/abs/10.1148/rg.344140010>.

Fritz J, Lurie B, Potter HG. MR imaging of knee arthroplasty implants. *Radiographics*. 2015 Aug; 35(5).

Fukushima W, Fujioka M, Kubo T, et al. Nationwide epidemiologic survey of idiopathic osteonecrosis of the femoral head. *Clin Orthop Relat Res*. 2010 Oct; 468(10):2715-24

Gaddey HL, Riegel AM. Unexplained lymphadenopathy: Evaluation and differential diagnosis. *Am Fam Physician*. 2016 Dec 1; 94(11):896-903.

Garras DN, Raikin SM, Bhat SB, et al. MRI is Unnecessary for Diagnosing Acute Achilles Tendon Ruptures: Clinical Diagnostic Criteria. *Clin Orthop Relat Res*. 2012 Aug; 470(8): 2268–2273.

Gill SK, Smith J, Fox R, Chessier TJ. Investigation of occult hip fractures: the use of CT and MRI. *Sci World J*. 2013; 2013:830319.

Glaser C. Tarsal Coalitions: A Practical Approach to a Not-So-Rare Entity. *J Belgian Soc Radiol*. 2016; 100(1):104. DOI: <http://doi.org/10.5334/jbr-btr.1224>.

Glaudemans AWJM, Jutte PC, et al. Consensus document for the diagnosis of peripheral bone infection in adults: a joint paper by the EANM, EBJIS, and ESR (with ESCMID endorsement). *Eur J Nucl Med Mol Imaging*. 2019; 46(4): 957–970.

Groh MM, Herrera J. A comprehensive review of hip labral tears. *Curr Rev Musculoskelet Med*. 2009; 2(2):105-17.

Guggenberger R, Pfirrmann CW, Koch PP, et al. Assessment of lower limb length and alignment by biplanar linear radiography: Comparison with supine CT and upright full-length radiography. *Am J Roentgenol*. 2014; 202(2):W161-W167.

Hananouchi T, Yasui Y, Yamamoto K, et al. Anterior impingement test for labral lesions has high positive predictive value. *Clin Orthop Relat Res*. 2012 Dec; 470(12):3524–3529.

Hesper T, Zilkens C, Bittersohl B, et al. Imaging modalities in patients with slipped capital femoral epiphysis. *J Child Orthop*. 2017; 11(2):99-106. <http://doi.org/10.1302/1863-2548-11-160276>. <https://www.ncbi.nlm.nih.gov/pubmed/28529656>.

Holzapfel K, Regler J, Baum T, et. al. Local Staging of Soft-Tissue Sarcoma: Emphasis on Assessment of Neurovascular Encasement—Value of MR Imaging in 174 Confirmed Cases. *Radiology*. 2015 Jan; 275(2).

Hussin P, Mawardi M, et al. The 'Chalky Culprit' of acute locked knee. *G Chir*. 2014; 35(9-10):239–240.

Iyer RS, Chapman T, et al. Pediatric bone imaging: Diagnostic imaging of osteoid osteoma. *Am J Roentgenol*. 2012; 198:1039-1052.

~~Johnson AJ, Costa CR, Mont MA. Do we need gender-specific total joint arthroplasty? *Clin Orthop Relat Res*. 2011; 469(7):1852-1858.~~

Kamegaya M, Saisu T, Nakamura J, et al. Drehmann sign and femoro-acetabular impingement in SCFE. *J Ped Orthop*. 2011 Dec; 31(8):853-857.

Katz JN, Brophy RH, Chaisson CE, et al. Surgery versus physical therapy for a meniscal tear and osteoarthritis. *N Engl J Med*. 2013; 368:1675-1684.

Kellar J, Givertz A, et al. Biphosphonate-related Femoral Shaft Fracture. *Clin Pract Cases Emerg Med*. 2020 Feb; 4(1):62-64.

Khan AN, Seriki DM, Hutchinson E, et al. Legg-Calve-Perthes Disease. *Emedicine*. 2011. <http://emedicine.medscape.com/article/410482-overview>.

~~Kamegaya M, Saisu T, Nakamura J, et al. Drehmann sign and femoro-acetabular impingement in SCFE. *J Ped Orthop*. 2011 Dec; 31(8):853-857.~~



~~Katz JN, Brophy RH, Chaisson CE, et al. Surgery versus physical therapy for a meniscal tear and osteoarthritis. *N Engl J Med*. 2013; 368:1675-1684.~~

Kircher MF, Willman JK. Molecular body imaging: MR imaging, CT, and US. Part II. Applications. *Radiology*. 2012; 264(2):349.

Kopf S, Beaufils P, Hirschmann MT, et al. Management of traumatic meniscus tears: the 2019 ESSKA meniscus consensus. *Knee Surg Sports Traumatol Arthrosc*. 2020; 28(4):1177-1194.

Laya BF, Restrepo R, Lee EY. Practical imaging evaluation of foreign bodies in children: An update. *Radiol Clin North Am*. 2017 Jul; 55(4):845-867.

Lefevre N, Naouri JF, Herman S, et al. A current review of the meniscus imaging: Proposition of a useful tool for its radiologic analysis. *Radiol Res Pract*. 2016; 2016:8329296.

Lipsky BA, Senneville E, Abbas ZG, et al. Guidelines on the diagnosis and treatment of foot infection in persons with diabetes (IWGDF 2019 update). *Diab Metab Res Rev*. 2020 Mar; 36 Suppl 1:-e3280.

Mandell JC, Khurana B, Smith JT, et al. Osteomyelitis of the lower extremity: Pathophysiology, imaging, and classification, with an emphasis on diabetic foot infection. *Emerg Radiol*. 2017.

Mintz DN, Roberts CC, et al. ACR Appropriateness Criteria Chronic Hip Pain. *J Am Coll Radiol*. 2017 May; 14(5S):S90-S102.

Mohankumar R, White L, Naraghi A. Review. Pitfalls and pearls in MRI of the knee. *AJR Am J Roentgenol*. 2014; 203:516-530. <https://www.ajronline.org/doi/full/10.2214/AJR.14.12969>.

Mohseni S, Shojaiepard A, Khorgami Z, et al. Peripheral lymphadenopathy: Approach and diagnostic tools. *Iran J Med Sci*. 2014 Mar; 39(2 Suppl):158-170.

Mordecai SC, Al-Hadithy N, Ware HE, et al. Treatment of meniscal tears: An evidence based approach. *World J Orthop*. 2014 Jul 18; 5(3):233-241.

~~Morrison WB, Weissman BN, Kransdorf MJ, et al. American College of Radiology ACR Appropriateness Criteria® Primary Bone Tumors. <https://acsearch.acr.org/docs/69421/Narrative/>. Published 2013.~~

Morshed S. Current Options for Determining Fracture Union. *Adv Med*. 2014; 2014:708574.

Mullan CP, Madan R, Trotman-Dickenson B, et al. Radiology of chest wall masses. *AJR*. 2011; 2011:197(3).

Murphey MD, Foreman KL, Klassen Fischer MK, et al. From the radiologic pathology archives imaging of osteonecrosis: Radiologic-pathologic correlation. *RadioGraphics*. 2014 Jul 14; 34(4)-



Murphey MD, Roberts CC, Bencardino JT, et al. American College of Radiology ACR Appropriateness Criteria® - Osteonecrosis of the Hip. *J Am Coll Radiol*. 2016; 13:147-155. <https://acsearch.acr.org/docs/69420/Narrative/>.

Murphey MD, Wessell DE, et al. ACR Appropriateness Criteria Soft-Tissue Masses. *J Am Coll Radiol*. 2018 May; 15(Suppl 5):S189-S197.

Naraghi A, White LM. MRI of labral and chondral lesions of the hip. *AJR Am J Roentgenol*. 2015; 205(3):479-490. <https://www.ajronline.org/doi/full/10.2214/AJR.14.12581>.

National Comprehensive Cancer Network (NCCN). Imaging guidelines. 2019. [https://www.nccn.org/professionals/physician\\_gls/default.aspx](https://www.nccn.org/professionals/physician_gls/default.aspx).

Patel DS, Roth M, Kapil N. Stress fractures: diagnosis, treatment, and prevention. *Am Fam Physician*. 2011; 83:39-46. <https://www.ncbi.nlm.nih.gov/pubmed/21888126>.

Peck DM, Voss LM, Voss TT. Slipped capital femoral epiphysis: Diagnosis and management. *Am Fam Physician*. 2017 Jun 15; 95(12):779-784.

Peck J, Gustafson KE, Bahner DP. Diagnosis of achilles tendon rupture with ultrasound in the emergency department setting. *Int J Academ Med*. 2017; 3(3):205-207.

**Pitocco D, Spanu T, Di Leo M, et al. Diabetic foot infections: A comprehensive overview. *Eur Rev Med Pharmacol Sci*. 2019 Apr; 23(2 Suppl):26-37.**

**Prat-Fabregat S, Camacho-Carrasco P. Treatment strategy for tibial plateau fractures: An update. *EFORT Open Rev*. 2016;1(5):225-232. doi:10.1302/2058-5241.1.000031**

Rabinovich RV, Haleem AM, Rozbruch SR. Complex ankle arthrodesis: Review of the literature. *World J Orthop*. 2015 Sep 18; 6(8):602–613.

**Rajani R, Quinn R. Synovial chondromatosis. *OrthoInfo*. 2016 Dec.**  
<https://orthoinfo.aaos.org/en/diseases--conditions/synovial-chondromatosis>

**Ross AB, Lee KS, Chang, EY, et al. American College of Radiology ACR Appropriateness Criteria® - Acute Hip Pain–Suspected Fracture. Revised 2018.**  
<https://acsearch.acr.org/docs/3082587/Narrative/>.

Ross AB, Lee KS, **Chang, EY** et al. ACR Appropriateness Criteria Acute Hip Pain-Suspected Fracture. *J Am Coll Radiol*. 2019 May; 16(5S):S18-S25.

Sabharwal S, Kumar A. Methods for Assessing Leg Length Discrepancy. *Clin Orthop Relat Res*. 2008 Dec; 466(12):2910–2922.

Sadineni RT, Psumarthy A, Bellapa NC, et al. Imaging patterns in MRI in recent bone injuries following negative or inconclusive plain radiographs. *J Clin Diagn Res*. 2015 Oct; 9(10):TC10–TC13.

Salih S, Blakey C, Chan D, et al. The callus fracture sign: a radiological predictor of progression to hypertrophic non-union in diaphyseal tibial fractures. *Strat Trauma Limb Reconstr*. 2015; 10(3):149–153.

Scalcione LR, Gimber LH, Ho AM, et al. Spectrum of carpal dislocations and fracture-dislocations: Imaging and management. *Am J Roentgenol*. 2014; 203:541-550.

Sinha S, Peach AH. Diagnosis and management of soft tissue sarcoma. *BMJ*. 2010; 341:c7170.

Slaughter AJ, Reynolds KA, Jambhekar K, et al. Clinical orthopedic examination findings in the lower extremity: Correlation with imaging studies and diagnostic efficacy. *Radiographics*. 2014 Mar; 34(2).

Smith TO, Drew BT, Toms AP, et al. Accuracy of magnetic resonance imaging, magnetic resonance arthrography and computed tomography for the detection of chondral lesions of the knee. *Knee Surg Sports Traumatol Arthrosc*. 2012 Dec; 20(12):2367-79. Epub 2012 Jan 24.

Smith BE, Thacker D, Crewesmith A, Hall M. Special tests for assessing meniscal tears within the knee: A systematic review and meta-analysis. *Evid Based Med*. 2015; 20:88-97.

Subhawong TK, Fishman EK, Swart JE, et al. Soft-tissue masses and masslike conditions: What does CT add to diagnosis and management? *AJR Am J Roentgenol*. 2010 Jun; 194(6):1559–1567.

[Taljanovic, MS, Chang, EY, Ha, AS, et al. American College of Radiology Appropriateness Criteria® - Acute Trauma to the Knee. Revised 2019. https://acsearch.acr.org/docs/69419/Narrative/.](https://acsearch.acr.org/docs/69419/Narrative/)

Thorpe SW, Wukich DK. Tarsal coalitions in the adult population: Does treatment differ from the adolescent? *Foot Ankle Clin*. 2012 Jun; 17(2):195-204.

Tos P, Crosio A, Pugliese P, et. el. Painful scar neuropathy: Principles of diagnosis and treatment. *Plast Aesthet Res*. 2015; 2:156-64.

~~Tuite MJ, Daffner RH, Weissman BN, et al. American College of Radiology Appropriateness Criteria® - Acute Trauma to the Knee. 2014. https://acsearch.acr.org/docs/69419/Narrative/.~~

Turan A, Celtikci P, Tufan A, et al. Basic radiological assessment of synovial diseases: A pictorial essay. *Eur J Rheumatol*. 2017 Jun; 4(2):166-74.

Van Bergen CJA, Van den Ende KIM, Ten Brinke B, et al. Osteochondritis dissecans of the capitellum in adolescents. *World J Orthop*. 2016 Feb 18; 7(2):102–108.

Van Dijk CN, Reilingh ML, Zengerink M, et al. Osteochondral defects in the ankle: Why painful? *Knee Surg Sports Traumatol Arthrosc*. 2010 May; 18(5):570–580.

Vuurberg G, Hoorntje A, Wink LM, et.al. Diagnosis, treatment, and prevention of ankle sprains: Update of an evidence-based clinical guideline. *Br J Sports Med*. 2018; 52(15):956. Epub 2018 Mar 7.

Wali Y, Almaskan S. Avascular necrosis of the hip in sickle cell disease in oman. Is it serious enough to warrant bone marrow transplantation? *Sultan Qaboos Univ Med J*. 2011 Feb; 11(1):127–128.

Walker EA, Beaman FD, et al. ACR Appropriateness Criteria Suspected Osteomyelitis of the Foot in Patients with Diabetes Mellitus. *J Am Coll Radiol*. 2019 Nov; 16(Suppl 11):S440-S450.


[Wheeless III CR. Wheelless' Textbook of Orthopaedics. Duke University Medical Center's Division of Orthopedic Surgery. © 1996-2018 Data Trace Internet Publishing, LLC. Updated 2018.](#)

~~Ward RJ, Weissman BN, Kransdorf MJ, et al. American College of Radiology ACR Appropriateness Criteria® Acute Hip Pain Suspected Fracture. 2013.~~  
~~<https://acsearch.acr.org/docs/3082587/Narrative/>.~~

Wilkins R, Bisson LJ. Operative versus nonoperative management of acute achilles tendon ruptures: A quantitative systematic review of randomized controlled trials. *Am J Sports Med*. 2012; 40(9):2154. Epub 2012 Jul 16.

Zoga AC, Weissman BN, Kransdorf MJ, et al. American College of Radiology ACR Appropriateness Criteria® – Soft Tissue Masses. 2017.  
<https://acsearch.acr.org/docs/69434/Narrative/>.

Zollars ES, Hyer M, Wolf B, et al. Measuring lupus arthritis activity using contrasted high-field MRI. Associations with clinical measures of disease activity and novel patterns of disease Lupus. *Science Med*. 2018; 5:e000264. doi: 10.1136/lupus-2018-000264.

Reviewed / Approved by  M. Atif Khalid, M.D., Medical Director, Radiology

### GENERAL INFORMATION

It is an expectation that all patients receive care/services from a licensed clinician. All appropriate supporting documentation, including recent pertinent office visit notes, laboratory data, and results of any special testing must be provided. If applicable: All prior relevant imaging results and the reason that alternative imaging cannot be performed must be included in the documentation submitted.

**Disclaimer:** Magellan Healthcare service authorization policies do not constitute medical advice and are not intended to govern or otherwise influence the practice of medicine. These policies are not meant to supplant your normal procedures, evaluation, diagnosis, treatment and/or care plans for your patients. Your professional judgement must be exercised and followed in all respects with regard to the treatment and care of your patients. These policies apply to all Magellan Healthcare subsidiaries including, but not limited to, National Imaging Associates (“Magellan”). The policies constitute only the reimbursement and coverage guidelines of Magellan. Coverage for services varies for individual members in accordance with the terms and conditions of applicable Certificates of Coverage, Summary Plan Descriptions, or contracts with governing regulatory agencies. Magellan reserves the right to review and update the guidelines at its sole discretion. Notice of such changes, if necessary, shall be provided in accordance with the terms and conditions of provider agreements and any applicable laws or regulations.