

AmeriHealth Caritas Louisiana

National Imaging Associates, Inc.*	
Clinical guidelines LOWER EXTREMITY CTA/CTV	Original Date: July 01, 2008
CPT Codes: 73706	Last Revised Date: May 2021
Guideline Number: NIA_CG_061-1	Implementation Date: January 2022

INDICATIONS FOR LOWER EXTREMITY CTA/CTV (COMPUTED TOMOGRAPHY ANGIOGRAM / COMPUTED TOMOGRAPHY VENOGRAM)

Abdominal Arteries CTA (CT Angiography) (CPT Code 75635) includes run-off so this is never approved when that one has been.

Peripheral Vascular Disease and Abdominal Arteries CTA (CT Angiography) (CPT Code 75635) has not been recently approved

- Critical Limb ischemia **ANY** of the below with clinical signs of peripheral artery disease. Ultrasound imaging is not needed. If done and negative, it should still be approved due to high false negative rate (Shishehbor, 2016; Weiss, 2017~~8~~)
 - Ischemic rest pain
 - Tissue loss
 - Gangrene
- Claudication with abnormal (ankle/brachial index, arterial Doppler) (Ahmed, 2017; Pollak, 2012, 2013)
- Clinical concern for vascular cause of ulcers with abnormal or indeterminate ultrasound (ankle/brachial index, arterial Doppler) (Rosyd, 2017)
- After stenting or surgery with signs of recurrent symptoms OR abnormal ankle/brachial index; abnormal or indeterminate arterial Doppler, OR pulse volume recording) (Pollak, 2012)

Popliteal Artery Entrapment Syndrome with abnormal arterial ultrasound (Williams, 2015)

Deep Venous Thrombosis with clinical suspicion of lower extremity DVT after abnormal or non-diagnostic ultrasound where a positive study would change management (Hanley, 2013; Karande, 2016; Katz, 2014)

* National Imaging Associates, Inc. (NIA) is a subsidiary of Magellan Healthcare, Inc.

Clinical suspicion of vascular disease with abnormal or indeterminate ultrasound or other imaging

- Tumor invasion (Kransdorf, 2018)
- Trauma (Wani, 2012)
- Vasculitis (Fonseca, 2017)
- Aneurysm (Verikokos, 2014)
- Stenosis/occlusions (Menke, 2010)

Hemodialysis Graft Dysfunction after Doppler ultrasound not adequate for treatment decisions (Murphy, 2017)

Vascular Malformation (Madani, 2015; Obara, 2019) - If MRA is contraindicated

- Non diagnostic doppler ultrasound

Note: CTA useful in delineating high flow lesions such as an arteriovenous malformation.

Evaluation of traumatic injuries to the LE with clinical findings suggestive of arterial injury (Wani, 2012)-

For a assessment/evaluation of known vascular disease/condition.

Pre-operative/procedural evaluation

- Pre-operative evaluation for a planned surgery or procedure (Ahmed, 2017)

Post-operative/procedural evaluation

- A follow-up study may be needed to help evaluate a patient's progress after treatment, procedure, intervention, or surgery. Documentation requires a medical reason that clearly indicates why additional imaging is needed for the type and area(s) requested (Conte, 2019; Cooper 2018)

Special Circumstances

(Weiss, 2017)

- High suspicion of an acute arterial obstruction - Arteriography preferred (the gold standard).
- Renal impairment
 - Not on dialysis
 - Mild to moderate, GFR 30-89 ml/min MRA can be done
 - Severe, GFR < 30 ml/min MRA without contrast
 - On dialysis
 - CTA with contrast can be done
- Doppler ultrasound can be useful in evaluating bypass grafts

BACKGROUND

Lower extremity computed tomography angiography (CTA) is an effective, noninvasive and robust imaging modality that is used in the assessment of symptomatic lower extremity vascular disease. It has excellent spatial resolution and shows accurate details of peripheral vasculature. CTA is an effective alternative to catheter-based angiography and allows accurate planning of open surgical and endovascular interventions.

OVERVIEW

Abdominal Arteries CTA – A—For imaging of the abdomen, pelvis **AND** both legs (CTA aorto-iliofemoral runoff; abdominal aorta and bilateral iliofemoral lower extremity runoff) use CPT code 75635.

Peripheral Arterial Disease — CTA is used in the evaluation of patients with peripheral arterial disease. It can be used to evaluate the patency after revascularization procedures. It is the modality of choice in patients with intermittent claudication. A drawback is its hampered vessel assessment caused by the depiction of arterial wall calcifications, resulting in a decreased accuracy in severely calcified arteries.

Chronic Limb Threatening Ischemia — Assessment and promotion of blood flow through the calf arteries is very important in patients with chronic limb threatening ischemia. CT Angiography allows for visualization of pedal vessels.

Surgical or Percutaneous Revascularization – CTA is accurate in the detection of graft-related complications, including stenosis and aneurysmal changes. It can reveal both vascular and extravascular complications.

CTA and screening for peripheral vascular disease: The USPSTF (U.S. Preventative Services Task Force) does not recommend routine screening for peripheral vascular disease in asymptomatic patients. High risk patients (e.g., diabetics) may be screened with ABI (ankle brachial index) and duplex ultrasound.

POLICY HISTORY

Date	Summary
May 2021	Reviewed literature for updates No changes
<u>May 2020</u>	<ul style="list-style-type: none">• <u>Clarified that CTA does not include a baseline CT exam</u>• <u>Expanded section about vascular malformation to include initial testing.</u>• <u>Added information about renal function and contrast agents</u>• <u>Added acute arterial obstruction and renal impairment</u>• <u>Simplified language</u>• <u>Updated references</u>

May 2019

- Added indication for deep venous thrombosis
- Reformatting and new references.

May 2019

- ~~Added indication for deep venous thrombosis~~
- ~~Reformatting and new references.~~

May 2020

- ~~Clarified that CTA does not include a baseline CT exam~~
- ~~Expanded section about vascular malformation to include initial testing.~~
- ~~Added information about renal function and contrast agents~~
- ~~Added acute arterial obstruction and renal impairment~~
- ~~Simplified language~~
- ~~Updated references~~

REFERENCES

Aboyans V, Ricco JB, Bartelink MLEL, et al. 2017 ESC guidelines on the diagnosis and treatment of peripheral arterial diseases, in collaboration with the European Society for Vascular Surgery (ESVS): document covering atherosclerotic disease of extracranial carotid and vertebral, mesenteric, renal, upper and lower extremity arteries endorsed by: the European Stroke Organization (ESO) the Task Force for the Diagnosis and Treatment of Peripheral Arterial Diseases of the European Society of Cardiology (ESC) and of the European Society for Vascular Surgery (ESVS). *Eur Heart J*. 2018 Mar; 39:763-816.

Ahmed O, Hanley M, Bennett SJ, et al. American College of Radiology ACR Appropriateness Criteria[®] - Vascular Claudication: Assessment for Revascularization. *J Am Coll Radiol*. 2017; 14(5 Suppl): S372-S379. <https://acsearch.acr.org/docs/69411/Narrative/>.

Conte MS, Bradbury AW, Kolh P, et al. Global vascular guidelines on the management of chronic limb-threatening ischemia. *J Vasc Surg*. 2019 Jun; 69(6S):3S-1255.e40. Epub 2019 May 28.

Cooper K, Majdalany BS, Kalva SP, et al. ACR Appropriateness Criteria lower extremity arterial revascularization – post-therapy. *J Am Coll Radiol*. 2018 May; 15(5Suppl):S104-S115.

Farber A. Chronic limb-threatening ischemia. *N Engl J Med*. 2018 Jul 12; 379(2):171-180. doi: 10.1056/NEJMcp1709326.

Fonseka CL, Galappaththi SR, Abeyaratne D, et al. A Case of Polyarteritis Nodosa Presenting as Rapidly Progressing Intermittent Claudication of Right Leg. *Case Reports in Medicine*. 2017; 2017:4219718.

Gerhard-Herman MD, Gornik HL. AHA/ACC guideline on the management of patients with lower extremity peripheral artery disease: A report of the American College of Cardiology/American Heart Association task force on clinical practice guidelines. *J Am Coll Cardiol*. 2017 Mar; 69(11).

Hanley M, Donahue J, Rybicki FJ, et al. American College of Radiology ACR Appropriateness Criteria® Clinical Condition: Suspected Lower-Extremity Deep Vein Thrombosis. <https://acsearch.acr.org/docs/69416/Narrative/>. Published 2013.

Jin T, Wu G, Li X, et al. Evaluation of vascular invasion in patients with musculoskeletal tumors of lower extremities: Use of time-resolved 3D MR angiography at 3-T. *Acta Radiol*. 2018 May; 59(5):586-592.

Karande GY, Hedgire SS, Sanchez Y, et al. Advanced imaging in acute and chronic deep vein thrombosis. *Cardiovasc Diagn Ther*. 2016; 6(6):493-507. <http://doi.org/10.21037/cdt.2016.12.06>.

Katz DS, Fruauff K, Kranz A, et al. Imaging of deep venous thrombosis: A multimodality overview. *Appl Radiol*. March 5, 2014.

Kransdorf MJ, Murphey MD, Wessell DE, et al. American College of Radiology ACR Appropriateness Criteria® - Soft Tissue Masses. <https://acsearch.acr.org/docs/69434/Narrative/>. Published 2017.

Lopera JE, Trimmer CK, Josephs SG, et al. Multidetector CT angiography of infrainguinal arterial bypass. *RadioGraphics*. 2008; 28(2):529. doi: 10.1148/rg.282075032. <https://pubs.rsna.org/doi/abs/10.1148/rg.282075032>.

Madani H, Farrant J, Chhaya N, et al. Peripheral limb vascular malformations: An update of appropriate imaging and treatment options of a challenging condition. *Br J Radiol*. 2015; 88(1047):20140406.

Menke J, Larsen J. Meta-analysis: Accuracy of contrast-enhanced magnetic resonance angiography for assessing steno-occlusions in peripheral arterial disease. *Ann Intern Med*. 2010; 153(5):325-334. doi: 10.7326/0003-4819-153-5-201009070-00007.

Met R, Bipat S, Legemate DA, et al. Diagnostic performance of computed tomography angiography in peripheral arterial disease: A systematic review and meta-analysis. *JAMA*. 2009; 301(4):415-424. doi: 10.1001/jama.301.4.415.

Murphy EA, Ross RA, Jones RG, et al. Imaging in vascular access. *Cardiovasc Eng Technol*. 2017; 8(3):255-272.

Obara P, McCool J, Kalva SP, et al. ACR Appropriateness Criteria clinically suspected vascular malformation of the extremities. *J Am Coll Radiol*. 2019 Nov; 16(11S):S340-S347.

Pollak AW, Kramer CM. MRI in lower extremity peripheral arterial disease: Recent advancements. *Curr Cardiovasc Imaging Rep*. 2013 Feb 1; 6(1):55–60.

Pollak AW, Norton P, Kramer CM. Multimodality imaging of lower extremity peripheral arterial disease: Current role and future directions. *Circ Cardiovasc Imaging*. 2012 Nov 1; 5(6):797–807.

Rosyd FN. Etiology, pathophysiology, diagnosis and management of diabetics' foot ulcer. *Int J Res Med Sci*. 2017 Oct; 5(10):4206-4213.

Shishehbor MH, White CJ, Gray BH, et al. Critical limb ischemia. *Am Coll Cardiol*. 2016; 68(18):2002–15.

Tuite M, Kransdorf M, Beaman F, et al. American College of Radiology ACR Appropriateness Criteria® - Acute Trauma to the Knee. [http://www.jacr.org/article/S1546-1440\(15\)00831-5/fulltext](http://www.jacr.org/article/S1546-1440(15)00831-5/fulltext). Published 2018.

Verikokos C, Karaolani G, Doulaptsis M, et al. Giant popliteal artery aneurysm: case report and review of the literature. *Case Rep Vasc Med*. 2014; 2014:780561.

Wani ML, Ahangar AG, Ganie FA, et al. Vascular injuries:Trends in management. *Trauma Mon*. 2012; 17(2):266–269.

Weiss C, Azene ER, Azene EM, et al. American College of Radiology (ACR). ACR Appropriateness Criteria® - Sudden Onset of Cold, Painful Leg. *J Am Coll Radiol*. 2017 May; 14(5Suppl):S307-S313.

Williams C, Kennedy D, Bastian-Jordan M, et al. A new diagnostic approach to popliteal artery entrapment syndrome. *J Med Radiat Sci*. 2015; 62(3):226–229.

Reviewed / Approved by NIA Clinical Guideline Committee

GENERAL INFORMATION

It is an expectation that a
supporting documentatio
of any special testing mu
reason that alternative imaging cannot be performed must be included in the documentation
submitted.

M. Atif Khalid MD

; from a licensed clinician. All appropriate
office visit notes, laboratory data, and results
I prior relevant imaging results and the

Disclaimer: Magellan Healthcare service authorization policies do not constitute medical advice and are not intended to govern or otherwise influence the practice of medicine. These policies are not meant to supplant your normal procedures, evaluation, diagnosis, treatment and/or care plans for your patients. Your professional judgement must be exercised and followed in all respects with regard to the treatment and care of your patients. These policies apply to all Magellan Healthcare subsidiaries including, but not limited to, National Imaging Associates (“Magellan”). The policies constitute only the reimbursement and coverage guidelines of Magellan. Coverage for services varies for individual members in accordance with the terms and conditions of applicable Certificates of Coverage, Summary Plan Descriptions, or contracts with governing regulatory agencies. Magellan reserves the right to review and update the guidelines at its sole discretion. Notice of such changes, if necessary, shall be provided in accordance with the terms and conditions of provider agreements and any applicable laws or regulations.