

AmeriHealth Caritas Louisiana

National Imaging Associates, Inc.*	
Clinical guidelines TEMPOROMANDIBULAR JOINT (TMJ) MRI	Original Date: May 23, 2003
CPT Code: 70336	Last Revised Date: June 2021
Guideline Number: NIA_CG_007	Implementation Date: January 2022

INDICATIONS FOR TEMPOROMANDIBULAR JOINT (TMJ) MRI

For evaluation of temporomandibular joint dysfunction (TMD) with suspected internal joint derangement with both: with ALL of the following

(Bag, 2014; Gauer, 2015; Petscavage-Thomas, 2014)

- Persistent symptoms of facial or jaw pain, restricted range of motion, pain and/or noise with TMJ function (i.e., chewing) ~~(Bag, 2014)~~ AND
- Conservative therapy with a trial of anti-inflammatory AND behavioral modification* has been unsuccessful for at least four (4) weeks

~~Initial X-rays have been performed~~ Note: X-ray should be the initial study if there is recent trauma, dislocation, malocclusion, or dental infection

* Behavioral modification includes patient education, self-care, cognitive behavior therapy, physical therapy, and occlusal devices. Muscle relaxants can be used for spasm.

For evaluation of Juvenile idiopathic arthritis (JIA)

(Granquist, 2018, Petscavage-Thomas, 2014)

Abnormal initial x-ray or ultrasound needing additional imaging

(Bag, 2014)

Pre-operative evaluation in candidates for orthognathic surgery

Post-operative Evaluation

(Hoffman, 2015)

* National Imaging Associates, Inc. (NIA) is a subsidiary of Magellan Healthcare, Inc.

- A follow-up study may be needed to help evaluate a patient’s progress after treatment, procedure, intervention, or surgery. Documentation requires a medical reason that clearly indicates why additional imaging is needed for the type and area(s) requested.

BACKGROUND

Temporomandibular joint (TMJ) dysfunction causes pain and dysfunction in the jaw joint and muscles controlling jaw movement. Symptoms may include jaw pain, masticator muscle stiffness, limited movement or locking of the jaw, clicking or popping in jaw joint when opening or closing the mouth, and a change in how the upper and lower teeth fit together. The cause of the condition is not always clear but may include acute or chronic trauma to the jaw or temporomandibular joint, e.g., grinding of teeth, clenching of jaw, or impact in an accident. Osteoarthritis or rheumatoid arthritis may also contribute to the condition.

Etiologies of TMJ dysfunction ([TMD](#)) include intra-articular (intracapsular) and extra-articular (extracapsular pathology). Intra-articular (intracapsular pathology), [such as](#) disc displacement and coexisting osteoarthritis or degenerative joint disease, is considered the most common cause of serious TMJ pain and dysfunction and the most likely to be treated surgically. Extra-articular (extracapsular pathology) includes musculoskeletal (bone, masticatory muscles and tendons) and central nervous system/peripheral nervous system (ASTMJS, 2001).

Imaging can assist in the diagnosis of TMD when history and physical examination findings are equivocal. The initial study should be plain radiography (transcranial and transmaxillary views) or panoramic radiography [when there is recent trauma, dislocation, malocclusion, or dental infection](#)-(Gauer, 2015). Ultrasound is an inexpensive and easily performed imaging modality that can also be used to evaluate the TMJ (Tu, 2018). CT is useful to evaluate the bony structures of the TMJ when there is suspicion of bony involvement (i.e., fractures, erosions, infection, invasion by tumor, as well as congenital anomalies) (Bag, 2014). Magnetic resonance imaging (MRI) has the highest sensitivity, specificity, and accuracy in the evaluation of temporomandibular joint dysfunction and provides tissue contrast for visualizing the soft tissue and periarticular structures of the TMJ.

Conservative care for TMD includes patient education, self-care, behavioral modification, cognitive behavioral therapy/biofeedback, medication, physical therapy, and occlusive devices. Medications include NSAIDs and muscle relaxants and in chronic cases, benzodiazepines, or antidepressants. There is lack of high-quality evidence and uncertainty about the effectiveness of manual therapy and therapeutic physical therapy in treating TMJ dysfunction- (Armijo-Olivo, 2016). The use of occlusive splints is thought to alleviate some of the degenerative forces on the TMJ which may be helpful in patients with bruxism or nocturnal teeth clenching. The preferred devices are unclear from the literature and dental consultation is required (Gauer, 2015). In systematic reviews, there has been [short-short](#)-term benefit observed from splinting

but no clear role in the overall long-term treatment of TMD patients (Ebrahim, 2012; Pficer, 2017).

POLICY HISTORY

Date	Summary
June 2021	<p>Updated reference and background section</p> <p>Deleted: Initial x-rays have been performed</p> <p>Added:</p> <p>Note: X-ray should be the initial study if there is recent trauma, dislocation, malocclusion, or dental infection</p> <p>* Behavioral modification includes patient education, self-care, cognitive behavior therapy, physical therapy, and occlusal devices.</p> <p>Muscle relaxants can be used for spasm.</p>
<u>May 2020</u>	<p><u>Added:</u></p> <ul style="list-style-type: none"> • <u>For evaluation of temporomandibular joint dysfunction (TMD) with suspected internal joint derangement with ALL of the following</u> <ul style="list-style-type: none"> ○ <u>Persistent symptoms of facial or jaw pain, restricted range of motion, pain and/or noise with TMJ function (i.e., chewing)</u> ○ <u>Conservative therapy with a trial of anti-inflammatory AND behavioral modification has been unsuccessful for at least four (4) weeks</u> ○ <u>Initial X-rays have been performed</u> • <u>For evaluation of Juvenile idiopathic arthritis (JIA)</u> • <u>Abnormal initial x-ray or ultrasound needing additional imaging</u> <p>-</p> <p><u>Deleted:</u></p> <ul style="list-style-type: none"> • <u>Locked or Frozen Jaw</u> <ul style="list-style-type: none"> ○ <u>For evaluation of dysfunctional temporomandibular joint after unsuccessful conservative therapy for at least four (4) weeks with bite block or splint and anti-inflammatory medicine</u>
<u>May 2019</u>	<ul style="list-style-type: none"> • <u>Updated background information and references</u>

May 2019

- ~~Updated background information and references~~

May 2020

Added:

- For evaluation of temporomandibular joint dysfunction (TMD) with suspected internal joint derangement with ALL of the following
 - Persistent symptoms of facial or jaw pain, restricted range of motion, pain and/or noise with TMJ function (i.e., chewing)
 - Conservative therapy with a trial of anti-inflammatory AND behavioral modification has been unsuccessful for at least four (4) weeks
 - Initial X-rays have been performed
- For evaluation of Juvenile idiopathic arthritis (JIA)
- Abnormal initial x ray or ultrasound needing additional imaging

-

Deleted:

- Locked or Frozen Jaw
 - For evaluation of dysfunctional temporomandibular joint after unsuccessful conservative therapy for at least four (4) weeks with bite block or splint and anti-inflammatory medicine

REFERENCES

Aiken A, Bouloux G, Hudgins P. MR imaging of the temporomandibular joint. *Magn Reson Imaging Clin N Am*. 2012 Aug;20(3):397-412. doi: 10.1016/j.mric.2012.05.002. Epub 2012 Jul 15. Review.

American Society of Temporomandibular Joint Surgeons (ASTMJS). Guidelines for Diagnosis and Management of Disorders involving the Temporomandibular Joint and related Musculoskeletal Structures. <http://astmjs.org/final%20guidelines-04-27-2005.pdf>. Published 2001.

Armijo-Olivo S, Pitance L, Singh V, et al. Effectiveness of manual therapy and therapeutic exercise for temporomandibular disorders: Systematic review and meta-analysis. *Phys Ther*. 2016; 96(1):9-25. doi:10.2522/ptj.20140548.

Bag AK, Gaddikeri S, Singhal A, et al. Imaging of the temporomandibular joint: An update. *World J Radiol*. 2014; 6(8):567-582. <http://doi.org/10.4329/wjr.v6.i8.567>.

Ebrahim S, Montoya L, Busse JW, et al. The effectiveness of splint therapy in patients with temporomandibular disorders: A systematic review and meta-analysis. *J Am Dent Assoc*. 2012 Aug; 143(8):847-57.

Gauer RL, Semidey MJ. Diagnosis and treatment of temporomandibular disorders. *Am Fam Physician*. 2015; 91(6):378-386. <https://www.aafp.org/afp/2015/0315/p378.html>.

Granquist E. Treatment of temporomandibular joint in a child with juvenile inflammatory arthritis. *Oral Maxillofac Surg Clin North Am*. 2018 Feb; 30(1):97-107.

Hoffman D, Puig L. Complications of TMJ surgery. *Oral Maxillofac Surg Clin North Am*. 2015 Feb; 27(1):109-24.

~~Murphy MK, MacBarb RF, Wong ME, et al. Temporomandibular disorders: a review of etiology, clinical management, and tissue engineering strategies. *Int J Oral Maxillofac Implants*. 2013; 28(6):e393-e414. doi:10.11607/jomi.te20~~

Petscavage-Thomas JM, Walker EA. Unlocking the jaw: Advanced imaging of the temporomandibular joint. *Am J Roentgenol*. 2014; 203(5):1047-1058.

Pficer JK, Dodic S, Lazic V, et al. Occlusal stabilization splint for patients with temporomandibular disorders: Meta-analysis of short and long term effects. *PLoS One*. 2017; 12(2):e0171296. <https://doi.org/10.1371/journal.pone.0171296>.

Stoll ML, Guleria S, Mannion ML, et al. Defining the normal appearance of the temporomandibular joints by magnetic resonance imaging with contrast: A comparative study of children with and without juvenile idiopathic arthritis. *Pediatr Rheumatol Online J*. 2018; 16(8).

Tu KH, Chuang HJ, Lai LA, et al. Ultrasound Imaging for Temporomandibular Joint Disc Anterior Displacement. *J Med Ultrasound*. 2018;26(2):109–110. doi:10.4103/JMU.JMU_18_18.

[Reviewed / Approved by NIA Clinical Guideline Committee](#)

GENERAL INFORMATION

It is an expectation that all patients receive care/services from a licensed clinician. All appropriate supporting documentation, including recent pertinent office visit notes, laboratory data, and results of any special testing must be provided. If applicable: All prior relevant imaging results and the reason that alternative imaging cannot be performed must be included in the documentation submitted.

Reviewed / Approved by  M. Atif Khalid, M.D., Medical Director, Radiology

Disclaimer: Magellan Healthcare service authorization policies do not constitute medical advice and are not intended to govern or otherwise influence the practice of medicine. These policies are not meant to supplant your normal procedures, evaluation, diagnosis, treatment and/or care plans for your patients. Your professional judgement must be exercised and followed in all respects with regard to the treatment and care of your patients. These policies apply to all Magellan Healthcare subsidiaries including, but not limited to, National Imaging Associates (“Magellan”). The policies constitute only the reimbursement and coverage guidelines of Magellan. Coverage for services varies for individual members in accordance with the terms and conditions of applicable Certificates of Coverage, Summary Plan Descriptions, or contracts with governing regulatory agencies. Magellan reserves the right to review and update the guidelines at its sole discretion. Notice of such changes, if necessary, shall be provided in accordance with the terms and conditions of provider agreements and any applicable laws or regulations.