

National Imaging Associates, Inc.	
Clinical guidelines	Original Date: November 2007
ORBIT, FACE, NECK, SINUS MRI	
CPT Codes: 70540, 70542, 70543	Last Revised Date: May April 20210
Guideline Number: NIA_CG_014	Implementation Date: January 20221



A single authorization for CPT code 70540, 70542, or 70543 includes imaging of the Orbit, Face, Sinuses, and Neck. Multiple authorizations are not required.

## **AmeriHealth Caritas Louisiana**

National Imaging Associates, Inc.*	
Clinical guidelines	Original Date: November 2007
ORBIT, FACE, NECK, SINUS MRI	
CPT Codes: 70540, 70542, 70543	Last Revised Date: April 2021
Guideline Number: NIA CG 014	Implementation Date: January 2022

#### **INDICATIONS FOR ORBIT MRI**

MRI is superior for the evaluation of the visual pathways, globe and soft tissues<sub>27</sub> CT is preferred for visualizing bony detail and calcifications (Hande, 2012; Kennedy, 2018)

#### Abnormal external or direct eye exam

- Exophthalmos (proptosis) or enophthalmos
- o Ophthalmoplegia with concern for orbital pathology
- Unilateral optic disk swelling (Hata, 2017; Margolin, 2019; Passi, 2013)
- Documented visual field defect (Fadzil, 2013; Kedar, 2011; Prasad, 2012; Sadun, 2011)

<sup>\*</sup> National Imaging Associates, Inc. (NIA) is a subsidiary of Magellan Healthcare, Inc.



- Unilateral or with optic disc abnormaliabnormaliabnormality optic disc(s) (-e.g., optic disc blurring, edema, or pallor); AND
- (i.e. optic disc blurring, edema, or pallor); AND
- \_\_\_\_Not explained by underlying diagnosis, glaucoma, or macular degeneration

#### • Optic <u>n</u>Neuritis

(CMSC, 2018; Gala, 2015; Srikajon, 2018; Voss, 2011; CMSC, 2018)

- If atypical presentation, severe visual impairment, or poor recovery following initial onset or treatment onset <u>OR</u>
- o If needed to confirm optic neuritis and rule out compressive lesions(CMSC, 2018)

#### Orbital trauma

(Lin, 2012; Sung, 2014)

- o Physical findings of direct eye injury
- o Suspected orbital trauma with indeterminate x-ray or ultrasound
- Orbital or ocular mass/tumor, suspected or known (Hande, 2012; Kedar, 2011)
- Clinical <u>s</u>Suspicion of orbital infection (Hande, 2012; Kennedy, 2018)

#### • Clinical <u>s</u>Suspicion of osteomyelitis

(Arunkumar, 2011; Lee, 2016)

- o Direct visualization of bony deformity OR
- o Abnormal x-rays
- Clinical <u>s</u>Suspicion of Orbital Inflammatory Disease (e.g., eye pain and restricted eye movement with suspected orbital pseudotumor) (Pakdaman, 2014)
- Congenital orbital anomalies
- <u>Complex strabismus to aid in diagnosis, treatment and/or surgical planning</u> (<u>Demer, 2002; Kadom, 2008; Demer, 2002</u>)

NOTE: FOR OTHER ORBIT MRI INDICATIONS, CLICK HERE

**INDICATIONS FOR ORBIT AND BRAIN MRI COMBINATION STUDIES:** 



- Optic neuropathy or unilateral optic disk swelling Unilateral optic disk swelling/optic neuropathy of unclear etiology to distinguish between a compressive lesion of the optic nerve, optic neuritis, ischemic optic neuropathy (arteritic or non-arteritic), central retinal vein occlusion or optic nerve infiltrative disorders (Behbehani, 2007)
- Bilateral optic disk swelling (papilledema) with vision loss (Margolin, 2019)
- Optic nNeuritis if atypical presentation, severe visual impairment, or poor recovery following initial onset or treatment onset (CMSC, 2018)
- Known or suspected neuromyelitis optica spectrum disorder with severe, recurrent, or bilateral optic neuritis (Wingerchuk, 2015)
- For approved indications as noted above and being performed in a child under 8 years of age who will need anesthesia for the procedure and there is a suspicion of concurrent intracranial pathology (Lawson, 2000).

#### **INDICATIONS FOR FACE/SINUS MRI:**

- Rhinosinusitis (Kirsch, 2017)
  - Clinical <u>s</u>Suspicion of fungal infection (Gavito-Higuera, 2016)
  - Clinical <u>s</u>-suspicion of orbital or intracranial complications (Arunkumar, 2011; Lee, 2016), such as (Arunkumar, 2011; Lee, 2016)
    - Preseptal, orbital, or central nervous system infection
    - Osteomyelitis
    - Cavernous sinus thrombosis
- Sinonasal obstruction, suspected-mass, based on exam, nasal endoscopy, or prior imaging (Kirsch, 2017; Rosenfeld, 2015)
- Suspected infection
  - Osteomyelitis (after x-rays) (Pincus, 2009)
  - o Abscess
- Anosmia <u>or Dysosmia based</u> on objective testing that is persistent and of unknown origin (Policeni, 2017; Rouby, 2011; Zaghouani, 2013)
- Granulomatosis with polyangiitis (Wegener's granulomatosis) disease (Pakalniskis, 2015)
- Face mass

(Kirsch, 2017; Koeller, 2016; Kuno, 2014):

- Present on physical exam and remains non-diagnostic after x-ray or ultrasound is completed (Kuno, 2014)
- Known or highly suspected head and neck cancer on examination (Kirsch, 2017)



• Failed 2 weeks of treatment for suspected infectious adenopathy (Haynes, 2015)-

#### Facial trauma

(Echo, 2010; Lin, 2012; Raju, 2017; Sung, 2014)

- Physical findings of direct facial bone injury
- o For further evaluation of a known fracture for treatment or surgical planning

Note:

CSF (cerebrospinal fluid) rhinorrhea - Sinus CT is indicated when looking to \_\_\_\_\_\_characterize a bony defect. CSF otorrhea - Temporal Bone CT is indicated. For \_\_\_\_\_\_intermittent leaks and complex cases, consider CT/MRI/Nuclear Cisternography). CSF -fluid should always be confirmed with laboratory testing (Beta-2 transferrin assay) -edPost traumatic CSF rhinorrhea (for CSF otorrhea Temporal Bone imaging is recommended) (Oh, 2017; Snetty, 2015) \_\_\_\_\_)ed

- Trigeminal neuralgia/neuropathy (for evaluation of the extracranial nerve course)
  - If < 40 years of age or atypical features (e.g., bilateral, hearing loss, dizziness/vertigo, visual changes, sensory loss, numbness, pain > 2min, pain outside trigeminal nerve distribution, progression) (ACR, 2017; Hughes, 2016; Policeni, 2017)

#### NOTE: FOR OTHER FACE/SINUS MRI INDICATIONS, CLICK HERE

#### **INDICATIONS FOR FACE/SINUS AND BRAIN MRI COMBINATION STUDIES:**

- Anosmia or -ondysosmia on objective testing that is persistent and of unknown origin (ACR, 2017; Decker, 2013; Policeni, 2017; Zaghouani, 2013, Decker 2013))
- Granulomatosis with polyangiitis (Wegener's granulomatosis) disease (Pakalniskis, 2015)
- Trigeminal neuralgia that meets the above criteria (Hughes 2016; Policeni, 2017)
- For approved indications as noted above and being performed in a child under 8 years of age who will need anesthesia for the procedure and there is a suspicion of concurrent intracranial pathology (Lawson, 2000).

#### INDICATIONS FOR NECK MRI:

Suspected tumor or cancer (ACR, 2018a): (ACR, 2018a)

- Suspicious lesions in mouth or throat (Kuno, 2014).
- Suspicious mass/tumor found on another imaging study and needing clarification Neck mass <u>or lymphadenopathy</u> (non-parotid or thyroid)

•



## • Present on physical exam and remains non-diagnostic after ultrasound is completed (Kuno, 2014)

Note: For discrete cystic lesions of the neck, an ultrasound should be performed as initial imaging unless there is a high suspicion of malignancy

Present on physical exam and remains non-diagnostic after ultrasound is completed (Kuno, 2014)

**NOTE:** For discrete cystic lesions of the neck, an ultrasound should be performed as initial imaging unless there is a high suspicion of malignancy

- Increased risk for malignancy -with one or more of the following findings (Pynnonen, 2017):
  - Fixation to adjacent tissues
  - Firm consistency
  - Size >1.5 cm
  - Ulceration of overlying skin
  - Mass present ≥ two weeks (or uncertain duration) without significant fluctuation and not considered of infectious cause
  - History of cancer
- Failed 2 weeks of treatment for suspected infectious adenopathy (Haynes, 2015).
- Neck Mass (parotid) (ACR, 2018a)
  - Parotid mass found on other imaging study and needing further evaluation (US is the initial imaging study of a parotid region mass)
- Neck Mass (thyroid) US is the initial imaging study of a thyroid region mass. CT is preferred to MRI in the evaluation of thyroid masses since there is less respiratory motion artifact (ACR, 2018b)
  - Staging and monitoring for recurrence of known thyroid cancer (ACR, 2018b).
  - To assess extent of thyroid tissue when other imaging suggests extension through the thoracic inlet into the mediastinum or concern for airway compression (Gharib 2016; Lin, 2016)

Note: US is the initial imaging study of a thyroid region mass. CT is preferred over MRI in the evaluation of thyroid masses since there is less respiratory motion artifact. Chest CT may be included for preoperative assessment in some cases NOTE: Chest CT may be included for preoperative assessment in some cases

#### Pediatric patients (≤ 18 years old) (Wai, 2020):

<u>(Wai, 2020)</u>

- Neck masses-in the pediatric population if ultrasound is inconclusive or suspicious (Brown, 2016)
- History of malignancy

#### Known or suspected deep space infections or abscesses of the pharynx or neck



(Meyer, 2009)

#### Other indications for a Neck MRI:

- MR Sialography to evaluate salivary ducts (Burke, 2011; Ren, 2015)
- Vocal cord lesions or vocal cord paralysis (Dankbaar, 2014).
- Unexplained ear pain when ordered by a specialist with all of the following (Earwood, 2018)
  - **Otoscopic exam, nasolaryngoscopy, lab evaluation (ESR, CBC) AND**
  - Risk factor for malignancy i.e., tobacco use, alcohol use, dysphagia, weight loss OR age older than 50 years
- •----
- Diagnosed primary hyperparathyroidism when surgery is planned
  - Previous nondiagnostic ultrasound or nuclear medicine scan (Khan, 2014; Piciucchi, 2012).
- Bell's palsy/hemifacial spasm (for evaluation of the extracranial nerve course)
  - If atypical signs, slow resolution beyond three weeks, no improvement at four months, or facial twitching/spasms prior to onset (Quesnel, 2010)
- Objective cranial nerve palsy (CN IX-XII) (for evaluation of the extracranial nerve course) (ACR, 2017; Mumtaz, 2014; Policeni, 2017)
- Brachial **p**Plexopathy if mechanism of injury or EMG/NCV studies are suggestive (Vijayasarathi, 2016)

Note: Chest MRI is preferred study, but neck and/or shoulder (upper extremity) MRI can be ordered depending on the suspected location of injury

NOTE: FOR OTHER NECK MRI INDICATIONS, CLICK HERE

**INDICATIONS FOR NECK AND BRAIN MRI COMBINATION STUDIES:** 

- Objective cranial nerve palsy (CN IX-XII) (for evaluation of the extracranial nerve course) (ACR, 2017; Mumtaz, 2014; Policeni, 2017)
- For approved indications as noted above and being performed in a child under 8 years of age who will need anesthesia for the procedure and there is a suspicion of concurrent intracranial pathology (Lawson, 2000).

#### **OTHER INDICATIONS FOR ORBIT/FACE/SINUS/NECK MRI**

Known tumor or cancer of skull base, orbits, sinuses, face, tongue, larynx, nasopharynx, pharynx, or salivary glands

• Initial staging (Kuno, 201<u>4</u>3)



- Restaging during treatment
- Suspected recurrence or new metastases based on symptoms or examination findings
  - o New mass
  - Change in lymph nodes (Hoang, 2013)
- Surveillance appropriate for tumor type and stage

Indication for combination studies for the initial pre-therapy staging of cancer, OR active monitoring for recurrence as clinically indicated OR evaluation of suspected metastases:

 < 5 concurrent studies to include CT or MRI of any of the following areas as appropriate depending on the cancer: Neck, Abdomen, Pelvis, Chest, Brain, Cervical Spine, Thoracic Spine or Lumbar Spine.

#### Pre-operative/procedural evaluation:

<u>Pre-operative evaluation for a planned surgery or procedure if the imaging provides diagnostic</u> <u>information that is not available on prior studies (provider should be referred to the health plan for</u> <u>nondiagnostic surgical planning studies).</u>Pre-operative evaluation for a planned surgery or procedure.

•

#### Post- operative/procedural evaluation:

 When imaging, physical, or laboratory findings indicate surgical or procedural complications.

#### INDICATIONS FOR COMBINATION STUDIES: ORBIT/FACE/ SINUS/NECK MRI WITH BRAIN MRI.

• For approved indications as noted above and being performed in a child under 8 years of age who will need anesthesia for the procedure and there is a suspicion of concurrent intracranial pathology (Lawson, 2000).

#### **Brain and Orbit MRI**

- Unilateral optic disk swelling/optic neuropathy of unclear etiology to distinguish between a compressive lesion of the optic nerve, optic neuritis, ischemic optic neuropathy (arteritic or non-arteritic), central retinal vein occlusion or optic nerve infitrative disorders (Behbehani, 2007)
- Bilateral optic disk swelling (papilledema) with vision loss (Margolin, 2019)
- Optic Neuritis if atypical presentation, severe visual impairment, or poor recovery following initial onset or treatment onset (CMSC, 2018)
- Known or suspected neuromyelitis optica spectrum disorder with severe, recurrent, or bilateral optic neuritis (Wingerchuk, 2015)

#### **Brain and Sinus MRI**

 Anosmia on objective testing that is persistent and of unknown origin (ACR, 2017; Policeni, 2017; Zaghouani, 2013)



Granulomatosis with polyangiitis (Wegener's granulomatosis) disease (Pakalniskis, 2015)

#### Brain and Neck MRI

 Objective cranial nerve palsy (CN IX XII) (for evaluation of the extracranial nerve course) (ACR, 2017; Mumtaz, 2014; Policeni, 2017)

#### BACKGROUND:

Magnetic resonance imaging (MRI) is used in the evaluation of face and neck region masses, trauma, and infection. The soft-tissue contrast between normal and abnormal tissues provided by MRI is sensitive for differentiating between inflammatory disease and malignant tumors and permits the precise delineation of tumor margins. MRI is used for therapy planning and follow-up of face and neck neoplasms. It is also used for the evaluation of neck lymphadenopathy and vocal cord lesions.

CT scanning remains the study of choice for the imaging evaluation of acute and chronic inflammatory diseases of the sinonasal cavities. MRI is not considered the first-line study for routine sinus imaging because of limitations in the definition of the bony anatomy and length of imaging time. MRI for confirmation of diagnosis of sinusitis is discouraged because of hypersensitivity (overdiagnosis) in comparison to CT without contrast. MRI, however, is superior to CT in differentiating inflammatory conditions from neoplastic processes. MRI may better depict intraorbital and intracranial complications in cases of aggressive sinus infection, as well as differentiating soft-tissue masses from inflammatory mucosal disease. MRI may also identify fungal invasive sinusitis or encephaloceles.

Anosmia - Nonstructural causes of anosmia include post viral symptoms, medications (Amitriptyline, Enalapril, Nifedipine, Propranolol, Penicillamine, Sumatriptan, Cisplatin, Trifleuoperazine, Propylthiouracil). These should be considered prior to advanced imaging to look for a structural cause. <u>Anosmia and dysgeusia have been reported as common early</u> <u>symptoms in patients with COVID-19, occurring in greater than 80 percent of patients. For</u> <u>isolated anosmia, imaging is typically not needed once the diagnosis of COVID has been made</u> <u>given the high association. As such, COVID testing should be done prior to imaging (Geyer,</u> <u>2008; Lechien, 2020; Saniasiaya, 2020).</u>

#### POLICY HISTORY:

<u>Date</u>	Summary
<u>May 2021</u>	Updated References
	Reordered Indications
	Added hyperlinks to OTHER indications
	Orbit



Added:
<ul> <li><u>Complex strabismus to ald in diagnosis, treatment and/or</u></li> </ul>
If needed to confirm ontic neuritis and rule out compressive
Clarified:
Documented visual defect if MRI is contraindicated or cannot
be performed - Unilateral or with abnormal optic disc(s) (i.e. Optic
disc blurring, edema, or pallor);
•
<ul> <li>Clinical Suspicion of osteomyelitis: Direct visualization of bony</li> </ul>
deformity OR Abnormal X-rays
•
<ul> <li>Optic neuropathy or unilateral optic disk swelling of unclear</li> </ul>
etiology (Combo Orbit/Brain CT)
Sinus/Face
Added:
Facial Trauma- For further evaluation of a known fracture for
treatment or surgical planning
• Dysosmia
Clarified:
• Sinonasal obstruction, suspected mass, based on exam, nasal
<u>endoscopy, or prior imaging</u>
Note: CSF (cerebrospinal fluid) rhinorrhea - Sinus CT is indicated
when looking to characterize a bony defect. CSF otorrhea -
Temporal Bone CT is indicated. For intermittent leaks and complex
cases consider CT/MRI/Nuclear Cisternography). CSF fluid should
always be confirmed with laboratory testing (Beta-2 transferrin
<u>assay)</u>
Deleted:
• Trigominal nouralgia if $A_{32} < 40$
<u>     Trigeninal neuraigia = n Age &lt; 40</u>
Neck
Auucu.



	<ul> <li>Unexplained ear pain when ordered by a specialist with all the</li> </ul>
	<u>following (Earwood, 2018)</u>
	•
	Otoscopic exam, nasolaryngoscopy, lab evaluation (ESR,
	CBC) AND
	<ul> <li>Risk factor for malignancy ie tobacco use, alcohol use,</li> </ul>
	dysphagia, weight loss OR age older than 50 years
	Brachial Dievenathy (Vijayasarathi, 2016) if mechanism of injury or
	ENG/NCV studies are suggestive
	Note: Chest MRI is preferred study, but neck and/or shoulder
	(upper extremity) MRI can be ordered depending on the
	suspected location of injury
	All
	Clarified:
	<ul> <li><u>Clarified: Pre-operative evaluation for a planned surgery or</u></li> </ul>
	procedure if the imaging provides diagnostic information that is not
	available on prior studies (provider should be referred to the health
	plan for nondiagnostic surgical planning studies)
	—— <u>Removed statement: A single authorization for CPT code 70540,</u>
	70542, or 70543 includes imaging of the Orbit, Face, Sinuses, and
	Neck. Multiple authorizations are not required
<u>May 2020</u>	Clarified:
	<u>Orbit</u>
	Ophthalmoplegia with concern for orbital pathology
	<ul> <li>Documented visual field defect if MRI is contraindicated or cannot</li> </ul>
	be performed
	Orbital or ocular mass/tumor suspected or known
	Clinical Suspicion of orbital infection
	Clinical Suspicion of Orbital Information: Disease (a.c. ava nain
	<ul> <li>Clinical Suspicion of Orbital Inflammatory Disease (e.g., eye pain and restricted ave movement with suspected orbital</li> </ul>
	and restricted eye movement with suspected orbital
	<u>Face/Sinus</u>
	Suspected infection
	<ul> <li>Osteomyelitis (after x-rays)</li> </ul>
	<u>o Abscess</u>



# NIA Magellan...

numbness, pain > 2min, pain outside trigeminal nerve
distribution, progression)
Added:
Neck
<ul> <li>Suspicious mass/tumor found on another imaging study and</li> </ul>
needing clarification
Under increased risk for malignancy
• History of cancer
O Mass present ≥ two weeks (or uncertain duration) without
significant fluctuation and not considered of infectious
<u>cause</u>
Neck Mass (parotid)
<ul> <li>Parotid mass found on other imaging study and needing</li> </ul>
further evaluation
Added:
Neck
<ul> <li>Neck Mass (thyroid) - US is the initial imaging study of a thyroid</li> </ul>
region mass. CT is preferred over MRI in the evaluation of thyroid
masses since there is less respiratory motion artifact
<ul> <li>Staging and monitoring for recurrence of known thyroid</li> </ul>
<u>cancer</u>
<ul> <li>To assess extent of thyroid tissue when other imaging</li> </ul>
suggests extension through the thoracic inlet into the
mediastinum or concern for airway compression (Lin, 2016;
Gnarib 2016) NOTE: Chest CT may be included for preeperative
assossment in some cases
Bodiatric patients (<18 years old)
<ul> <li>Neck masses in the pediatric population if ultrasound is</li> </ul>
inconclusive or suspicious
<ul> <li>History of malignancy</li> </ul>
Added:
Neck
<ul> <li>Known or suspected deep space infections or abscesses of the</li> </ul>
pharynx or neck
Combo
• Known tumor or cancer of skull base, orbits, sinuses, face, tongue,
larynx, nasopharynx, pharynx, or salivary glands
<ul> <li>Surveillance appropriate for tumor type and stage</li> </ul>
• For approved indications as noted above and being performed in a
child under 8 years of age who will need anesthesia for the



	procedure and there is a suspicion of concurrent intracranial
	pathology
	Added:
	Combo
	Added sub Combo sections
	• Brain and Orbit
	<ul> <li>Optic Neuritis if atypical presentation, severe visual</li> </ul>
	impairment or poor recovery following initial onset
	or treatment onset
	<ul> <li>Brain and Sinus</li> </ul>
	<ul> <li>Brain and Neck</li> </ul>
	Deleted:
	<u>Orbit</u>
	Unilateral optic disk swelling papilledema approve dedicated
	Orbits MRI even if Brain MRI approved
	Deleted:
	Face/Sinus
	Clinical Suspicion of osteomyelitis
	<ul> <li>Direct visualization of lesion over bone</li> </ul>
	<ul> <li>Abnormal x-ray</li> </ul>
	Face Mass
	<ul> <li>Prior history of tumor with suspicion of recurrence</li> </ul>
	Facial trauma
	<ul> <li>Suspected orbital trauma with indeterminate x-ray or</li> </ul>
	ultrasound
	Neck
	Palpable from Palpable suspicious lesions in mouth or throat
	<ul> <li>Salivary gland stones or clinical concern for abscess</li> </ul>
	Thoracic Outlet Syndrome
	Combo
	Trigeminal neuralgia
	Cranial neuropathy (weakness or sensory abnormalities of the
	head and neck
July 2019	ORBIT MRI:
	Removed: Orbital asymmetry and Suspected hyperthyroidism
	(such as Graves' disease)
	Added: Clinical suspicion of osteomyelitis
	Face/Sinus MRI
	Added specifics to Face Mass:
	O Present on physical exam and remains non-diagnostic after
	x-ray or ultrasound is completed (Kuno, 2014)



<ul> <li>Clinical concern for abscess</li> </ul>
<ul> <li>Failed 2 weeks of treatment for suspected infectious</li> </ul>
adenopathy (Haynes, 2015).
• Prior history of tumor with suspicion of recurrence
Added: Facial trauma with physical findings of direct facial bone
injury: suspected orbital trauma w/indeterminate x-ray or US: CSF
leak (rhinorrhea or otorrhea)
Other Indications
<ul> <li>Added: Suspected recurrence or new metastases based on</li> </ul>
Added. Suspected recurrence of new metastases based on
symptoms or examination findings with new mass or change in
lymph nodes; Anosmia on objective testing; Trigeminal neuralgia
if <40 years of age or atypical features; Objective cranial nerve
palsy: and Granulomatosis with polyangiitis (Wegener's
granulomatosis) disease
Indications for combo studies orbit/face/sinus neck MRI with brain
MRI
WIN
Added: Bilateral papilledema with vision loss AND Known or
suspected neuromyelitis optica spectrum disorder with severe,
recurrent, or bilateral optic neuritis



#### Review Date: July 2019 Review Summary:

#### ORBIT MRI:

- Removed: Orbital asymmetry and Suspected hyperthyroidism (such as Graves' disease)
- Added: Clinical suspicion of osteomyelitis

#### Face/Sinus MRI

- Added specifics to Face Mass:
  - Present on physical exam and remains non-diagnostic after x ray or ultrasound is completed (Kuno, 2014)
  - ⊖ Clinical concern for abscess
  - Failed 2 weeks of treatment for suspected infectious adenopathy (Haynes, 2015).
  - Prior history of tumor with suspicion of recurrence
- Added: Facial trauma with physical findings of direct facial bone injury; suspected orbital trauma w/indeterminate x-ray or US; CSF leak (rhinorrhea or otorrhea)

#### **Other Indications**

 Added: Suspected recurrence or new metastases based on symptoms or examination findings with new mass or change in lymph nodes; Anosmia on objective testing; Trigeminal neuralgia if <40 years of age or atypical features; Objective cranial nerve palsy; and Granulomatosis with polyangiitis (Wegener's granulomatosis) disease

Indications for combo studies orbit/face/sinus neck MRI with brain MRI

 Added: Bilateral papilledema with vision loss AND Known or suspected neuromyelitis optica spectrum disorder with severe, recurrent, or bilateral optic neuritis

#### Review Date: May 2020

**Review Summary:** 

**Clarified:** 

**Orbit** 

- Ophthalmoplegia with concern for orbital pathology
- Documented visual field defect if MRI is contraindicated or cannot be performed
- Orbital or ocular mass/tumor, suspected or known
- Clinical Suspicion of orbital infection
- Clinical Suspicion of Orbital Inflammatory Disease (e.g., eye pain and restricted eye
  movement with suspected orbital pseudotumor)

#### Face/Sinus

- Suspected infection
  - Osteomyelitis (after x-rays)
  - ⊖ Abscess
- Facial Trauma



- Post traumatic CSF rhinorrhea (for CSF otorrhea Temporal Bone imaging is recommended)
- Anosmia on objective testing that is persistent and of unknown origin (also in Brain and Sinus combo section)

#### Neck

- Neck mass (non-parotid or thyroid)
  - Note: For discrete cystic lesions of the neck, an ultrasound should be performed as initial imaging unless there is a high suspicion of malignancy
- MR Sialography to evaluate salivary ducts
- Objective cranial nerve palsy (CN IX-XII) (for evaluation of the extracranial nerve course)
   (also in Brain and Neck combo section)
- **Combo Brain and Orbit**
- Reworded: Unilateral optic disk swelling/optic neuropathy of unclear etiology to distinguish between a compressive lesion of the optic nerve, optic neuritis, ischemic optic neuropathy (arteritic or non-arteritic), central retinal vein occlusion or optic nerve infiltrative disorders
- Bilateral optic disk swelling (papilledema) with vision loss

#### Added:

#### **Orbit**

- MRI is superior for the evaluation of the visual pathways, globe and soft tissues, CT is preferred for visualizing bony detail and calcifications
- Unilateral optic disk swelling
- Under documented visual field defect
  - Unilateral or with optic disc abnormality
- Congenital orbital anomalies

#### Added:

#### Face/Sinus

- Examples of orbital or intracranial complications
  - Preseptal, orbital, or central nervous system infection
  - ─ Osteomyelitis
  - ⊖ Cavernous sinus thrombosis
- Face mass
  - Known or highly suspected head and neck cancer on examination
- Trigeminal neuralgia/neuropathy (for evaluation of the extracranial nerve course)
  - If < 40 years of age or atypical features (e.g. bilateral, hearing loss, dizziness/vertigo, visual changes, sensory loss, numbness, pain > 2min, pain outside trigeminal nerve distribution, progression)

#### <u>Added</u>:

#### **Neck**

- Suspicious mass/tumor found on another imaging study and needing clarification
- Under increased risk for malignancy
  - ⊖ History of cancer



- → Mass present ≥ two weeks (or uncertain duration) without significant fluctuation and not considered of infectious cause
- Neck Mass (parotid)
  - Parotid mass found on other imaging study and needing further evaluation

Added:

Neck

- Neck Mass (thyroid) US is the initial imaging study of a thyroid region mass. CT is preferred over MRI in the evaluation of thyroid masses since there is less respiratory motion artifact
  - Staging and monitoring for recurrence of known thyroid cancer
  - To assess extent of thyroid tissue when other imaging suggests extension through the thoracic inlet into the mediastinum or concern for airway compression (Lin, 2016; Gharib 2016)
- NOTE: Chest CT may be included for preoperative assessment in some cases
   Pediatric patients (≤18 years old)
  - Neck masses in the pediatric population if ultrasound is inconclusive or suspicious
     History of malignancy

### <u>Added</u>:

**Neck** 

- Known or suspected deep space infections or abscesses of the pharynx or neck
- Combo
- Known tumor or cancer of skull base, orbits, sinuses, face, tongue, larynx, nasopharynx, pharynx, or salivary glands
  - Surveillance appropriate for tumor type and stage
- For approved indications as noted above and being performed in a child under 8 years of age who will need anesthesia for the procedure and there is a suspicion of concurrent intracranial pathology

#### <u>Added</u>:

#### Combo

- Added sub Combo sections
  - ⊖ Brain and Orbit
    - Optic Neuritis if atypical presentation, severe visual impairment or poor recovery following initial onset or treatment onset
      - Brain and Sinus
      - Brain and Neck

#### **Deleted:**

#### **Orbit**

 Unilateral optic disk swelling papilledema approve dedicated Orbits MRI even if Brain MRI approved

#### Deleted:

#### Face/Sinus

- Clinical Suspicion of osteomyelitis
  - Direct visualization of lesion over bone

# NIA Magellan...

#### ⊖ Abnormal x-ray

- Face Mass
  - Prior history of tumor with suspicion of recurrence
- Facial trauma
  - Suspected orbital trauma with indeterminate x-ray or ultrasound

#### Neck

- Palpable from Palpable suspicious lesions in mouth or throat
- Salivary gland stones or clinical concern for abscess
- Thoracic Outlet Syndrome

#### **Combo**

- Trigeminal neuralgia
- Cranial neuropathy (weakness or sensory abnormalities of the head and neck



#### **REFERENCES**:

Abuabara A. Cerebrospinal fluid rhinorrhoea: diagnosis and management. *Med Oral Patol* Oral Cir Bucal. 2007;12(5):E397-400.

American College of Radiology (ACR). ACR Appropriateness Criteria<sup>®</sup> Cranial Neuropathy. 2017. https://acsearch.acr.org/docs/69509/Narrative/.

American College of Radiology (ACR). ACR Appropriateness Criteria<sup>®</sup> Plexopathy. 2016. https://acsearch.acr.org/docs/69487/Narrative/.

American College of Radiology (ACR). ACR Appropriateness Criteria<sup>®</sup> Neck Mass/Adenopathy 2018a. https://acsearch.acr.org/docs/69504/Narrative/.

American College of Radiology (ACR). ACR Appropriateness Criteria<sup>®</sup> Thyroid Disease. 2018b. https://acsearch.acr.org/docs/3102386/Narrative/.

Arunkumar JS, Naik AS, Prasad KC, et al. Role of nasal endoscopy in chronic osteomyelitis of maxilla and zygoma: A case report. *Case Rep Med*. 2011; Article ID 802964.

Behbehani R. Clinical approach to optic neuropathies. *Clin Ophthalmol*. 2007; 1(3):233-246.

Borges A, Casselman J. Imaging the trigeminal nerve. *Eur J Radiol*. 2010;74(2):323-340. doi:10.1016/j.ejrad.2010.02.006.

Brown RE, Harave S. Diagnostic imaging of benign and malignant neck masses in children – A pictorial review. *Quant Imaging Med Surg.* 2016 Oct; 6(5):591-604.

Burke CJ, Thomas RH, Howlett D. Imaging the major salivary glands. *Br J Oral Maxillofac Surg*. 2011; 49(4):261.

Consortium of Multiple Sclerosis Centers (CMSC). MRI protocol and clinical guidelines for the diagnosis and follow-up of MS. February 2018.

Dankbaar JW, Pameijer FA. Vocal cord paralysis: Anatomy, imaging and pathology. *Insights Imagings*. December 2014; 5(6):743-751.

Decker JR, Meen EK, Kern RC, Chandra RK. Cost effectiveness of magnetic resonance imaging in the workup of the dysosmia patient. *International Forum of Allergy & Rhinology*. 2013;3(1):56-61. doi:10.1002/alr.21066



Demer JL, Clark RA, Kono R, Wright W, Velez F, Rosenbaum AL. A 12-year, prospective study of extraocular muscle imaging in complex strabismus. *J AAPOS*. 2002;6(6):337-347. doi:10.1067/mpa.2002.129040.

Earwood JS, Rogers TS, Rathjen NA. Ear pain: Diagnosing common and uncommon causes. *Am Fam Physician*. 2018;97(1):20-27.

Dankbaar JW, van Bemmel AJ, Pameijer FA. Imaging findings of the orbital and intracranial complications of acute bacterial rhinosinusitis. *Insights Imaging*. 2015; 6(5):509-518.

Echo A, Troy JS, Hollier LH. Frontal sinus fractures. *Semin Plast Surg*. 2010; 24(4):375-382. http://doi.org/10.1055/s-0030-1269766. https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3324222/.

Fadzil F, Ramli N, Ramli NM. MRI of optic tract lesions: review and correlation with visual field defects. *Clin Radiol*. 2013 Oct; 68(10):e538-51.

Gala F. Magnetic resonance imaging of optic nerve. *Indian J Radiol Imaging*. 2015 Oct-Dec; 25(4):421-438.

Gavito-Higuera J, Mullins CB, Ramos-Duran L, et al. Sinonasal fungal infections and complications: A pictorial review. *J Clin Imaging Sci*. 2016 Jun 14; 6:23.

<u>Geyer M, Nilssen E. Evidence-based management of a patient with anosmia. *Clin Otolaryngol.* 2008; 33(5).</u>

Gharib H, Papini E, Garber JR, et al. American Association of Clinical Endocrinologists, American College of Endocrinology, and Associazione Medici Endocrinologi Medical Guidelines for Clinical Practice for the Diagnosis and Management of Thyroid Nodules – 2016 Update. *Endocr Pract*. 2016 May; 22(5):622-39.

Hande PC, Talwar I. Multimodality imaging of the orbit. *Indian J Radiol Imaging*. 2012; 22(3):227-239. http://doi.org/10.4103/0971-3026.107184.

Hata M, Miyamoto K. Causes and prognosis of unilateral and bilateral optic disc swelling. *Neuroophthalmology*. 2017 Aug; 41(4):187-191.

Haynes J, Arnold K, Aguirre-Oskins C, et al. Evaluation of neck masses in adults. *Am Fam Physician*. May 2015; 91(10):698-706. https://www.aafp.org/afp/2015/0515/p698.html.



Hata M, Miyamoto K. Causes and prognosis of unilateral and bilateral optic disc swelling. *Neuroophthalmology*. 2017 Aug; 41(4):187-191.

Hoang JK, Vanka J, Ludwig BJ, et al. Evaluation of cervical lymph nodes in head and neck cancer with CT and MRI: Tips, traps, and a systematic approach. *Am J Roentgenol*. 2013; 200:W17-W25.

Hughes MA, Frederickson AM, Branstetter BF, et al. MRI of the trigeminal nerve in patients with trigeminal neuralgia secondary to vascular compression. *Am J Roentgenol*. 2016; 206:595-600.

Kadom N. Pediatric strabismus imaging. *Current Opinion in Ophthalmology*. 2008;19(5):371-378. doi:10.1097/ICU.0b013e328309f165.

Kedar S, Ghate D, Corbett JJ. Visual fields in neuro-ophthalmology. *Indian J Ophthalmol*. 2011 Mar-Apr; 59(2):103-109.

Kennedy TA, Corey AS, et al. ACR Appropriateness Criteria Orbits Vision and Visual Loss. *J Am Coll Radiol.* 2018 May; 15(5S):S116-S131.

Khan MA, Rafiq S, Lanitis S, et al. Surgical treatment of primary hyperparathyroidism: Description of techniques and advances in the field. Indian J Surg. 2014; 76(4):308-315.

Kirsch CFE, Bykowski J, et al. ACR Appropriateness Criteria Sinonasal Disease. *J Am Coll Radiol.* 2017 Nov; 14(11S):S550-S559.

Koeller KK. Radiologic features of sinonasal tumors. *Head Neck Pathol*. 2016; 10(1):1-12.

Kransdorf MJ, Murphey MD, Wessell DE, et al. American College of Radiology (ACR) Appropriateness Criteria. Expert Panel on Musculoskeletal Imaging: Soft-Tissue Masses. https://acsearch.acr.org/docs/69434/Narrative/. Published 2017.

Kuno H, Onaya H, Fujii S, et al. Primary staging of laryngeal and hypopharyngeal cancer: CT, MR imaging and dual-energy CT. [Epub October 27, 2013]. Eur J Radiol. January 2014; 83(1):e23-35. https://www.ncbi.nlm.nih.gov/pubmed/24239239

Lawson, GR. Sedation of children for magnetic resonance imaging. *Archives Dis Childhood*. 2000; 82(2).

Lechien JR, Chiesa-Estomba CM, De Siati DR, et al. Olfactory and gustatory dysfunctions as a clinical presentation of mild-to-moderate forms of the coronavirus disease (COVID-19): a



multicenter European study. *Eur Arch Otorhinolaryngol*. 2020;277(8):2251-2261. doi:10.1007/s00405-020-05965-1.

Lee YJ, Sadigh S, Mankad K, et al. The imaging of osteomyelitis. *Quant Imaging Med Surg*. 2016; 6(2):184-198. http://doi.org/10.21037/qims.2016.04.01. https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4858469/.

Lin YS, Wu HY, Lee CW, et al. Surgical management of substernal goiters at a tertiary referral centre: A retrospective cohort study of 2,104 patients. *Int J Surg.* 2016 Mar; 27:46-52.

Lin KY, Ngai P, Echegoyen JC. Et. al. Imaging in orbital trauma. *Saudi J Ophthalmol*. 2012 Oct; 26(4):427-432.

Mantur M, Łukaszewicz-Zając M, Mroczko B, et al. Cerebrospinal fluid leakage—Reliable diagnostic methods. *Clinica Chimica Acta*. 2011;412(11-12):837-840. doi:10.1016/j.cca.2011.02.017

Margolin E. Swollen optic nerve: An approach to diagnosis and management. Pract Neurol. 2019 Jun 13. Epub ahead of print.

Meyer AC, Kimbrough TG, Finkelstein M, et al. Symptom duration and CT findings in pediatric deep neck infection. *Otolaryngol Head Neck Surg*. 2009; 140(2):183-186. doi: 10.1016/j.otohns.2008.11.005.

Mumtaz S, Jensen MB. Facial neuropathy with imaging enhancement of the facial nerve: A case report. *Future Neurol*. 2014; 9(6):571-576. doi:10.2217/fnl.14.55.

Oh JW, Kim SH, Whang K. Traumatic cerebrospinal fluid leak: Diagnosis and management. *Korean J Neurotrauma*. 2017; 13(2):63-67.

Pakalniskis MG, Berg AD, Policeni BA, et al. The many faces of granulomatosis with polyangiitis: A review of the head and neck imaging manifestations. *Am J Roentgenol*. 2015; 205:W619-W629.

Pakdaman MN, Sepahdari AR, Elkhanary SM. Orbital inflammatory disease: Pictorial review and differential diagnosis. *World J Radiol.* 2014 Apr 28; 6(4):106-115.

Passi N, Degnan AJ, Levy LM. MR imaging of papilledema and visual pathways: Effects of increased intracranial pressure and pathophysiologic mechanisms. *Am J Neuroradiol*. 2013 May; 34(5):919-924.

Piciucchi S, Barone D, Gavelli G, et al. Primary hyperparathyroidism: Imaging to pathology. *J Clin Imaging Sci.* 2012; 2:59.



Pincus DJ, Armstrong MB, Thaller SR. Osteomyelitis of the craniofacial skeleton. *Semin Plast Surg*. May 2009; 23(2):73-79.

Policeni B, Corey AS, Burns J, et al. American College of Radiology (ACR) Appropriateness Criteria. Expert Panel on Neurologic Imaging: Cranial Neuropathy. https://acsearch.acr.org/docs/69509/Narrative/. Published 2017.

Prasad S, Galetta SL. Approach to the patient with acute monocular visual loss. *Neurol Clin Pract*. 2012; 2(1):14-23.

Pynnonen MA, Gillespie MB, Roman B, et al. Clinical practice guideline: Evaluation of the Neck Mass in Adults. *Otolaryngol Head Neck Surg.* 2017; 157(2\_suppl):S1.

Quesnel AM, Lindsay RW, Hadlock TA. When the bell tolls on Bell's palsy: Finding occult malignancy in acute-onset facial paralysis. Am J Otolaryngol. 2010 Sep-Oct; 31(5):339-42. Epub 2009 Jun 24.

Raju N S, Ishwar P, Banerjee R. Role of multislice computed tomography and three-dimensional rendering in the evaluation of maxillofacial injuries. *J Oral Maxillofac Radiol*. 2017; 5:67-73.

Razek AA, Huang BY. Soft Tissue Tumors of the Head and Neck: Imaging-based Review of the WHO Classification. *RadioGraphics*. 2011; 31:7:1923-1954. http://pubs.rsna.org/doi/pdf/10.1148/rg.317115095.

Ren YD, Li XR, Zhang J, et al. Conventional MRI techniques combined with MR sialography on T2-3D-DRIVE in Sjögren syndrome. *Int J Clin Exp Med*. 2015; 8(3):3974-3982.

Romano N, Federici M, Castaldi A. Imaging of cranial nerves: a pictorial overview. *Insights Imaging*. 2019 Mar 15; 10(1):33. doi:10.1186/s13244-019-0719-5.

Rosenfeld RM, Piccirillo JF, Chandrasekhar SS, et al. American Academy of Otolaryngology-Head and Neck Surgery. Clinical Practice Guideline (update): Adult Sinusitis. *Otolaryngol Head Neck Surg*. 2015; 152(2 Suppl):S1-S39.

Rouby C, Thomas-Danquin T, Vigouroux M, et al. The Lyon clinical olfactory test: Validation and measurement of hyposmia and anosmia in healthy and diseased populations. *Int J Otolaryngol*. 2011; Article ID 203805.

Sadun AA, Wang MY. Abnormalities of the optic disc. *Handb Clin Neurol*. 2011; 102:117-57. doi: 10.1016/B978-0-444-52903-9.00011-X.



Saniasiaya J, Islam MA, Abdullah B. Prevalence of olfactory dysfunction in coronavirus disease 2019 (COVID-19): a meta-analysis of 27,492 patients. *Laryngoscope*. 2021;131(4):865-878. doi:10.1002/lary.29286.

Salvolini U. Traumatic injuries: Imaging of facial injuries. Eur Radiol. 2002 Jun; 12(6):1253-61. Epub 2002 Apr 24.

Snetty VS, Reis MN, Aulino JM, et al. American College of Radiology (ACR) Appropriateness Criteria. Expert Panel on Neurologic Imaging: Head trauma. https://acsearch.acr.org/docs/69481/Narrative/. Published 2015.

Srikajon J, Siritho S, Ngamsombat C, et al. Differences in clinical features between optic neuritis in neuromyelitis optica spectrum disorders and in multiple sclerosis. *Mult Scler J Exp Transl Clin*. 2018 Aug 11; 4(3):2055217318791196. doi: 10.1177/2055217318791196.

Stalcup ST, Tuan AS, Hesselink JR. Intracranial causes of ophthalmoplegia: The visual reflex pathways. *RadioGraphics*. 2013; 33(5).

Sun JK, LeMay DR. Imaging of facial trauma. Neuroimaging Clin N Am. 2002 May; 12(2):295-309.

Sung EK, Nadgir RN, Fujita A, et al. Injuries of the globe: What can the radiologist offer? *RadioGraphics*. 2014; 34(3).

Tharin BD, Kini JA, York GE, et al. Brachial plexopathy: A review of traumatic and nontraumatic causes. *AJR Am J Roentgenol.* 2014; 202(1):W67.

<u>Vijayasarathi A, Chokshi C. MRI of the brachial Plexus: A practical review. Applied Radiology.</u> 2016 May: 9-18.

Voss E, Raab P, Trebst C, et al. Clinical approach to optic neuritis: Pitfalls, red flags and differential diagnosis. *Ther Adv Neurol Disord*. 2011; 4(2):123-134.

Wai K, Wang T, Lee E, et al. Management of persistent pediatric cervical lymphadenopathy. *Arch Otorhinolaryngol Head Neck Surg*. 2020; 4(1):1. DOI: 10.24983/scitemed.aohns.2020.00121.

Wingerchuk DM, Banwell B, Bennett JL, et al. International consensus diagnostic criteria for neuromyelitis optica spectrum disorders. *Neurology*. 2015; 85:177.



Zaghouani H, Slim I, Zina NB, et al. Kallmann syndrome: MRI findings. *Indian J Endocrinol Metab*. 2013; 17(Suppl 1):S142–S145. doi:10.4103/2230-8210.119536.

Reviewed / Approved by M. Arg Klarlic ML Be M. Atif Khalid, M.D., Medical Director, Radiology



#### **GENERAL INFORMATION**

It is an expectation that all patients receive care/services from a licensed clinician. All appropriate supporting documentation, including recent pertinent office visit notes, laboratory data, and results of any special testing must be provided. If applicable: All prior relevant imaging results and the reason that alternative imaging cannot be performed must be included in the documentation submitted.

**Disclaimer:** Magellan Healthcare service authorization policies do not constitute medical advice and are not intended to govern or otherwise influence the practice of medicine. These policies are not meant to supplant your normal procedures, evaluation, diagnosis, treatment and/or care plans for your patients. Your professional judgement must be exercised and followed in all respects with regard to the treatment and care of your patients. These policies apply to all Magellan Healthcare subsidiaries including, but not limited to, National Imaging Associates ("Magellan"). The policies constitute only the reimbursement and coverage guidelines of Magellan. Coverage for services varies for individual members in accordance with the terms and conditions of applicable Certificates of Coverage, Summary Plan Descriptions, or contracts with governing regulatory agencies. Magellan reserves the right to review and update the guidelines at its sole discretion. Notice of such changes, if necessary, shall be provided in accordance with the terms and conditions of provider agreements and any applicable laws or regulations.

