

Clinical Policy: Burn Surgery

Reference Number: LA.CP.MP.186

Date of Last Revision: ~~2/25~~1/26

[Coding Implications](#)

[Revision Log](#)

See [Important Reminder](#) at the end of this policy for important regulatory and legal information.

Description

Early burn debridement is vital to the overall survivability and outcome of burn patients. Early grafting is also directly related to improved survival rates.¹² Grafts used to cover the wound bed include skin replacements (i.e., autograft and allograft) and skin substitutes. Autografts (split or full thickness skin grafts) are the current standard of care in burn surgery. When the total body surface area is larger than the available donor sites or tissues are too edematous to allow successful acceptance of autografts, allograft and skin substitutes are an alternative. Skin substitutes are tissue-engineered products designed to replace, either temporarily or permanently, the form and function of the skin.⁹ This policy addresses the medical necessity criteria for burn debridement and/or excision and the use of skin substitutes for burns during the acute phase of treatment.

Note: For skin substitutes for chronic wounds, refer to *LA.CP.MP.185 Skin Substitutes for Chronic Wounds*.

Policy/Criteria

- I. It is the policy of Louisiana Healthcare Connections that burn treatment with debridement and/or excision is **medically necessary** for either of the following:
 - A. Deep partial-thickness burn;
 - B. Full-thickness burn or deeper.
- II. It is the policy of Louisiana Healthcare Connections that burn treatment with skin replacement/substitutes (including the procedure, product, service) is **medically necessary** when meeting all of the following:
 - A. Sufficient autograft is not available at the time of excision or is not feasible due to the physiological condition of the member/enrollee;
 - B. No evidence of burn wound infection;
 - C. Burn is either deep partial-thickness or full-thickness;
 - D. Treatment with any of the following skin replacement/substitutes:
 1. Allograft (human cadaver);
 2. Xenograft (porcine);
 3. Tissue-engineered skin substitute used according to FDA indications (e.g., Biobrane®, Transcyte®, Apligraf®, TheraSkin®, Integra® Wound Matrix, Integra® meshed Bilayer Wound Matrix, Integra® Dermal Regeneration Template, or Epicel®, if used per FDA ~~HDE~~); Humanitarian Device Exemption [HDE]).

Background

~~Johns Hopkins Health Library defines a burn as a type of painful wound caused by thermal, electrical, chemical, or electromagnetic energy. They cite smoking and open flame as the leading causes of burn injury for older adults and scalding as the leading cause of burn injury for~~

~~children.~~¹ A burn is defined as a traumatic injury to the skin or other organic tissue primarily caused by heat or exposure to electrical discharge, friction, chemicals, and radiation.¹⁵ According to the American Burn Association, burn injuries result in more than 500,000 hospital emergency department visits and approximately 50,000 acute admissions per year in the United States. and are a leading cause of accidental injury and death.^{2,15} The most severe burn injuries require admission to a specialty hospital or burn center.^{2,15}

~~A severe or major burn is classified as any burn that is accompanied by a major trauma, inhalation injury, or a chemical or high-voltage electrical burn. Also considered severe are any burns involving over 20 percent of the total body surface area (TBSA), with the exception of first-degree burns. Burns to high-risk individuals such as older adults, young children and anyone with a major comorbidity may be considered severe even if less than 20 percent of their TBSA is involved. Burns to areas like the eyes, ears, face, hands, feet or perineum may require specialized burn center care due to the high risk of functional impairment.~~³ In addition, circumferential burns of the extremities or thorax require a consultation with a burn center as they are an indicator of decreased blood flow. Deep circumferential burns of the chest may impair or prevent mechanical ventilation of the burn victim.⁴

Burns are classified in terms of degrees. First-degree burns, also called superficial partial thickness, only involve the outer layer of skin, the epidermis. These burns ~~will be~~ red and painful but remain dry and without blisters. First-degree burns typically heal within about one week. Second degree, or partial thickness burns, extend deeper into the dermis, include blisters, and have a wet appearance. Second-degree burns are extremely painful and can take two to three weeks to heal. Third-degree, or full thickness, burns have a white or leathery appearance and ~~will be~~ dry to the touch. These burns are often without sensation due to nerve damage. They extend the full depth of the skin. Skin grafts are typically required for healing third-degree burns. The most severe burns are called fourth-degree or are classified as with extension to deep tissues. These burns will extend to the muscles, tendons, and/or bone. Skin grafting and even more intensive surgeries or amputations may be required for healing.⁴

A severe or major burn is classified as any burn that is accompanied by a major trauma, inhalation injury, or a chemical or high-voltage electrical burn, and, generally, any burns involving over 20 percent of the total body surface area (TBSA), excluding first-degree burns.³ Burns to high-risk individuals such as older adults, young children and anyone with a major comorbidity may be considered severe even if less than 20 percent of their TBSA is involved. Burns to areas like the eyes, ears, face, hands, feet or perineum may require specialized burn center care due to the high risk of functional impairment.³ In addition, circumferential burns of the extremities or thorax require consultation with a burn center as they are an indicator of decreased blood flow. Deep circumferential burns of the chest may impair or prevent mechanical ventilation of the burn victim.⁴

In the past, burns were treated with painful debridement of blisters, daily soaking and scrubbing, and frequent bandage changes with topical medications. Today, tissue regeneration and grafting is rapidly becoming the new standard of care in burn injuries.^{3,5,6} The goal of burn treatment is to replace damaged or missing tissue with similar, healthy tissue and restore full function to the involved area with minimal to no scar tissue formation.^{9,13} Second-, third- and fourth-degree

burns most often require a surgical procedure to allow for healing. Although some of the most severe burns may require multisystem surgeries or amputations, most burn injuries are treated with the application of skin grafts.^{3,5,6}

One of the greatest advances in burn treatment has been early excision of necrotic tissue and closure of thermal burn wounds. Early excision and grafting provide a skin substitute for the wound, but further reconstructive surgeries may still be required to restore a normal appearance and function.⁷ According to the International Society for Burn Injury (ISBI), management of major burns by debridement with dressings and subsequent delayed grafting may be the safest approach when resources are limited. By waiting until sloughing of the eschar ~~has happened~~, minimal surgery is needed, with harvest of the skin grafts being the key surgical intervention.¹³

Hydrosurgery describes the use of high-pressure water jets to debride necrotic tissue. This technique ~~aids~~ can aid in dermal skin preservation and scar reduction ~~with~~. In a 137-person within-patient trial, twice as many patients ~~who noted a difference, reporting~~ reported the hydrosurgically debrided area to be better or much better compared to conventional debridement at 12 months, ~~and but~~ one-fourth of patients saw no difference. Although encouraging, additional larger trials are necessary prior to recommendations for routine use.⁹

By performing early excision and grafting, the patient's length of stay in the hospital is significantly reduced, as is the risk for hypertrophic scarring, joint contracture, infection and stiffness. Early closure also allows for quicker rehabilitation and lower mortality rates. As with any surgical procedure, there are also risks and challenges. Major challenges associated with burn surgery include extensive tissue loss and limited availability of tissue, exposure of other structures, scarring and limited tissue pliability.⁷

Skin Grafts

Skin grafting consists of taking tissue from another source and placing it over a wound. Sources include unaffected skin from another location on the burn victim's body, cadaveric skin grafts and amniotic chorion/membrane. The success of a skin graft relies on many factors. The graft bed must be suitable to sustain the graft during the imbibition phase of healing ~~and be free from necrotic or ischemic tissue, inflammation, and infection~~. Also, there must be sufficient perfusion to the graft, either from the graft bed or from another supply such as a flap repair.^{7,8,16} Skin grafts are used for coverage of exposed bone and tendons only if there is a vascularized layer of periosteum or the paratenon is intact.^{7,8}

Coding Implications

This clinical policy references Current Procedural Terminology (CPT®). CPT® is a registered trademark of the American Medical Association. All CPT codes and descriptions are copyrighted 2023/2024, American Medical Association. All rights reserved. CPT codes and CPT descriptions are from the current manuals and those included herein are not intended to be all-inclusive and are included for informational purposes only. Codes referenced in this clinical policy are for informational purposes only and may not support medical necessity. Inclusion or exclusion of any codes does not guarantee coverage. Providers should reference the most up-to-date sources of professional coding guidance prior to the submission of claims for reimbursement of covered services.

NOTE: Coverage is subject to each requested code's inclusion on the corresponding LDH fee schedule. Non-covered codes are denoted (*) and are reviewed for Medical Necessity for members under 21 years of age on a per case basis.

CPT Codes that Support Coverage Medical Necessity Criteria

CPT® Codes	Description
11000	Debridement of extensive eczematous or infected skin; up to 10% of body surface
11001	Debridement of extensive eczematous or infected skin; each additional 10% of the body surface, or part thereof (List separately in addition to code for primary procedure)
11004	Debridement of skin, subcutaneous tissue, muscle and fascia for necrotizing soft tissue infection; external genitalia and perineum
11005	Debridement of skin, subcutaneous tissue, muscle and fascia for necrotizing soft tissue infection; abdominal wall, with or without fascial closure
11006	Debridement of skin, subcutaneous tissue, muscle and fascia for necrotizing soft tissue infection; external genitalia, perineum and abdominal wall, with or without fascial closure
11008	Removal of prosthetic material or mesh, abdominal wall for infection (eg, for chronic or recurrent mesh infection or necrotizing soft tissue infection) (List separately in addition to code for primary procedure)
11010	Debridement including removal of foreign material at the site of an open fracture and/or an open dislocation (eg, excisional debridement); skin and subcutaneous tissues
11011	Debridement including removal of foreign material at the site of an open fracture and/or an open dislocation (eg, excisional debridement); skin, subcutaneous tissue, muscle fascia, and muscle
11012	Debridement including removal of foreign material at the site of an open fracture and/or an open dislocation (eg, excisional debridement); skin, subcutaneous tissue, muscle fascia, muscle, and bone
11042	Debridement, subcutaneous tissue (includes epidermis and dermis, if performed); first 20 sq cm or less
11043	Debridement, muscle and/or fascia (includes epidermis, dermis, and subcutaneous tissue, if performed); first 20 sq cm or less
11044	Debridement, bone (includes epidermis, dermis, subcutaneous tissue, muscle and/or fascia, if performed); first 20 sq cm or less
11045	Debridement, subcutaneous tissue (includes epidermis and dermis, if performed); each additional 20 sq cm, or part thereof (List separately in addition to code for primary procedure)
11046	Debridement, muscle and/or fascia (includes epidermis, dermis, and subcutaneous tissue, if performed); each additional 20 sq cm, or part thereof (List separately in addition to code for primary procedure)

CPT® Codes	Description
11047	Debridement, bone (includes epidermis, dermis, subcutaneous tissue, muscle and/or fascia, if performed); each additional 20 sq cm, or part thereof (List separately in addition to code for primary procedure)
15271	Application of skin substitute graft to trunk, arms, legs, total wound surface area up to 100 sq cm; first 25 sq cm or less wound surface area
15272	Application of skin substitute graft to trunk, arms, legs, total wound surface area up to 100 sq cm; each additional 25 sq cm wound surface area, or part thereof (List separately in addition to code for primary procedure)
15273	Application of skin substitute graft to trunk, arms, legs, total wound surface area greater than or equal to 100 sq cm; first 100 sq cm wound surface area, or 1% of body area of infants and children
15274	Application of skin substitute graft to trunk, arms, legs, total wound surface area greater than or equal to 100 sq cm; each additional 100 sq cm wound surface area, or part thereof, or each additional 1% of body area of infants and children, or part thereof (List separately in addition to code for primary procedure)
15275	Application of skin substitute graft to face, scalp, eyelids, mouth, neck, ears, orbits, genitalia, hands, feet, and/or multiple digits, total wound surface area up to 100 sq cm; first 25 sq cm or less wound surface area
15276	Application of skin substitute graft to face, scalp, eyelids, mouth, neck, ears, orbits, genitalia, hands, feet, and/or multiple digits, total wound surface area up to 100 sq cm; each additional 25 sq cm wound surface area, or part thereof (List separately in addition to code for primary procedure)
15277	Application of skin substitute graft to face, scalp, eyelids, mouth, neck, ears, orbits, genitalia, hands, feet, and/or multiple digits, total wound surface area greater than or equal to 100 sq cm; first 100 sq cm wound surface area, or 1% of body area of infants and children
15278	Application of skin substitute graft to face, scalp, eyelids, mouth, neck, ears, orbits, genitalia, hands, feet, and/or multiple digits, total wound surface area greater than or equal to 100 sq cm; each additional 100 sq cm wound surface area, or part thereof, or each additional 1% of body area of infants and children, or part thereof (List separately in addition to code for primary procedure)

HCPCS Codes that Support ~~Coverage~~Medical Necessity Criteria

HCPCS Codes	Description
Q4100*	Skin substitute, not otherwise specified
Q4101*	Apligraf, per sq cm
Q4104*	Integra bilayer matrix wound dressing (BMWD), per sq cm
Q4105*	Integra dermal regeneration template (DRT) or Integra Omnigraft dermal regeneration matrix, per sq cm
Q4108*	Integra matrix, per sq cm

HCPCs Codes	Description
Q4121*	TheraSkin, per sq cm
Q4182*	Transeyte TransCyte, per sq cm
C9363*	Skin substitute (Integra Meshed Bilayer Wound Matrix), per sq cm

Reviews, Revisions, and Approvals	Revision Date	Approval Date	Effective Date
Converted corporate to local policy.	8/15/20		
Removed ICD-10 codes from policy. References reviewed and updated. Replaced “member” with “member/enrollee.” Annual review. References reviewed and updated. Changed, “review date,” in the header to, “date of last revision,” and, “date,” in the revision log header to, “revision date.” Added “and may not support medical necessity” in coding implications. Removed criteria III. Stating burn surgery was, “not medically necessary when duplicating another provider’s procedure, product, or service.” Reviewed by specialist.	2/22		
Annual review completed. Background updated and minor rewording with no clinical significance. References reviewed, reformatted and updated. Adding additional skin debridement CPT in the same family as 11000 that were missing. All codes are on LDH FS	1/23	4/3/23	
Annual review. Added criteria II.C. that burn must be deep partial-thickness or full-thickness. Added used according to FDA indications to II.D.3. References reviewed and updated. Reviewed by internal specialist.	1/24	3/25/24	
Annual review. References reviewed and updated. Reviewed by external specialist.	2/25	4/28/25	5/29/25
<u>Annual review. Description and background updated with no clinical significance. Updated II.D.3. to clarify HDE abbreviation with no clinical significance. Coding reviewed and updated. Replaced “coverage” to “medical necessity” in the code table titles. References reviewed and updated.</u>	<u>1/26</u>		

References

1. John Hopkins Medicine website. Burns and wounds overview.
http://www.hopkinsmedicine.org/healthlibrary/conditions/adult/physical_medicine_and_rehabilitation/burns_85,p01146. Accessed October 28, 2024, 2025.
2. American Burn Association website. Guidelines for trauma centers caring for burn patients.
<http://ameriburn.org/wp-content/uploads/2017/05/acs-resources-burn-chapter-14.pdf>. Published 2014. Accessed October 28, 2024, 2025.

3. Gauglitz GG, Williams FN. Overview of the management of the severely burned patient. UpToDate. www.uptodate.com. Updated ~~June 15, 2023~~July 25, 2025. Accessed October ~~28, 2024~~28, 2025.
4. Herndon DN, Branski LK. Contemporary methods allowing for safe and convenient use of amniotic membrane as a biologic wound dressing for burns. *Ann Plast Surg*. 2017 Feb;78(2 Suppl 1):S9 to S10. doi: 10.1097/SAP.0000000000000979.
5. Glat PM. The evolution of burn injury management: using dehydrated human amnion/chorion membrane allografts in clinical practice. *Ann Plast Surg*. 2017;78(2 Suppl 1):S1. doi: 10.1097/SAP.0000000000000982
6. Glat PM, Davenport T. Current techniques for burn reconstruction: using dehydrated human amnion/chorion membrane allografts as an adjunctive treatment along the reconstructive ladder. *Ann Plast Surg*. 2017;78(2 Suppl 1):S14 to S18. doi:10.1097/SAP.0000000000000980
7. Liu HF, Zhang F, Lineaweaver WC. History and advancement of burn treatments. *Ann Plast Surg*. 2017;78(2 Suppl 1):S2 to S8. doi:10.1097/SAP.0000000000000896
8. Leon-Villapalos J, Dziewulski P. Overview of surgical procedures used in the management of burn injuries. UpToDate. www.uptodate.com. Updated March 29, 2024. Accessed October ~~28, 2024~~28, 2025.
9. Phelan HA, ~~Bernal~~Bernal E. Treatment of deep ~~burns~~burn injury. UpToDate. www.uptodate.com. Updated March 12, 2024. Accessed October ~~28, 2024~~28, 2025.
10. Shahrokhi S. Skin substitutes. UpToDate. www.uptodate.com. Updated ~~August 2, 2023~~July 31, 2025. Accessed October ~~28, 2024~~28, 2025.
11. Browning JA, Cindass R. Burn. Debridement, grafting, and reconstruction. In: *StatPearls*. Treasure Island (FL): StatPearls Publishing; June 5, 2022.
12. Vyas KS, Vasconez HC. Wound healing: biologics, skin substitutes, biomembranes and scaffolds. *Healthcare (Basel)*. 2014;2(3):356 to 400. Published 2014 Sep 10. doi:10.3390/healthcare2030356
13. International Society for Burn Injury (ISBI). ISBI practice guidelines committee. ISBI practice guidelines for burn care. *Burns*. 2016;42(5):953 to 1021. <http://dx.doi.org/10.1016/j.burns.2016.05.013> doi: 10.1016/j.burns.2016.05.013.
14. Vig K, Chaudhari A, Tripathi S, et al. Advances in Skin Regeneration Using Tissue Engineering. *Int J Mol Sci*. 2017;18(4):789. Published 2017 Apr 7. doi:10.3390/ijms18040789
15. Orgill DP. Assessment and classification of burn injury. UpToDate. www.uptodate.com. Updated December 10, 2024. Accessed October 6, 2025.
16. Leon-Villapalos J, Dziewulski P. Skin autografting. UpToDate. www.uptodate.com. Updated October 28, 2024. Accessed October 6, 2025.
17. ISBI Practice Guidelines Committee; Advisory Subcommittee; Steering Subcommittee. ISBI Practice Guidelines for Burn Care, Part 2. *Burns*. 2018 Nov;44(7):1617 to 1706. doi: 10.1016/j.burns.2018.09.012.

Important Reminder

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program approval status; evidence-based guidelines and positions of leading national health professional

organizations; views of physicians practicing in relevant clinical areas affected by this clinical policy; and other available clinical information. LHCC makes no representations and accepts no liability with respect to the content of any external information used or relied upon in developing this clinical policy. This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved.

The purpose of this clinical policy is to provide a guide to medical necessity, which is a component of the guidelines used to assist in making coverage decisions and administering benefits. It does not constitute a contract or guarantee regarding payment or results. Coverage decisions and the administration of benefits are subject to all terms, conditions, exclusions and limitations of the coverage documents (e.g., evidence of coverage, certificate of coverage, policy, contract of insurance, etc.), as well as to state and federal requirements and applicable LHCC administrative policies and procedures.

This clinical policy is effective as of the date determined by LHCC. The date of posting may not be the effective date of this clinical policy. This clinical policy may be subject to applicable legal and regulatory requirements relating to provider notification. If there is a discrepancy between the effective date of this clinical policy and any applicable legal or regulatory requirement, the requirements of law and regulation shall govern. LHCC retains the right to change, amend or withdraw this clinical policy, and additional clinical policies may be developed and adopted as needed, at any time.

This clinical policy does not constitute medical advice, medical treatment or medical care. It is not intended to dictate to providers how to practice medicine. Providers are expected to exercise professional medical judgment in providing the most appropriate care, and are solely responsible for the medical advice and treatment of members/enrollees. This clinical policy is not intended to recommend treatment for members/enrollees. Members/enrollees should consult with their treating physician in connection with diagnosis and treatment decisions.

Providers referred to in this clinical policy are independent contractors who exercise independent judgment and over whom LHCC has no control or right of control. Providers are not agents or employees of LHCC.

This clinical policy is the property of LHCC. Unauthorized copying, use, and distribution of this clinical policy or any information contained herein are strictly prohibited. Providers, members/enrollees and their representatives are bound to the terms and conditions expressed herein through the terms of their contracts. Where no such contract exists, providers, members/enrollees and their representatives agree to be bound by such terms and conditions by providing services to members/enrollees and/or submitting claims for payment for such services.

©2024 Louisiana Healthcare Connections. All rights reserved. All materials are exclusively owned by Louisiana Healthcare Connections and are protected by United States copyright law and international copyright law. No part of this publication may be reproduced, copied, modified, distributed, displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise published without the prior written permission of Louisiana Healthcare Connections. You may not alter or remove any trademark, copyright or other notice contained

herein. Louisiana Healthcare Connections is a registered trademarks exclusively owned by Louisiana Healthcare Connections.