

UnitedHealthcare, Inc. ("UHC") Proprietary and Confidential Information: The information contained in this document is confidential, proprietary and the sole property of UHC. The recipient of this information agrees not to disclose or use it for any purpose other than to facilitate UHC's compliance with applicable State Medicaid contractual requirements. Any other use or disclosure is strictly prohibited and requires the express written consent of UHC.



Chemotherapy Observation or Inpatient Hospitalization (for Louisiana Only)

Guideline Number: CS198LA.A

Effective Date: TBD

[➔ Instructions for Use](#)

<u>Table of Contents</u>	<u>Page</u>
Application.....	1
Coverage Rationale.....	1
References.....	4
Guideline History/Revision Information.....	4
Instructions for Use.....	4

Application

This Medical Policy only applies to the state of Louisiana.

Coverage Rationale

[➔ See Benefit Considerations](#)

Most cancer chemotherapies can be administered safely and effectively in a physician office or through home healthcare services. However, because of the risk of certain toxicities or individual comorbidities, some cancer chemotherapy may be administered either in a facility observation or inpatient unit.

This guideline does not apply to individuals under 18 years of age.

An inpatient stay is medically necessary for drug regimens that require inpatient monitoring or complex administration over multiple days:

<u>Regimen</u>	<u>Drugs</u>	<u>Cancer Type</u>	<u>Factors contributing to the need for inpatient stay:</u>
<u>EPOCH or DA-EPOCH or R-EPOCH</u>	<ul style="list-style-type: none"> <u>Etoposide 50 mg/m²/day continuous infusion on days 1 to 4</u> <u>Prednisone</u> <u>Vincristine (Oncovine) IV days 1-4</u> <u>Cyclophosphamide 750 mg/m² IV on day 5</u> 	<u>Lymphoma</u>	<u>Coordination of multiple infusions or multiple drugs over 96 hours</u>

UnitedHealthcare, Inc. ("UHC") Proprietary and Confidential Information: The information contained in this document is confidential, proprietary and the sole property of UHC. The recipient of this information agrees not to disclose or use it for any purpose other than to facilitate UHC's compliance with applicable State Medicaid contractual requirements. Any other use or disclosure is strictly prohibited and requires the express written consent of UHC.

<u>Regimen</u>	<u>Drugs</u>	<u>Cancer Type</u>	<u>Factors contributing to the need for inpatient stay:</u>
	<ul style="list-style-type: none"> <u>Doxorubicin (Hydroxydaunorubicin) 10 mg/m²/day continuous infusion on days 1-4</u> <u>With or without Rituximab</u> 		
<u>ESHAP or R-ESHAP</u>	<ul style="list-style-type: none"> <u>Etoposide 40 mg/m²/day continuous infusion on days 1 to 4</u> <u>Methylprednisolone (solumedrol)</u> <u>Cytarabine ("High-dose Ara-c") 2g/m²</u> <u>Cisplatin (platinol) 25mg/m continuous infusion days 1-4</u> <u>With or without Rituximab</u> 	<u>Lymphoma</u>	<ul style="list-style-type: none"> <u>Coordination of multiple infusions or multiple drugs over 96 hours</u> <u>Monitor for CNS toxicity with cytarabine</u>
<u>Interleukin 2 infusion</u>	<u>Interleukin 2 600,000 IU/kg IV every 8 hours for up to 14 consecutive doses over 5 days</u>	<u>Melanoma</u> <u>Renal Cell Cancer</u>	<ul style="list-style-type: none"> <u>Continuous cardiac monitoring</u> <u>Close monitoring of serum electrolytes, creatinine, bilirubin, urine output</u> <u>Vasopressor support with dopamine</u> <u>Proximity to intensive care unit</u>
<u>High dose Ifosphamide</u>	<u>Ifosphamide infusion > 1g/m²/day</u>	<u>Sarcoma</u>	<ul style="list-style-type: none"> <u>Close monitoring of serum electrolytes and urine pH</u> <u>Replacement of electrolytes</u> <u>Alkalinization of urine</u>
<u>High dose methotrexate with leucovorin rescue</u>	<ul style="list-style-type: none"> <u>Methotrexate dose at > 500 mg/m²</u> <u>Leucovorin 15 mg every 6 hours for eight doses beginning 12 hours after the completion of methotrexate infusion, and increased to 50 mg IV every 6 hours if methotrexate levels are >20 µmol/L at 0 hour, are >1.0 µmol/L at 24 hours, or are >0.1 µmol/L at 48 hours after the end of methotrexate infusion, until levels are <0.1 µmol/L plus</u> 	<u>Lymphoma</u> <u>Sarcoma</u>	<u>Close monitoring of serum methotrexate levels</u>
<u>Hyper-CVAD</u>	<ul style="list-style-type: none"> <u>Cycles 1, 3, 5, and 7 (3-4 weeks between cycles):</u> <u>Cyclophosphamide 300 mg/m² IV over 2 hours every 12 hours for 6 doses</u> 	<u>Lymphoma</u> <u>Leukemia</u>	<ul style="list-style-type: none"> <u>Coordination of multiple infusions or multiple drugs over 96 hours</u>

UnitedHealthcare, Inc. ("UHC") Proprietary and Confidential Information: The information contained in this document is confidential, proprietary and the sole property of UHC. The recipient of this information agrees not to disclose or use it for any purpose other than to facilitate UHC's compliance with applicable State Medicaid contractual requirements. Any other use or disclosure is strictly prohibited and requires the express written consent of UHC.

<u>Regimen</u>	<u>Drugs</u>	<u>Cancer Type</u>	<u>Factors contributing to the need for inpatient stay:</u>
	<ul style="list-style-type: none"> • <u>Mesna 600 mg/m²/day continuous infusion on days 1-3, starting 1 hour before cyclophosphamide</u> • <u>Vincristine</u> • <u>Doxorubicin 50 mg/m² IV on day 4</u> • <u>Dexamethasone</u> • <u>Cycles 2, 4, 6, and 8 (3-4 weeks between cycles):</u> • <u>Methotrexate 200 mg/m² IV over 2 hours followed by 800 mg/m² IV over 22 hours on day 1 plus</u> • <u>Cytarabine 3 g/m² (1 g/m² for patients older than 60 years) IV over 2 hours every 12 hours for four doses starting on day 2</u> • <u>Leucovorin 15 mg every 6 hours for eight doses beginning 12 hours after the completion of methotrexate infusion, and increased to 50 mg IV every 6 hours if methotrexate levels are >20 µmol/L at 0 hour, are >1.0 µmol/L at 24 hours, or are >0.1 µmol/L at 48 hours after the end of methotrexate infusion, until levels are <0.1 µmol/L plus</u> • <u>Methylprednisolone 50 mg</u> 		<ul style="list-style-type: none"> • <u>Bladder irrigation with cyclophosphamide</u> • <u>Close monitoring of serum methotrexate levels</u>

The following are clinical conditions or complications of cancer chemotherapy which, when present, may require an observation stay:

- Known hypersensitivity reactions from previous infusion
- Congestive heart failure or chronic renal failure requiring high volume fluid infusions
- Transcatheter arterial chemoembolization (TACE) or intra-arterial chemotherapy infusion
- Comorbidities that require an observation or overnight stay
- Cancer chemotherapy administered during a hospitalization for an unrelated problem

The following are clinical conditions which require an inpatient hospital stay:

- Acute leukemia
- Intra-arterial infusion of chemotherapy
- Prophylaxis of tumor lysis syndrome in cases of high grade lymphoma with large masses
- Comorbidities that require an inpatient stay

Conditions requiring observation or inpatient hospital treatment other than those noted above will be reviewed on a case-by-case basis.

UnitedHealthcare, Inc. ("UHC") Proprietary and Confidential Information: The information contained in this document is confidential, proprietary and the sole property of UHC. The recipient of this information agrees not to disclose or use it for any purpose other than to facilitate UHC's compliance with applicable State Medicaid contractual requirements. Any other use or disclosure is strictly prohibited and requires the express written consent of UHC.

For medical necessity clinical coverage criteria, see InterQual® 2020, Apr. 2020 Release, LOC:Acute Adult Hematology/Oncology: Chemotherapy.

Click [here](#) to view the InterQual® criteria.

Additional Review Points

- A written protocol will be expected to be followed by the provider administering the chemotherapy drug.
- Any requests for an extension of the inpatient stay beyond the recommended day(s) must be clinically reviewed.

References

Chemocare (Cleveland Clinic Foundation). <http://www.chemocare.com/bio/interleukin.asp>. Accessed January 22, 2020.

Drugs at FDA. <http://www.accessdata.fda.gov/scripts/cder/daf/>. Accessed January 22, 2020.

Medicare Benefit Policy Manual, Chapter 6 – Hospital Services Covered under Part B, section 20.6 – Outpatient Observation Services, A. Outpatient Observation Services Defined. <http://www.cms.gov/Regulations-and-Guidance/Guidance/Manuals/downloads/bp102c06.pdf>. Accessed January 22, 2020.

Guideline History/Revision Information

<u>Date</u>	<u>Summary of Changes</u>
<u>TBD</u>	• <u>New Policy</u>

Instructions for Use

This Utilization Review Guideline provides assistance in interpreting UnitedHealthcare standard benefit plans. When deciding coverage, the federal, state or contractual requirements for benefit plan coverage must be referenced as the terms of the federal, state or contractual requirements for benefit plan coverage may differ from the standard benefit plan. In the event of a conflict, the federal, state or contractual requirements for benefit plan coverage govern. Before using this guideline, please check the federal, state or contractual requirements for benefit plan coverage. UnitedHealthcare reserves the right to modify its Policies and Guidelines as necessary. This Utilization Review Guideline is provided for informational purposes. It does not constitute medical advice.

UnitedHealthcare may also use tools developed by third parties, such as the InterQual® criteria, to assist us in administering health benefits. The UnitedHealthcare Utilization Review Guidelines are intended to be used in connection with the independent professional medical judgment of a qualified health care provider and do not constitute the practice of medicine or medical advice.