

Clinical Policy: Hyperhidrosis Treatments

Reference Number: LA.CP.MP.62

Date of Last Revision: ~~42~~/23~~2~~

Coding Implications

Revision Log

See [Important Reminder](#) at the end of this policy for important regulatory and legal information.

Description

Hyperhidrosis is defined as excessive sweating beyond a level required to maintain normal body temperature in response to heat exposure or exercise.

[Refer to the Louisiana Medicaid Preferred Drugs List, \(PDL\) and the LDH guidelines for coverage criteria for the medications referenced in this clinical policy](#)

- [AbobotulinumtoxinA \(Dysport®\)](#)
- [OnabotulinumtoxinA \(Botox®\)](#)
- [Qbrexza \(glycopyrronium\)](#)

Policy/Criteria

- I. It is the policy of Louisiana Healthcare Connections that treatment with iontophoresis (electrophoresis, Drionic device) is **medically necessary** when *all* of the following criteria are met:
 - A. Diagnosis of primary hyperhidrosis;
 - B. Development of medical complications, such as skin maceration with secondary skin infections; *or* has a significant constant disruption of professional and/or social life (e.g., recurrent changing of clothes, affecting job/social function, etc.) which has occurred because of excessive sweating;
 - C. Unresponsive or unable to tolerate at least one of the pharmacotherapies prescribed for excessive sweating (e.g., anticholinergics, beta-blockers, or benzodiazepines);
 - D. Failed a ~~six~~6-month trial of conservative management including the adherent application of aluminum chloride hexahydrate [Drysol by prescription] or topical agents have resulted in a severe rash;
 - E. Has none of the following contraindications:
 1. Cardiac pacemaker;
 2. Cardiac arrhythmias;
 3. Pregnancy (hyperhidrosis often improves during pregnancy);
 4. Metal implants, depending on size and position (may divert the electric current);
 5. Cracked skin near the treatment area.
- II. It is the policy of Louisiana Healthcare Connections that endoscopic thoracic sympathectomy (ETS) for palmar or palmar and axillary hyperhidrosis is **medically necessary** when *all* of the following criteria are met:
 - A. Meets all of the iontophoresis criteria in I.A-D;
 - B. Has a resting heart rate \geq 55 beats per minute;
 - C. Hyperhidrosis symptoms started at an early age (usually < 16 years), and surgery is requested for a young member/enrollee (usually <25 years of age);

- D. Body mass index <28;
- E. Reports no sweating during sleep;
- F. The member/enrollee ~~has~~ **relatively healthy with** no significant comorbidities;
- G. Has persistent and severe primary hyperhidrosis;
- H. Has failed one of the following:
 - 1. Iontophoresis;
 - 2. Trial of botulinum toxin for predominantly axillary hyperhidrosis.

III. It is the policy of Louisiana Healthcare Connections that surgical excision of axillary sweat glands for axillary hyperhidrosis are **medically necessary** when *all* of the following criteria are met:

- A. Meets all of the iontophoresis criteria in I.A ~~through~~ -D;
- B. Has persistent and severe primary hyperhidrosis;
- C. Has failed one of the following:
 - 1. Iontophoresis;
 - 2. Trial of botulinum toxin.

Note: The normal line of medical therapy is:

- 1. Drysol, then ~~B~~ **bot**ox or topical glycopyrronium for axillary hyperhidrosis
- 2. Drysol, then iontophoresis for palmo-plantar hyperhidrosis
- 3. Other treatments are third-line therapies (iontophoresis and surgery for axillary hyperhidrosis, and Botox and surgery for palmo-plantar hyperhidrosis).

IV. There is insufficient evidence in published peer-reviewed literature to support all other treatments for hyperhidrosis, including, but not limited to, microwave therapy, or liposuction as the sole method of removing axillary sweat glands.

Background

Hyperhidrosis can be classified as either primary or secondary.¹² ~~12~~ Primary focal hyperhidrosis is idiopathic in nature and is defined as excessive sweating induced by sympathetic hyperactivity in selected areas that is not associated with an underlying disease process.³³ The most common locations are underarms (axillary hyperhidrosis), hands (palmar hyperhidrosis), and feet (plantar hyperhidrosis). Primary focal hyperhidrosis is a condition that is characterized by visible, excessive sweating of at least six months' duration without apparent cause. Hyperhidrosis can ruin clothing, produce emotional distress, and lead to occupational disability.¹² ~~12~~

Secondary hyperhidrosis can result from a variety of drugs, such as tricyclic antidepressants, selective serotonin reuptake inhibitors (SSRIs), or underlying diseases/conditions, such as febrile diseases, diabetes mellitus, or menopause. Secondary hyperhidrosis is usually generalized or craniofacial sweating. Secondary gustatory hyperhidrosis is excessive sweating on ingesting highly spiced foods. This trigeminovascular reflex typically occurs symmetrically on scalp or face and predominately over forehead, lips, and nose. Secondary facial gustatory sweating, in contrast, is usually asymmetrical and occurs independently of the nature of the ingested food. This phenomenon frequently occurs after injury or surgery in the region of the parotid gland.

A variety of therapies have been investigated for primary hyperhidrosis, including topical therapy with aluminum chloride, iontophoresis, intradermal injections of botulinum toxin type A,

endoscopic transthoracic sympathectomy, and surgical excision of axillary sweat glands. ^{1,2,12}
~~1,2,12~~ Endoscopic thoracic sympathectomy (ETS) is an invasive procedure intended to arrest the symptoms of hyperhidrosis and involves interrupting the upper thoracic sympathetic chain through clipping, cauterization, or cutting. ¹²~~12~~ Treatment of secondary hyperhidrosis focuses on the treatment of the underlying cause, such as discontinuing certain drugs or hormone replacement therapy as a treatment of menopausal symptoms.

Microwave energy has been proposed for the treatment of primary axillary hyperhidrosis. The miraDry System (Mirimar Labs, Inc) is a Food and Drug Administration (FDA) approved device indicated for treatment of primary axillary hyperhidrosis. It is not indicated for treating hyperhidrosis related to other body areas or generalized hyperhidrosis. Evidence is still emerging in the published peer-reviewed literature to support the safety and efficacy of microwave energy for the treatment of primary axillary hyperhidrosis. ¹⁷~~17~~ Hyperhidrosis can be classified as either primary or secondary. Primary focal hyperhidrosis is idiopathic in nature and is defined as excessive sweating induced by sympathetic hyperactivity in selected areas that is not associated with an underlying disease process. The most common locations are underarms (axillary hyperhidrosis), hands (palmar hyperhidrosis), and feet (plantar hyperhidrosis). Primary focal hyperhidrosis is a condition that is characterized by visible, excessive sweating of at least 6 months' duration without apparent cause. Hyperhidrosis can ruin clothing, produce emotional distress, and lead to occupational disability.

Secondary hyperhidrosis can result from a variety of drugs, such as tricyclic antidepressants, selective serotonin reuptake inhibitors (SSRIs), or underlying diseases/conditions, such as febrile diseases, diabetes mellitus, or menopause. Secondary hyperhidrosis is usually generalized or craniofacial sweating. Secondary gustatory hyperhidrosis is excessive sweating on ingesting highly spiced foods. This trigeminovascular reflex typically occurs symmetrically on scalp or face and predominately over forehead, lips, and nose. Secondary facial gustatory sweating, in contrast, is usually asymmetrical and occurs independently of the nature of the ingested food. This phenomenon frequently occurs after injury or surgery in the region of the parotid gland.

A variety of therapies have been investigated for primary hyperhidrosis, including topical therapy with aluminum chloride, iontophoresis, intradermal injections of botulinum toxin type A, endoscopic transthoracic sympathectomy, and surgical excision of axillary sweat glands. Thoracic sympathectomy is an invasive procedure intended to arrest the symptoms of hyperhidrosis. Treatment of secondary hyperhidrosis focuses on the treatment of the underlying cause, such as discontinuing certain drugs or hormone replacement therapy as a treatment of menopausal symptoms.

Microwave energy has been proposed for the treatment of primary axillary hyperhidrosis. The miraDry System (Mirimar Labs, Inc) is an FDA approved device indicated for treatment of primary axillary hyperhidrosis. It is not indicated for treating hyperhidrosis related to other body areas or generalized hyperhidrosis. There is insufficient evidence in published peer-reviewed literature to support the safety and efficacy of microwave energy for the treatment of primary axillary hyperhidrosis. Most of the studies are limited by small sample size with data on long-term health outcomes lacking.

Coding Implications

This clinical policy references Current Procedural Terminology (CPT®). CPT® is a registered trademark of the American Medical Association. All CPT codes and descriptions are copyrighted 2021, American Medical Association. All rights reserved. CPT codes and CPT descriptions are from the current manuals and those included herein are not intended to be all-inclusive and are included for informational purposes only. Codes referenced in this clinical policy are for informational purposes only and may not support medical necessity. Inclusion or exclusion of any codes does not guarantee coverage. Providers should reference the most up-to-date sources of professional coding guidance prior to the submission of claims for reimbursement of covered services.

~~This clinical policy references Current Procedural Terminology (CPT®). CPT® is a registered trademark of the American Medical Association. All CPT codes and descriptions are copyrighted 2020, American Medical Association. All rights reserved. CPT codes and CPT descriptions are from the current manuals and those included herein are not intended to be all-inclusive and are included for informational purposes only. Codes referenced in this clinical policy are for informational purposes only and may not support medical necessity. Inclusion or exclusion of any codes does not guarantee coverage. Providers should reference the most up-to-date sources of professional coding guidance prior to the submission of claims for reimbursement of covered services.~~

CPT® Codes	Description
11450	Excision of skin and subcutaneous tissue for hidradenitis, axillary; with simple or intermediate repair
11451	Excision of skin and subcutaneous tissue for hidradenitis, axillary; with complex repair
15877*	Suction assisted lipectomy; trunk
15878*	Suction assisted lipectomy; upper extremity
32664	Thoracoscopy, surgical; with thoracic sympathectomy
64802 <u>through–</u> 64823	Sympathectomy, sympathetic nerves
97024*	Application of a modality to 1 or more areas; diathermy (eg, microwave)
97033	Application of a modality to 1 or more areas; iontophoresis, each 15 minutes

* Insufficient evidence in published peer-reviewed literature to support suction assisted liposuction as the sole method of removing axillary sweat glands.

* Considered investigational, not medically necessary when used to report liposuction as the sole method of removing axillary sweat glands.

ICD-10-CM Diagnosis Codes that Support Coverage Criteria

ICD-10-CM Code	Description
L74.510-L74.519	Primary focal hyperhidrosis
L74.52	Secondary hyperhidrosis
R61	Generalized hyperhidrosis

* Insufficient evidence in published peer-reviewed literature to support suction assisted liposuction as the sole method of removing axillary sweat glands.

Reviews, Revisions, and Approvals	Revision Date	Approval Date
Converted corporate to local policy.	1/21/2020	
Annual review. References reviewed and updated. Reviewed by specialist. Changed "Last Review Date" in the header to "Date of Last Revision" and "Date" in revision log to "Revision Date". Added "and may not support medical necessity" to coding implications. "Experimental/investigational" verbiage replaced in policy statement and background with descriptive language.	2/22	
<u>Annual review. Updated Criteria II.B. to greater than 55 beats per minute. Removed "is relatively healthy" in criteria II.F. Background updated with no impact on criteria. ICD-10 codes removed. References reviewed and updated. References made to include LDH coverage criteria for medications described in the policy</u>	<u>42/23</u>	

References

1. Cerfolio RJ, De Campos JR, Bryant AS, et al. The Society of Thoracic Surgeons expert consensus for the surgical treatment of hyperhidrosis. *Ann Thorac Surg*. 2011;91(5):1642 to 1648 doi:10.1016/j.athoracsur.2011.01.105
2. Eisenach JH, Atkinson JL, Fealey RD. Hyperhidrosis: evolving therapies for a well-established phenomenon. [published correction appears in *Mayo Clin Proc*. 2005 Jun;80(6):828]. *Mayo Clin Proc*. 2005;80(5):657 to 666. doi:10.4065/80.5.657
3. Glaser DA. The use of botulinum toxins to treat hyperhidrosis and gustatory sweating syndrome. *Neurotox Res*. 2006;9(2 to 3):173 to 177. doi:10.1007/BF03033936
4. Glaser DA, Coleman WP 3rd, Fan LK, et al. A randomized, blinded clinical evaluation of a novel microwave device for treating axillary hyperhidrosis: the dermatologic reduction in underarm perspiration study. *Dermatol Surg*. 2012;38(2):185 to -191. doi:10.1111/j.1524-4725.2011.02250.x
5. Hong HC, Lupin M, O'Shaughnessy KF. Clinical evaluation of a microwave device for treating axillary hyperhidrosis. *Dermatol Surg*. 2012;38(5):728 to 735. doi:10.1111/j.1524-4725.2012.02375.x
6. Hsu TH, Chen YT, Tu YK, Li CN. A systematic review of microwave-based therapy for axillary hyperhidrosis. *J Cosmet Laser Ther*. 2017;19(5):275 to 282. doi:10.1080/14764172.2017.1303168
7. Hyperhidrosis Treatment Overview. International Hyperhidrosis Society. <https://www.sweathelp.org/hyperhidrosis-treatments/treatment-overview.html>. Accessed December 19, 2022.
8. Karpinski RHS. Surgical treatment of axillary hyperhidrosis treatment & management. Medscape. <https://emedicine.medscape.com/article/1296530-treatment>. Updated February 12, 2019. Accessed December 14, 2022.
9. Lakraj, AA, Moghimi N, Jabbari B. Hyperhidrosis: anatomy, pathophysiology and treatment with emphasis on the role of botulinum toxins. *Toxins (Basel)*. 2013;5(4):821 to 840. Published 2013 Apr 23. doi:10.3390/toxins5040821

10. Oakley A. Hyperhidrosis. DermNet NZ. <https://dermnetnz.org/topics/hyperhidrosis>. Accessed December 14, 2022.
11. Cole A, Oakley A. DermNet NZ. <https://dermnetnz.org/topics/iontophoresis>. Updated April 2015. Accessed December 14, 2022.
12. Smith CC, Pariser, D. Primary focal hyperhidrosis. UpToDate. www.uptodate.com. Published December 01, 2022. Accessed December 12, 2022.
13. Pariser DM, Hebert AA, Drew J, Quiring J, Gopalan R, Glaser DA. Topical Glycopyrronium Tosylate for the Treatment of Primary Axillary Hyperhidrosis: Patient-Reported Outcomes from the ATMOS-1 and ATMOS-2 Phase III Randomized Controlled Trials. *Am J Clin Dermatol*. 2019;20(1):135 to 145. doi:10.1007/s40257-018-0395-0
14. Glaser DA, Hebert AA, Nast A, et al. Topical glycopyrronium tosylate for the treatment of primary axillary hyperhidrosis: Results from the ATMOS-1 and ATMOS-2 phase 3 randomized controlled trials. *J Am Acad Dermatol*. 2019;80(1):128 to 138.e2. doi:10.1016/j.jaad.2018.07.002
15. Sheikh NK, Dua A. Iontophoresis Analgesic Medications. In: *StatPearls*. Treasure Island (FL): StatPearls Publishing; August 1, 2022. Accessed December 09, 2022.
16. Vannucci F, Araújo JA. Thoracic sympathectomy for hyperhidrosis: from surgical indications to clinical results. *J Thorac Dis*. 2017;9(Suppl 3):S178 to S192. doi:10.21037/jtd.2017.04.04
17. Evidence Analysis Research Brief. Microwave therapy for management of hyperhidrosis. Hayes. www.hayesinc.com. Published April 19, 2022. Accessed December 12, 2022.
1. Cerfolio RJ, De Campos JR, Bryant AS, et al. The Society of Thoracic Surgeons expert consensus for the surgical treatment of hyperhidrosis. *Ann Thorac Surg*. 2011;91(5):1642–1648 doi:10.1016/j.athoracsur.2011.01.105
2. Eisenach JH, Atkinson JL, Fealey RD. Hyperhidrosis: Evolving therapies for a well-established phenomenon. [published correction appears in *Mayo Clin Proc*. 2005 Jun;80(6):828]. *Mayo Clin Proc*. 2005;80(5):657–666. doi:10.4065/80.5.657
3. Glaser DA. The use of botulinum toxins to treat hyperhidrosis and gustatory sweating syndrome. *Neurotox Res*. 2006;9(2–3):173–177. doi:10.1007/BF03033936
4. Glaser DA, Coleman WP 3rd, Fan LK, et al. A randomized, blinded clinical evaluation of a novel microwave device for treating axillary hyperhidrosis: the dermatologic reduction in underarm perspiration study. *Dermatol Surg*. 2012;38(2):185–191. doi:10.1111/j.1524-4725.2011.02250.x
5. Hong HC, Lupin M, O'Shaughnessy KF. Clinical evaluation of a microwave device for treating axillary hyperhidrosis. *Dermatol Surg*. 2012;38(5):728–735. doi:10.1111/j.1524-4725.2012.02375.x
6. Hsu TH, Chen YT, Tu YK, Li CN. A systematic review of microwave-based therapy for axillary hyperhidrosis. *J Cosmet Laser Ther*. 2017;19(5):275–282. doi:10.1080/14764172.2017.1303168
7. International Hyperhidrosis Society. Hyperhidrosis Treatment Overview. <https://www.sweathelp.org/hyperhidrosis-treatments/treatment-overview.html>. Published 2017. Accessed January 4, 2020.
8. Karpinski RHS. Surgical treatment of axillary hyperhidrosis treatment & management. Medscape. <https://emedicine.medscape.com/article/1296530-treatment> Updated February 12, 2019. Accessed December 16, 2021.

9. Lakraj, A.D; Moghimi, N; Jabbari, B. Hyperhidrosis: Anatomy, Pathophysiology and Treatment with Emphasis on the Role of Botulinum Toxins. *Toxins (Basel)*. 2013;5(4):821–840. Published 2013 Apr 23. doi:10.3390/toxins5040821
10. Oakley A. Hyperhidrosis. DermNet NZ. <https://dermnetnz.org/topics/hyperhidrosis>. Updated July 2015. Accessed January 4, 2022.
11. Cole A, Oakley A. DermNet NZ. <https://dermnetnz.org/topics/iontophoresis>. Updated April 2015. Accessed December 16, 2021.
12. Smith CC, Pariser, D. Primary focal hyperhidrosis. UpToDate. www.uptodate.com. Published January 2, 2020. Accessed December 16, 2021.
13. Pariser DM, Hebert AA, Drew J, Quiring J, Gopalan R, Glaser DA. Topical Glycopyrronium Tosylate for the Treatment of Primary Axillary Hyperhidrosis: Patient Reported Outcomes from the ATMOS-1 and ATMOS-2 Phase III Randomized Controlled Trials. *Am J Clin Dermatol*. 2019;20(1):135–145. doi:10.1007/s40257-018-0395-0
14. Glaser DA, Hebert AA, Nast A, et al. Topical glycopyrronium tosylate for the treatment of primary axillary hyperhidrosis: Results from the ATMOS-1 and ATMOS-2 phase 3 randomized controlled trials. *J Am Acad Dermatol*. 2019;80(1):128–138.e2. doi:10.1016/j.jaad.2018.07.002
15. Sheikh NK, Dua A. Iontophoresis Analgesic Medications. In: StatPearls. Treasure Island (FL): StatPearls Publishing; August 3, 2021. Accessed December 16, 2021.
16. Vannucci F, Araújo JA. Thoracic sympathectomy for hyperhidrosis: from surgical indications to clinical results. *J Thorac Dis*. 2017;9(Suppl 3):S178–S192. doi:10.21037/jtd.2017.04.04

Important Reminder

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program approval status; evidence-based guidelines and positions of leading national health professional organizations; views of physicians practicing in relevant clinical areas affected by this clinical policy; and other available clinical information. LHCC makes no representations and accepts no liability with respect to the content of any external information used or relied upon in developing this clinical policy. This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved.

The purpose of this clinical policy is to provide a guide to medical necessity, which is a component of the guidelines used to assist in making coverage decisions and administering benefits. It does not constitute a contract or guarantee regarding payment or results. Coverage decisions and the administration of benefits are subject to all terms, conditions, exclusions and limitations of the coverage documents (e.g., evidence of coverage, certificate of coverage, policy, contract of insurance, etc.), as well as to state and federal requirements and applicable LHCC administrative policies and procedures.

This clinical policy is effective as of the date determined by LHCC. The date of posting may not be the effective date of this clinical policy. This clinical policy may be subject to applicable legal and regulatory requirements relating to provider notification. If there is a discrepancy between the effective date of this clinical policy and any applicable legal or regulatory requirement, the requirements of law and regulation shall govern. LHCC retains the right to change, amend or

withdraw this clinical policy, and additional clinical policies may be developed and adopted as needed, at any time.

This clinical policy does not constitute medical advice, medical treatment or medical care. It is not intended to dictate to providers how to practice medicine. Providers are expected to exercise professional medical judgment in providing the most appropriate care, and are solely responsible for the medical advice and treatment of members/enrollees. This clinical policy is not intended to recommend treatment for members/enrollees. Members/enrollees should consult with their treating physician in connection with diagnosis and treatment decisions.

Providers referred to in this clinical policy are independent contractors who exercise independent judgment and over whom LHCC has no control or right of control. Providers are not agents or employees of LHCC.

This clinical policy is the property of LHCC. Unauthorized copying, use, and distribution of this clinical policy or any information contained herein are strictly prohibited. Providers, members/enrollees and their representatives are bound to the terms and conditions expressed herein through the terms of their contracts. Where no such contract exists, providers, members/enrollees and their representatives agree to be bound by such terms and conditions by providing services to members/enrollees and/or submitting claims for payment for such services.

| ©2023⁹ Louisiana Healthcare Connections. All rights reserved. All materials are exclusively owned by Louisiana Healthcare Connections and are protected by United States copyright law and international copyright law. No part of this publication may be reproduced, copied, modified, distributed, displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise published without the prior written permission of Louisiana Healthcare Connections. You may not alter or remove any trademark, copyright or other notice contained herein. Louisiana Healthcare Connections is a registered trademark exclusively owned by Louisiana Healthcare Connections.