



BENLYSTA® (BELIMUMAB)

Policy Number: CSLA2020D0046K

Effective Date: TBD

Table of Contents	Page
APPLICATION	1
COVERAGE RATIONALE	1
APPLICABLE CODES	2
BACKGROUND	2
CLINICAL EVIDENCE	2
U.S. FOOD AND DRUG ADMINISTRATION	54
CENTERS FOR MEDICARE AND MEDICAID SERVICES	65
REFERENCES	65
POLICY HISTORY/REVISION INFORMATION	76
INSTRUCTIONS FOR USE	7

Commercial Policy

- [Benlysta® \(Belimumab\)](#)

APPLICATION

This Medical Benefit Drug Policy only applies to state of Louisiana.

COVERAGE RATIONALE

This policy refers only to Benlysta (belimumab) injection for intravenous infusion for the treatment of systemic lupus erythematosus (SLE). Benlysta (belimumab) for self-administered subcutaneous injection is obtained under the pharmacy benefit and is indicated for systemic lupus erythematosus.

Benlysta (belimumab) is proven and medically necessary for the treatment of active systemic lupus erythematosus when ALL of the following criteria are met:

- For **initial therapy**, all of the following:
 - Diagnosis of active systemic lupus erythematosus; without severe active lupus nephritis or severe active central nervous system lupus; **and**
 - Laboratory testing has documented the presence of autoantibodies [e.g., ANA, Anti-dsDNA, Anti-Sm, Anti-Ro/SSA, Anti-La/SSB]; **and**
 - Currently receiving at least one standard of care treatment for active systemic lupus erythematosus (e.g., antimalarials, corticosteroids, or immunosuppressants); that is not a biologic or intravenous cyclophosphamide;^{1-7,10} **and**
 - Benlysta is initiated and titrated according to U.S. Food and Drug Administration labeled dosing for SLE up to a maximum of 10mg/kg every 4 weeks;¹ **and**
 - Initial authorization is for no more than 12 months
- For **continuation of therapy**, all of the following:
 - Patient has previously received Benlysta injection for intravenous infusion; **and**
 - Documentation of positive clinical response; **and**
 - Currently receiving at least **one** standard of care treatment for active systemic lupus erythematosus (e.g., antimalarials, corticosteroids, or immunosuppressants); that is not a biologic or intravenous cyclophosphamide;^{1-7,10} **and**

Proprietary Information of United Healthcare: The information contained in this document is proprietary and the sole property of United HealthCare Services, Inc. Unauthorized copying, use and distribution of this information are strictly prohibited. Copyright 2020 United HealthCare Services, Inc.

- Benlysta is dosed according to U.S. Food and Drug Administration labeled dosing for SLE up to a maximum of 10mg/kg every 4 weeks; ¹ **and**
- Authorization is for no more than 12 months

Benlysta is unproven and not medically necessary for:

• **Antineutrophil Cytoplasmic Antibody-Associated Vasculitis**

- Severe active lupus nephritis¹
- Severe active central nervous system (CNS) lupus¹
- Use in combination with other biologics or intravenous cyclophosphamide¹
- Waldenström macroglobulinemia
- Sjögren's syndrome
- Rheumatoid arthritis

APPLICABLE CODES

The following list(s) of procedure and/or diagnosis codes is provided for reference purposes only and may not be all inclusive. Listing of a code in this policy does not imply that the service described by the code is a covered or non-covered health service. Benefit coverage for health services is determined by federal, state or contractual requirements and applicable laws that may require coverage for a specific service. The inclusion of a code does not imply any right to reimbursement or guarantee claim payment. Other Policies and Coverage Determination Guidelines may apply.

HCPCS Code	Description
J0490	Injection, belimumab, 10 mg

ICD-10 Diagnosis Code	Description
M32.0	Drug-induced systemic lupus erythematosus
M32.10	Systemic lupus erythematosus, organ or system involvement unspecified
M32.11	Endocarditis in systemic lupus erythematosus
M32.12	Pericarditis in systemic lupus erythematosus
M32.13	Lung involvement in systemic lupus erythematosus
M32.14	Glomerular disease in systemic lupus erythematosus
M32.15	Tubulo-interstitial nephropathy in systemic lupus erythematosus
M32.19	Other organ or system involvement in systemic lupus erythematosus
M32.8	Other forms of systemic lupus erythematosus
M32.9	Systemic lupus erythematosus, unspecified

BACKGROUND

Benlysta (belimumab) is a recombinant human IgG1λ monoclonal antibody specific for soluble human B lymphocyte stimulator protein (BLyS, also referred to as BAFF and TNFSF13B). Belimumab blocks the binding of soluble BLyS, a B-cell survival factor, to its receptors on B cells. Belimumab does not bind B cells directly, but by binding BLyS, belimumab inhibits the survival of B cells, including autoreactive B cells, and reduces the differentiation of B cells into immunoglobulin-producing plasma cells.

CLINICAL EVIDENCE

Proven

Systemic Lupus Erythematosus

Belimumab is indicated for the treatment of active, autoantibody-positive, systemic lupus erythematosus (SLE) who are receiving standard therapy.¹

Ginzler et al evaluated the efficacy/safety of belimumab plus standard therapy in patients (n=449) with active SLE treated up to 7 years (n=177, currently ongoing).¹⁴ Patients (n=345) who completed a double-blind, placebo-controlled, 52-week study of belimumab 1, 4, or 10 mg/kg and 24-week extension of belimumab (placebo switched to 10 mg/kg; belimumab same dose or switched to 10 mg/kg) could receive belimumab 10 mg/kg in an open-label continuation study (n=296). Disease activity was analyzed in patients with active SLE at baseline of the initial study. Efficacy endpoints measured included percentage change in the Safety of Estrogen in Lupus Erythematosus National Assessment-SLE Disease Activity Index (SELENA-SLEDAI), frequency of 1 new British Isles Lupus Assessment Group (BILAG) A or 2 new B scores, frequencies of mild-moderate and severe flares as defined by SELENA-SLEDAI Flair Index (SFI), and change in corticosteroid use. Total belimumab exposure over 7 years (double-blind⁵ and open-label periods¹⁴) was 1746 patient-years. SLE Responder Index (SRI) response rates reported at Week 52 in autoantibody-positive patients was placebo, 29%; belimumab, 46% (p<0.05). Researchers reported the following in the continuation study: 57% of auto-antibody-positive patients had an SRI response by Year 2 and 65% by Year 7; severe flares occurred in 19% with placebo and 17% with belimumab during the first year, with the annual rate declining to 2%-9% during years 2-7. Anti-dsDNA autoantibodies in patients positive for them at baseline had a progressive decline of 40%-60% from baseline over 2-7 years with belimumab. Corticosteroid use decreased over time with ≥ 50-55% reduction in median dose during years 5-7. Serious and overall annual AE rates, including infections, were generally stable or decreased during 7-year treatment. Researchers concluded that the data showed that belimumab administered over the long term with standard therapy was generally well tolerated, and sustained disease control was maintained for up to 7 years in patients with active SLE at baseline.

In a post hoc, pooled analysis of the BLISS-52 and BLISS-76 studies, Strand et al assessed the effects of belimumab treatment on health-related quality of life (HRQOL) in patients with active, autoantibody-positive systemic lupus erythematosus (SLE).¹⁷ The authors analyzed data from the major secondary endpoints of the two studies, which were the mean change in SF-36v2 Health Survey Physical Component Summary (PCS) scores at week 24. Additional pre-specified secondary endpoints included mean changes from baseline in Short Form-36 (SF-36) PCS, Mental Component Summary (MCS), and domains, Functional Assessment of Chronic Illness Therapy-Fatigue (FACIT-Fatigue) V.4, and EuroQol-5D (EQ-5D) scores at weeks 12, 24, 52 and 76 (BLISS-76 only). The SF-36, FACIT-Fatigue, and EQ-5D were administered at baseline, and weeks 4, 8, 12, 24 and 52 in both trials, and additionally at weeks 20, 32, 40, 48, 68 and 76 in BLISS-76 and week 36 in BLISS-52. Baseline SF-36 scores were 1.5 standard deviations (SDs) below age-/sex-matched US norms with similar improvement at week 24 across treatment groups. Mean changes from baseline in PCS scores were significantly (p<0.05) greater with belimumab 1 mg/kg (4.20) and 10 mg/kg (4.18) versus placebo (2.96) in BLISS-52, week 52. In BLISS-76, significantly (p<0.05) greater improvements were seen with belimumab 1 mg/kg in PCS (belimumab 1 mg/kg=4.37, 10 mg/kg=3.41 vs placebo=2.85) and Mental Component Summary (MCS) scores (belimumab 1 mg/kg=3.14, 10 mg/kg=2.70 vs placebo=1.40) at week 52, and in MCS score at week 76 (belimumab 1 mg/kg=3.05, 10 mg/kg=2.28 vs placebo=1.36), however, mean changes in PCS and MCS scores with belimumab 10mg/kg were not significantly different (week 52: PCS=3.41, MCS=2.70, and MCS week 76=2.28). In pooled analysis, there were significantly greater improvements in PCS scores with both belimumab doses versus placebo (p<0.05), and MCS scores with 1mg/kg (p<0.01). FACIT-Fatigue scores were not significantly different at week 24, however at week 52, scores improved significantly (p<0.05) with belimumab 1 and 10mg/kg vs. placebo in BLISS-52, and with 1mg/kg at weeks 52 and 76 in BLISS-76. In pooled analysis, FACIT-Fatigue scores were significantly improved (p<0.05) with both dosages at week 52, as well as weeks 8 and 12. EQ-5D utility index and VAS scores were not significantly different between treatment groups in BLISS-52. In BLISS-76, the EQ-5D VAS score was only significantly improved with belimumab 1mg/kg at week 52. The authors concluded that patients receiving belimumab reported clinically meaningful improvements in HRQOL and fatigue versus placebo, in both individual BLISS studies and by pooled analyses, that are consistent with the reductions in disease activity observed in the trials.

Unproven

Efficacy of belimumab has not been established in patients with severe active lupus nephritis or severe active CNS lupus, and belimumab has not been studied in combination with other biologic agents or IV cyclophosphamide.¹ Therefore, use of belimumab in these situations is unproven.

The use of belimumab is also being investigated for treatment of other conditions, such as, [antineutrophil cytoplasmic antibody \(ANCA\)-associated vasculitis](#)²², Waldenström macroglobulinemia^{9,16}, Sjögren's syndrome¹¹, and rheumatoid arthritis.^{12,15} Use of belimumab is considered unproven for these indications due to a lack of large, controlled clinical trials and published evidence demonstrating improved health outcomes.

Professional Societies

The European League Against Rheumatism (EULAR)

In ~~2008~~2019, EULAR published their updated recommendations for the treatment management of systemic lupus erythematosus (SLE). Their recommendations applicable to belimumab are as follows.¹⁸

General Management

Treatment of SLE

• Biologics:

- In patients with inadequate response to standard-of-care (combinations of hydroxychloroquine (HCQ) and glucocorticoids (GC) with or without immunosuppressive agents), defined as residual disease activity not allowing tapering of glucocorticoids and/or frequent relapses, add-on treatment with belimumab should be considered (1a/A).
- In organ-threatening disease refractory or with intolerance/contraindications to standard immunosuppressive agents, rituximab can be considered (2b/C). In the treatment of SLE without major organ manifestations antimalarials and/or glucocorticoids are of benefit and may be used. NSAIDs may be used judiciously for limited periods of time at patients at low risk for their complications. In non-responsive patients or patients not being able to reduce steroids below doses acceptable for chronic use, immunosuppressive agents such as azathioprine, mycophenolate mofetil, and methotrexate should also be considered.

Adjunct Therapy

- ~~Photo-protection may be beneficial in patients with skin manifestations and should be considered. Lifestyle modifications (smoking cessation, weight control, exercise) are likely to be beneficial for patient outcomes and should be encouraged. Depending on the individual medication and the clinical situation, other agents (low-dose aspirin, calcium/vitamin D, bisphosphonates, statins, anti-hypertensives (including angiotensin converting enzyme inhibitors)) should be considered. Estrogens (oral contraceptives, hormonal replacement therapy) may be used but accompanying risks should be assessed.~~

Neuropsychiatric Lupus

More expansive EULAR guidelines for neuropsychiatric lupus were published in 2010.¹⁹ Treatment guidelines are below:

- Treatment:
 - SLE patients with major neuropsychiatric manifestations considered to be of inflammatory origin (optic neuritis, acute confusional state/coma, cranial or peripheral neuropathy, psychosis, and transverse myelitis/myelopathy) may benefit from immunosuppressive therapy.

Pregnancy in Lupus

Pregnancy affects mothers with SLE and their off-springs in several ways.

- Mother:
 - There is no significant difference in fertility in lupus patients. Pregnancy may increase lupus disease activity but these flares are usually mild. Patients with lupus nephritis and anti-phospholipid antibodies are more at risk of developing pre-eclampsia and should be monitored more closely.
- Fetus:
 - SLE may affect the fetus in several ways, especially if the mother has a history of lupus nephritis, antiphospholipid, anti-Ro and/or anti-La antibodies. These conditions are associated with an increase of the risk of miscarriage, stillbirth, premature delivery, intrauterine growth restriction and fetal heart block. Prednisolone, azathioprine, hydroxychloroquine, and low dose aspirin may be used in lupus pregnancies. At present evidence suggests that mycophenolate mofetil, cyclophosphamide and methotrexate must be avoided.

Anti-Phospholipid Syndrome

In patients with SLE and anti-phospholipid antibodies, low-dose aspirin may be considered for primary prevention of thrombosis and pregnancy loss. Other risk factors for thrombosis should also be assessed. Estrogen-containing drugs increase the risk for thrombosis. In non-pregnant patients with SLE and APS-associated thrombosis, long-term anticoagulation with oral anticoagulants is effective for secondary prevention of thrombosis. In pregnant patients with SLE and anti-phospholipid syndrome combined unfractionated or LMW heparin and aspirin reduce pregnancy loss and thrombosis and should be considered.

Lupus Nephritis

More expansive EULAR guidelines for lupus nephritis were published in 2012.²⁰ Treatment guidelines are below:

- Treatment:
 - In patients with proliferative lupus nephritis, glucocorticoids in combination with immunosuppressive agents are effective against progression to end-stage renal disease. Long-term efficacy has been demonstrated only for cyclophosphamide-based regimens, which are however, associated with considerable adverse effects. In short- and medium-term trials, mycophenolate mofetil has demonstrated at least similar efficacy compared to pulse cyclophosphamide and a more favorable toxicity profile: failure to respond by 6 months should evoke discussions for intensification of therapy. Flares following remission are not uncommon and require diligent follow-up.
- End-stage renal disease:
 - Dialysis and transplantation in SLE have comparable rates for long-term patient and graft-survival as those observed in non-diabetic non-SLE patients, with transplantation being the method of choice.

U.S. FOOD AND DRUG ADMINISTRATION (FDA)

This section is to be used for informational purposes only. FDA approval alone is not a basis for coverage.

Benlysta is a B-lymphocyte stimulator (BLyS)-specific inhibitor indicated for the treatment of adult patients with active, autoantibody-positive, systemic lupus erythematosus who are receiving standard therapy.¹

Limitations of Use¹

- Benlysta is not recommended in patients with severe active lupus nephritis or severe active central nervous system lupus.
- Benlysta is not recommended to be used in combination with other biologics or intravenous cyclophosphamide.

The efficacy of Benlysta has not been evaluated in patients in these situations.

Progressive Multifocal Leukoencephalopathy (PML)¹

Cases of JC virus-associated PML resulting in neurological deficits, including fatal cases, have been reported in patients with SLE receiving immunosuppressants, including Benlysta. Risk factors for PML include:

- Testing positive for anti-JC virus (JCV) antibodies
- Longer duration of treatment with immunosuppressant therapies, including Benlysta
- Impairment of immune function

The risks and benefits of continuing treatment with Benlysta should be carefully considered in those patients who are found to be anti-JCV antibody positive and have one or more of these risk factors for PML.

Consider the diagnosis of PML in any patient presenting with new-onset or deteriorating neurological signs and symptoms and consult with a neurologist or other appropriate specialist as clinically indicated. In patients with confirmed PML, consider stopping immunosuppressant therapy, including Benlysta.

A patient's anti-JCV antibody status may be determined using an anti-JCV antibody detection test that has been analytically and clinically validated, and has been ordered by a healthcare professional. The [Stratify JCV® DxSelect™ Antibody ELISA Test](#) was cleared by FDA on January 20, 2012.²¹

The safety and efficacy of Benlysta has not been established in children.¹

In phase 3 trials, response rates for the primary endpoint were lower for African-American subjects in the Benlysta group relative to African-American subjects in the placebo group. Therefore, Benlysta should be used with caution in African-American patients.¹

Benlysta should be administered by healthcare providers prepared to manage anaphylaxis.¹

CENTERS FOR MEDICARE AND MEDICAID SERVICES (CMS)

[Medicare does not have a National Coverage Determination \(NCD\) specifically for BENLYSTA® \(belimumab\). Local Coverage Determinations \(LCDs\) do not exist at this time.](#)

[In general, Medicare covers outpatient \(Part B\) drugs that are furnished "incident to" a physician's service provided that the drugs are not usually self-administered by the patients who take them. Refer to the Medicare Benefit Policy Manual, Chapter 15, §50 - Drugs and Biologicals.](#)

[\(Accessed March 6, 2020\)](#)

~~Medicare does not have a National Coverage Determination (NCD) specifically for BENLYSTA® (belimumab). Local Coverage Determinations (LCDs) exist; see the LCDs for [Drugs and Biologics \(Non-chemotherapy\)](#).~~

~~In general, Medicare covers outpatient (Part B) drugs that are furnished "incident to" a physician's service provided that the drugs are not usually self-administered by the patients who take them. Refer to the [Medicare Benefit Policy Manual, Chapter 15, §50 - Drugs and Biologicals](#). (Accessed February 21, 2019)~~

REFERENCES

1. Benlysta [prescribing information]. Rockville, MD: Human Genome Sciences, Inc.; ~~June~~ January 2018 2020.
2. FDA Briefing Document for the Arthritis Advisory Committee Meeting: Benlysta/Belimumab. November 16, 2010. Available at: <https://wayback.archive-it.org/7993/20170404145649/https://www.fda.gov/AdvisoryCommittees/CommitteesMeetingMaterials/Drugs/ArthritisAdvisoryCommittee/ucm233578.htm>. Accessed March 11, 2019.
3. Furie R, Petri M, Zamani O, et al. A phase III, randomized, placebo-controlled study of belimumab, a monoclonal antibody that inhibits B lymphocyte stimulator, in patients with systemic lupus erythematosus. *Arthritis Rheum*. 2011 Dec;63(12):3918-30.
4. Navarra SV, Guzmán RM, Gallacher AE, et al. Efficacy and safety of belimumab in patients with active systemic lupus erythematosus: a randomised, placebo-controlled, phase 3 trial. *Lancet*. 2011 Feb 26;377(9767):721-31.
5. Wallace DJ, Stohl W, Furie RA, et al. A phase II, randomized, double-blind, placebo-controlled, dose-ranging study of belimumab in patients with active systemic lupus erythematosus. *Arthritis Rheum*. 2009 Sep 15;61(9):1168-78.
6. American College of Rheumatology Ad Hoc Committee on Systemic Lupus Erythematosus Guidelines. Guidelines for referral and management of systemic lupus erythematosus in adults. *Arthritis Rheum*. 1999 Sep;42(9):1785-96.
7. Gold Standard, Inc. Benlysta. Clinical Pharmacology [database online]. Available at: <http://www.clinicalpharmacology.com>. Accessed March 11, 2019.
8. Belimumab. In: Lexi-Drugs Online® [database on the Internet]. Hudson (OH): Lexi-Comp, Inc.; 2019 [cited March 11, 2019]. Available from: <http://online.lexi.com>. Subscription required to view.
9. Cancer Trials Australia; Human Genome Sciences. A study of belimumab in treating symptomatic Waldenstroms macroglobulinaemia. In: ClinicalTrials.gov [Internet]. Bethesda (MD): National Library of Medicine (US). 2000- [cited March 18, 2019]. Available from: <http://clinicaltrials.gov/ct2/show/NCT01142011?term=belimumab+macroglobulinemia&rank=1>. NLM Identifier: NCT01142011.
10. MCG™ Care Guidelines, ~~23rd~~ 24th edition, 2019 2020, Belimumab ACG:A-0666 (AC).
11. Mariette X, Seror R, Quartuccio L, Baron G, Salvin S, Fabris M, Desmoulin F, Nocturne G, Ravaud P, De Vita S. Efficacy and safety of belimumab in primary Sjögren's syndrome: results of the BELISS open-label phase II study. *Ann Rheum Dis*. 2015 Mar;74(3):526-31.
12. McKay J, Chwalinska-Sadowska H, Boling E, et al. Belimumab, a fully human monoclonal antibody to B-lymphocyte stimulator (BLyS), combined with standard of care therapy reduces the signs and symptoms of rheumatoid arthritis in a heterogeneous subject population. 69th Annual Scientific Meeting of the American College of Rheumatology/Association of Rheumatology Health Professionals. November 16, 2005. Oral Presentation #1920.

13. Merrill JT, Ginzler EM, Wallace DJ, et al: Long-term safety profile of belimumab plus standard therapy in patients with systemic lupus erythematosus. *Arthritis Rheum* 2012; 64(10):3364-3373.
14. Ginzler EM, Wallace DJ, Merrill JT, et al. Disease control and safety of belimumab plus standard therapy over 7 years in patients with systemic lupus erythematosus. *J Rheumatol.* 2014 Feb;41(2):300-9.
15. Stohl W, Merrill JT, McKay JD, et al. Efficacy and safety of belimumab in patients with rheumatoid arthritis: a phase II, randomized, double-blind, placebo-controlled, dose-ranging study. *J Rheumatol.* 2013 May;40(5):579-89.
16. Bishton M, Spencer A, Dickinson M, et al. A single-arm, phase II study of the anti-Blys monoclonal antibody belimumab in symptomatic Waldenstrom macroglobulinemia. *Clin Lymphoma Myeloma Leuk.* 2013 Oct;13(5):575-8.
17. Strand V, Levy RA, Cervera R, Petri MA, Birch H, Freimuth WW, Zhong ZJ, Clarke AE; BLISS-52 and -76 Study Groups. Improvements in health-related quality of life with belimumab, a B-lymphocyte stimulator-specific inhibitor, in patients with autoantibody-positive systemic lupus erythematosus from the randomized controlled BLISS trials. *Ann Rheum Dis.* 2014 May;73(5):838-44.
18. [Fanouriakis A, Kostopoulou M, Alunno A, Bertsias G, Ioannidis JP, Boletis J, et al. 2019 update of the EULAR recommendations for the management of systemic lupus erythematosus. EULAR recommendations for the management of systemic lupus erythematosus. Report of a Task Force of the EULAR Standing Committee for International Clinical Studies Including Therapeutics. *Ann Rheum Dis.* 2019 Jun;78\(6\):736-745. *Ann Rheum Dis.* 2008 Feb;67\(2\):195-205. Epub 2007 May 15.](#)
19. Bertsias GK, Ioannidis JP, Aringer M, et al. EULAR recommendations for the management of systemic lupus erythematosus with neuropsychiatric manifestations: report of a task force of the EULAR standing committee for clinical affairs. *Ann Rheum Dis.* 2010 Dec;69(12):2074-82.
20. Bertsias GK, Tektonidou M, Amoura Z, et al. Joint European League Against Rheumatism and European Renal Association-European Dialysis and Transplant Association (EULAR/ERA-EDTA) recommendations for the management of adult and paediatric lupus nephritis. *Ann Rheum Dis.* 2012 Nov;71(11):1771-82.
21. Stratify JCV® DxSelect™ [package insert]. Focus Diagnostics: Cypress, CA: November 2015.
- 21-22. [Jayne D, Blockmans D, Lugmani R, et al. Efficacy and Safety of Belimumab and Azathioprine for Maintenance of Remission in Antineutrophil Cytoplasmic Antibody-Associated Vasculitis: A Randomized Controlled Study. *Arthritis Rheumatol.* 2019 Jun;71\(6\):952-963.](#)

POLICY HISTORY/REVISION INFORMATION

Date	Action/Description
TBD	Annual review. Added ANCA-associated vasculitis as an unproven indication. Updated clinical evidence, CMS statement, and references.

INSTRUCTIONS FOR USE

This Medical Benefit Drug Policy provides assistance in interpreting UnitedHealthcare standard benefit plans. When deciding coverage, the federal, state or contractual requirements for benefit plan coverage must be referenced as the terms of the federal, state or contractual requirements for benefit plan coverage may differ from the standard benefit plan. In the event of a conflict, the federal, state or contractual requirements for benefit plan coverage govern. Before using this policy, please check the federal, state or contractual requirements for benefit plan coverage. UnitedHealthcare reserves the right to modify its Policies and Guidelines as necessary. This Medical Benefit Drug Policy is provided for informational purposes. It does not constitute medical advice.

UnitedHealthcare may also use tools developed by third parties, such as the MCG™ Care Guidelines, to assist us in administering health benefits. The UnitedHealthcare Medical Benefit Drug Policies are intended to be used in connection with the independent professional medical judgment of a qualified health care provider and do not constitute the practice of medicine or medical advice.