

CONCERT GENETIC TESTING: DERMATOLOGIC CONDITIONS

Reference Number: LA.CP.CG.03 Date of Last Revision <u>12/2306/24</u> Revision Log **Coding implications**

See <u>Important Reminder</u> at the end of this policy for important regulatory and legal information.

OVERVIEW

Genetic testing for dermatologic conditions and disorders that have many dermatologic findings may be used to confirm a diagnosis in a patient who has signs and/or symptoms of the disease. Confirming the diagnosis may alter some aspects of management and may eliminate the need for further diagnostic workup. This document addresses genetic testing for dermatologic conditions.

POLICY REFERENCE TABLE

Coding Implications

This clinical policy references Current Procedural Terminology (CPT®). CPT® is a registered trademark of the American Medical Association. All CPT codes and descriptions are copyrighted 20222023, American Medical Association. All rights reserved. CPT codes and CPT descriptions are from the current manuals and those included herein are not intended to be all-inclusive and are included for informational purposes only. -Codes referenced in this clinical policy are for informational purposes only and may not support medical necessity. Inclusion or exclusion of any codes does not guarantee coverage. -Providers should reference the most up-to-date sources of professional coding guidance prior to the submission of claims for reimbursement of covered services.

The tests and associated laboratories and CPT codes contained within this document serve only as examples to help users navigate claims and corresponding criteria; as such, they are not comprehensive and are not a guarantee of coverage or non-coverage. Please see the Concert Genetics Platform for a comprehensive list of registered tests.



NOTE: Coverage is subject to each requested code's inclusion on the corresponding LDH fee schedule. Non-covered codes are denoted (*) and are reviewed for Medical Necessity for members under 21 years of age on a per case basis. The non-covered codes will only be denoted in the table below and not throughout the policy. Please only reference the policy reference table for covered and non-covered codes.

Please see the Concert Genetics Platform for a comprehensive list of registered tests.

<u>Criteria Sections</u>	Example Tests (Labs)	Common CPT Codes	Common ICD Codes	Ref		
Known Familial Variant Analysis for Dermatologic Conditions						
Known Familial Variant Analysis for Dermatologic Conditions	Targeted Mutation Analysis for a Known Familial Variant	81403*		8		
Capillary Malforma	tion-Arteriovenous Malformation Sync	lrome (CM-AVM)			
Capillary Malformation Arteriovenous Malformation Syndrome (CM-AVM)	Capillary Malformation—Arteriovenous Malformation Syndrome (CM AVM) Panel, Sequencing and Deletion/Duplication (ARUP Laboratories)	81479	Q27.3, Q27.9	1		
Criteria Sections	Example Tests (Labs)	Common CPT Codes	Common ICD Codes	Ref		
Capillary Malforma	tion-Arteriovenous Malformation Synd	lrome (CM-AVM)			
Capillary Malformation- Arteriovenous Malformation Syndrome (CM- AVM)	Capillary Malformation- Arteriovenous Malformation Syndrome (CM-AVM) Panel, Sequencing and Deletion/Duplication (ARUP Laboratories)	81479	Q27.3, Q27.9	1		
	Vascular Malformation Sequencing Panel (Greenwood Genetic Center)					
	RASA1 Full Gene Sequencing and Deletion/Duplication (Invitae)					
	EPHB4 Full Gene Sequencing and Deletion/Duplication (Invitae)					
Congenital Ichthyos	Congenital Ichthyosis Congenital Ichthyosis					
	Ichthyosis Panel (Blueprint Genetics)	81405*, 81479	Q80	2		



Congenital Lehthyosis Multigene PanelsCongenital Lehthyosis	Ichthyosis NGS Panel (Connective Tissue Gene Tests) Invitae Congenital Ichthyosis Panel (Invitae)			
Multigene Panels Epidermolysis Bullo	osaCovered Dermatologic Conditions			
Epidermolysis Bullosa Multigene	Epidermolysis Bullosa Panel (Blueprint Genetics)	81406*, 81479	Q81	3, 4
Panels	Epidermolysis Bullosa NGS Panel (Connective Tissue Gene Tests)			
	Invitae Epidermolysis Bullosa and Palmoplantar Keratoderma Panel (Invitae)			
Covered Dermatologic Conditions				
Other Covered Dermatologic ConditionsOther Covered Dermatologic Conditions	See Below	81401*-81408*, 81479	Varies	3, 4, 5, 6, 7

OTHER RELATED POLICIES

This policy document provides criteria for Genetic Testing for Dermatologic Conditions. Please refer to:

- *Genetic Testing: Hereditary Cancer Susceptibility* for criteria related to hereditary cancer syndromes that may have or present with dermatologic findings.
- Genetic Testing: Multisystem Inherited Disorders, Intellectual Disability, and Developmental Delay for criteria related to tuberous sclerosis, neurofibromatosis, HHT, incontinentia pigmenti, proteus syndrome, pseudoxanthoma elasticum, and other disorders that affect the skin and other organ systems.
- Genetic Testing: General Approach to Genetic and Molecular Testing for criteria related to genetic testing for a dermatologic condition that is not specifically discussed in this or another more specific policy.

back to top



CRITERIA

It is the policy of Louisiana Healthcare Connections that the specific genetic testing noted below is **medically necessary** when meeting the related criteria:

KNOWN FAMILIAL VARIANT ANALYSIS FOR DERMATOLOGIC CONDITIONS

- I. Targeted mutation analysis for a known familial variant (81403*) in a dermatologic condition may be considered **medically necessary** when:
 - A. The member/enrollee has a <u>close relative</u> with a known pathogenic or likely pathogenic variant causing the condition.
- II. Targeted mutation analysis for a known familial variant (81403*) in a dermatologic condition is considered **investigational** for all other indications.

back to top

CAPILLARY MALFORMATION-ARTERIOVENOUS MALFORMATION (CM-AVM) SYNDROME

RASA1 and EPHB4 Sequencing and/or Deletion/Duplication Analysis or Multigene Panel

- I. *RASA1* and *EPHB4* sequencing and/or deletion/duplication analysis or multi-gene panel analysis (81479) to establish a diagnosis of capillary malformation-arteriovenous malformation (CM-AVM) syndrome is considered **medically necessary** when:
 - A. The member/enrollee displays one or more of the following:
 - 1. Capillary malformations, **OR**
 - 2. Arteriovenous malformations/arteriovenous fistulas, **OR**
 - 3. Parkes Weber syndrome phenotype, a cutaneous capillary malformation associated with underlying multiple micro-AVFs and soft-tissue and skeletal hypertrophy of the affected limb.
- II. *RASA1* and *EPHB4* sequencing and/or deletion/duplication analysis or multi-gene panel analysis (81479) to establish a diagnosis of capillary malformation-arteriovenous malformation (CM-AVM) syndrome is considered **investigational** for all other indications.



CONGENITAL ICHTHYOSIS

Congenital Ichthyosis Multigene Panels

- I. Multigene panel analysis to establish or confirm a diagnosis of congenital ichthyosis (81405*, 81479) is considered **medically necessary** when:
 - A. The member/enrollee has scaly skin with or without a history of harlequin ichthyosis, collodion membrane, or thick, hyperkeratotic skin, **AND**
 - B. One or more of the following:
 - 1. Ectropion (eversion of eyelids), **OR**
 - 2. Eclabium (eversion of lips), **OR**
 - 3. Scarring alopecia, OR
 - 4. Palmar and/or plantar hyperkeratosis, **OR**
 - 5. Erythroderma (red skin), AND)
 - C. The panel includes, at a minimum, the following genes: *ABCA12*, *SLC27A4*, and *TGM1*.
- II. Multigene panel analysis to establish or confirm a diagnosis of congenital ichthyosis $(81405 *_{2} 81479)$ is considered **investigational** for all other indications.

back to top

EPIDERMOLYSIS BULLOSA

Epidermolysis Bullosa Multigene Panels

- l. Multigene panel analysis to establish or confirm a diagnosis of epidermolysis bullosa (81406*, 81479) is considered **medically necessary** when:
 - A. The member/enrollee has fragility of the skin manifested by blistering with little or no trauma, **OR**
 - B. The member/enrollee has the presence of blistering that:



- 1. Is present in the neonatal period, **OR**
- 2. Primarily affects the hands and feet but can affect the whole body, OR
- 3. Occurs in annular or curvilinear groups or clusters, OR
- 4. Leads to progressive brown pigmentation interspersed with hypopigmented spots on the trunk and extremities (frequently disappears in adult life), **OR**
- 5. Is associated with palmar and plantar hyperkeratosis that may be severe, **OR**
- C. The member/enrollee has one or more of the following:
 - 1. Nail dystrophy, OR
 - 2. Milia. OR
 - 3. Congenital pyloric atresia, OR
 - 4. Natal teeth, OR
 - 5. Ureteral and renal anomalies, including hydronephrosis, ureterocele, absent bladder, dysplastic kidneys, urinary collecting system/kidney duplication, obstructive uropathy, and glomerulosclerosis, **AND**
- D. The panel includes, at a minimum, the following genes: *EXPH5*, *KRT5*, *KRT14*, *PLEC*.
- II. Multigene panel analysis to establish or confirm a diagnosis of epidermolysis bullosa (81406*, 81479) is considered **investigational** for all other indications.

back to top

OTHER COVERED DERMATOLOGIC CONDITIONS

The following is a list of conditions that have a known genetic association. Due to their relative rareness, it may be appropriate to cover these genetic tests to establish or confirm a diagnosis.

I. Genetic testing to establish or confirm one of the following dermatologic conditions to guide management is considered **medically necessary** when the member/enrollee demonstrates clinical features* consistent with the condition (the list is not meant to be comprehensive, see II below):



- A. Hidrotic Ectodermal Dysplasia 2 (Clouston Syndrome)
- B. Hypohidrotic Ectodermal Dysplasia
- C. Ocular albinism, X-linked
- D. Oculocutaneous albinism
- C. Ocular albinism, X-linked
- D. Oculocutaneous albinism
- II. Genetic testing to establish or confirm the diagnosis of all other dermatologic conditions not specifically discussed within this or another medical policy will be evaluated by the criteria outlined in *General Approach to Genetic and Molecular Testing* (see policy criteria).

*Clinical features for a specific disorder may be outlined in resources such as <u>GeneReviews</u>, <u>OMIM</u>, <u>National Library</u> of <u>Medicine</u>, <u>Genetics Home Reference</u> or other scholarly sources.

back to top

NOTES AND DEFINITIONS

- 1. Close relatives include first, second, and third degree blood relatives:
 - a. First-degree relatives are parents, siblings, and children
 - b. Second-degree relatives are grandparents, aunts, uncles, nieces, nephews, grandchildren, and half siblings
 - **C.** Third-degree relatives are great grandparents, great aunts, great uncles, great grandchildren, and first cousins

back to top

back to top

BACKGROUND AND RATIONALE

Known Familial Variant Analysis for Dermatologic Conditions

Genetic Support Foundation

The Genetic Support Foundation's Genetics 101 information on genetic testing says the following about testing for familial pathogenic variants:

Genetic testing for someone who may be at risk for an inherited disease is always easier if we know the specific genetic cause. Oftentimes, the best way to find the genetic cause is to start by testing someone in the family who is known or strongly suspected to have the disease. If their testing is positive, then we can say that we have found the familial pathogenic (harmful) variant. We can use this as a marker to test other members of the family to see who is also at risk.



Capillary Malformation-Arteriovenous Malformation Syndrome (CM-AVM)

GeneReviews: Capillary Malformation-Arteriovenous Malformation Syndrome

GeneReviews is an expert-authored review of current literature on a genetic disease, and goes through a rigorous editing and peer review process before being published online. The recommended diagnostic testing for CM-AVM is as follows:

"CM-AVM syndrome should be suspected in individuals who have any of the following:

- Capillary malformations (CMs), the hallmark of CM-AVM syndrome. CMs are generally:
 - Multifocal, atypical pink-to-reddish brown, multiple, small (1-2 cm in diameter), round-to-oval lesions sometimes with a white halo;
 - Composed of dilated capillaries in the papillary dermis
 - Mostly localized on the face and limbs;
 - Seen in combination with arteriovenous malformations (AVMs) or arteriovenous fistulas (AVF), but may be the only finding.
- AVMs/AVFs in soft tissue, bone, and brain that may be associated with overgrowth
- Parkes Weber syndrome phenotype, a cutaneous capillary malformation associated with underlying multiple micro-AVFs and soft-tissue and skeletal hypertrophy of the affected limb"

"The diagnosis of CM-AVM syndrome is established in a proband with suggestive clinical findings and a heterozygous pathogenic variant in *EPHB4* or *RASA1* identified by molecular genetic testing."

"When the phenotypic and laboratory findings suggest the diagnosis of CM-AVM syndrome, molecular genetic testing approaches can include use of a multigene panel. A multigene panel that includes *EPHB4*, *RASA1*, and other genes of interest is most likely to identify the genetic cause of the condition at the most reasonable cost while limiting identification of variants of uncertain significance and pathogenic variants in genes that do not explain the underlying phenotype."

Congenital Ichthyosis Multigene Panels

GeneReviews: Autosomal Recessive Congenital Ichthyosis

GeneReviews is an expert-authored review of current literature on a genetic disease, and goes through a rigorous editing and peer review process before being published online. The recommended diagnostic testing for nonsyndromic congenital ichthyosis is as follows:

"Autosomal recessive congenital ichthyosis (ARCI) encompasses several forms of nonsyndromic ichthyosis. Although most neonates with ARCI are collodion babies, the clinical presentation and



severity of ARCI may vary significantly, ranging from harlequin ichthyosis, the most severe and often fatal form, to lamellar ichthyosis (LI) and (nonbullous) congenital ichthyosiform erythroderma (CIE). These phenotypes are now recognized to fall on a continuum; however, the phenotypic descriptions are clinically useful for clarification of prognosis and management."

- The diagnosis of ARCI is established in a proband (typically an infant):
 - With scaly skin with or without a history of harlequin ichthyosis, collodion membrane, or thick, hyperkeratotic skin AND the later development of ONE of the following:
 - Classic lamellar ichthyosis (LI). Brown, plate-like scale over the entire body, associated with ectropion (eversion of eyelids), eclabium (eversion of lips), scarring alopecia, and palmar and plantar hyperkeratosis
 - (Nonbullous) congenital ichthyosiform erythroderma (CIE).
 Erythroderma (red skin) with fine, white scale and often with palmoplantar hyperkeratosis
 - Intermediate forms with some features of both LI and CIE, or nonLI/nonCIE form with mild hyperkeratosis;

AND/OR

• By identification of biallelic pathogenic variants in one of the genes listed below.

"The twelve genes known to be associated with ARCI are *ABCA12*, *ALOX12B*, *ALOXE3*, *CASP14*, *CERS3*, *CYP4F22*, *LIPN*, *NIPAL4*, *PNPLA1*, *SDR9C7*, *SLC27A4*, *SULT2B1*, and *TGM1*. A multigene panel that includes these genes is the diagnostic test of choice. If such testing is not available, single-gene testing can be considered starting with *ABCA12* in individuals with harlequin ichthyosis, *TGM1* in individuals with ARCI without harlequin presentation at birth and *SLC27A4* in those presenting with ichthyosis-prematurity syndrome."

Epidermolysis Bullosa Multigene Panels

GeneReviews: Epidermolysis Bullosa Simplex

GeneReviews is an expert-authored review of current literature on a genetic disease, and goes through a rigorous editing and peer review process before being published online. The recommended diagnostic testing for epidermolysis bullosa simplex and epidermolysis bullosa with pyloric atresia is as follows:

The diagnosis of epidermolysis bullosa simplex (EBS) is best established in a proband by the identification of biallelic pathogenic variants in *EXPH5* or *TGM5* or heterozygous (or rarely biallelic) pathogenic variants in *KRT5* or *KRT14* by molecular genetic testing

"The diagnosis of epidermolysis bullosa simplex (EBS) should be suspected in individuals with the following clinical findings:



- Fragility of the skin manifested by blistering with little or no trauma, which typically heals without scarring
- Blistering that:
 - May be present in the neonatal period
 - Primarily affects the hands and feet but can affect the whole body
 - Occurs in annular or curvilinear groups or clusters
 - Can lead to progressive brown pigmentation interspersed with hypopigmented spots on the trunk and extremities that frequently disappears in adult life
 - Is associated with palmar and plantar hyperkeratosis that may be severe
- Nail dystrophy
- Milia

Family history that is consistent with either an autosomal recessive or autosomal dominant inheritance pattern back to top

Note: Absence of a known family history of EBS does not preclude the diagnosis."

GeneReviews: Epidermolysis Bullosa - Pyloric Atresia

"The diagnosis of epidermolysis bullosa simplex (EBS) is best established in a proband by the identification of biallelic pathogenic variants in *EXPH5* or *TGM5* or heterozygous (or rarely biallelic) pathogenic variants in *KRT5* or *KRT14* by molecular genetic testing. A multigene panel that includes *EXPH5*, *KRT5*, *KRT14*, *TGM5* and other genes of interest may also be considered."

"Epidermolysis bullosa with pyloric atresia (EB-PA) should be suspected in newborns with the following clinical features:

- Congenital pyloric atresia with vomiting and abdominal distension resulting from complete obstruction of the gastric outlet. Radiographs reveal that the stomach is distended and filled with air
- Fragility of the skin with:
 - Blistering with little or no trauma. Blistering may be mild or severe; however, blisters generally heal with no significant scarring
 - Significant oral and mucous membrane involvement
 - Large areas of absent skin (aplasia cutis congenita), often with a thin membranous covering, affecting the extremities or head

Ureteral and renal anomalies, including hydronephrosis, ureterocele, absent bladder, dysplastic kidneys, urinary collecting system/kidney duplication, obstructive uropathy, and glomerulosclerosis."

back to top

Reviews, Revisions, and Approvals	Revision Date	Approval Date
Converted to local policy	09/23	11/27/23



Reviews, Revisions, and Approvals	Revision Date	Approval Date
Semi-annual review. Overview, coding, reference-table, background and references updated. Throughout policy: replaced "coverage criteria" with "criteria. For Other Related Policies: added "Molecular". For Congenital Ichthyosis Multigene Panels: removed "81252" throughout. For Epidermolysis Bullosa Multigene Panels: in I.A. replaced "AND" with "OR"; in I.B.1 replaced with "May be" with "Is"; in I.B.4. replaced "Can lead" with "Leads"; in I.B.5. replaced "AND" with "OR"; in I.C. added "4. Natal teeth, OR". For Other Covered Dermatologic Conditions: added "and Molecular". For Background and Rationale: replaced "inheritance patterns" with "genetic testing".	12/23	2/27/24

Semi-annual review. In Known Familial Variant Analysis for Dermatologic Conditions	06/24	
criteria, moved criteria to policy "Genetic Testing: General Approach to Genetic and		
Molecular Testing" to consolidate criteria for known familial variant tests. In		
Epidermolysis Bullosa Multigene Panels criteria, retired criteria set based on rarity of		
testing (low order volume and low claim volume). In Congenital Ichthyosis Multigene		
Panels criteria, removed minimum gene list; at present there is limited rationale for		
inclusion. Minor rewording for clarity throughout. Coding, reference-table,		
background and references updated.		

REFERENCES

- 1. Bayrak-Toydemir P, Stevenson DA. Capillary Malformation-Arteriovenous Malformation Syndrome. 2011 Feb 22 [Updated 2019 Sep 12]. In: Adam MP, Ardinger HH, Pagon RA, et al., editors. GeneReviews [Internet]. Seattle (WA): University of Washington, Seattle; 1993-2023. Available from: https://www.ncbi.nlm.nih.gov/books/NBK52764/
- 2. Richard G. Autosomal Recessive Congenital Ichthyosis. 2001 Jan 10 [Updated 2017 May 182023 April 20]. In: Adam MP, Ardinger HH, Pagon RA, et al., editors. GeneReviews [Internet]. Seattle (WA): University of Washington, Seattle; 1993-2023. Available from: https://www.ncbi.nlm.nih.gov/books/NBK1420/
- 3. So JY, Teng J. Epidermolysis Bullosa Simplex. 1998 Oct 7 [Updated 2022 Aug 4]. In: Adam MP, Ardinger HH, Pagon RA, et al., editors. GeneReviews [Internet]. Seattle (WA): University of Washington, Seattle; 1993–2023. Available from: https://www.ncbi.nlm.nih.gov/books/NBK1369/
- 4. Pfendner EG, Lucky AW. Epidermolysis Bullosa with Pyloric Atresia. 2008 Feb 22 [Updated 2017 Sep 7]. In: Adam MP, Ardinger HH, Pagon RA, et al., editors. GeneReviews [Internet]. Seattle (WA): University of Washington, Seattle; 1993–2023. Available from: https://www.ncbi.nlm.nih.gov/books/NBK1157/
- 5.3. Adam MP, Ardinger HH, Pagon RA, et al., editors. GeneReviews [Internet]. Seattle (WA): University of Washington, Seattle; 1993-2023. Available from: https://www.ncbi.nlm.nih.gov/books/NBK1116/



- 6.4. Online Mendelian Inheritance in Man, OMIM. McKusick-Nathans Institute of Genetic Medicine, Johns Hopkins University (Baltimore, MD). World Wide Web URL: https://omim.org/
- 7.5. MedlinePlus [Internet]. Bethesda (MD): National Library of Medicine (US). Available from: https://medlineplus.gov/genetics/
- 8. Genetic Support Foundation. Genetics 101 Genetic Testing: Familial Pathogenic Variant. Accessed 10/4/2022. https://geneticsupportfoundation.org/genetics-101/#

back to top

back to top

Important Reminder

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program approval status; evidence-based guidelines and positions of leading national health professional organizations; views of physicians practicing in relevant clinical areas affected by this clinical policy; and other available clinical information. LHCC makes no representations and accepts no liability with respect to the content of any external information used or relied upon in developing this clinical policy. This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved.

The purpose of this clinical policy is to provide a guide to medical necessity, which is a component of the guidelines used to assist in making coverage decisions and administering benefits. It does not constitute a contract or guarantee regarding payment or results. Coverage decisions and the administration of benefits are subject to all terms, conditions, exclusions, and limitations of the coverage documents (e.g., evidence of coverage, certificate of coverage, policy, contract of insurance, etc.), as well as to state and federal requirements and applicable LHCC administrative policies and procedures.

This clinical policy is effective as of the date determined by LHCC. The date of posting may not be the effective date of this clinical policy. This clinical policy may be subject to applicable legal and regulatory requirements relating to provider notification. If there is a discrepancy between the effective date of this clinical policy and any applicable legal or regulatory requirement, the requirements of law and regulation shall govern. LHCC retains the right to change, amend or withdraw this clinical policy, and additional clinical policies may be developed and adopted as needed, at any time.

This clinical policy does not constitute medical advice, medical treatment, or medical care.- It is not intended to dictate to providers how to practice medicine. Providers are expected to exercise professional medical judgment in providing the most appropriate care, and are solely responsible for the medical advice and treatment of member/enrollees. This clinical policy is not intended to recommend treatment for member/enrollees. Member/enrollees should consult with their treating physician in connection with diagnosis and treatment decisions.



Providers referred to in this clinical policy are independent contractors who exercise independent judgment and over whom LHCC has no control or right of control. Providers are not agents or employees of LHCC.

This clinical policy is the property of LHCC. Unauthorized copying, use, and distribution of this clinical policy or any information contained herein are strictly prohibited. -Providers, member/enrollees, and their representatives are bound to the terms and conditions expressed herein through the terms of their contracts. Where no such contract exists, providers, member/enrollees and their representatives agree to be bound by such terms and conditions by providing services to member/enrollees and/or submitting claims for payment for such services.

<u>Note:</u> ©2023 Louisiana Healthcare Connections. All rights reserved. All materials are exclusively owned by Louisiana Healthcare Connections and are protected by United States copyright law and international copyright law. No part of this publication may be reproduced, copied, modified, distributed, displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise published without the prior written permission of Louisiana Healthcare Connections. You may not alter or remove any trademark, copyright or other notice contained herein. Louisiana Healthcare Connections is a registered trademarks exclusively owned by Louisiana Healthcare Connections.