

UnitedHealthcare, Inc. ("UHC") Proprietary and Confidential Information: The information contained in this document is confidential, proprietary and the sole property of UHC. The recipient of this information agrees not to disclose or use it for any purpose other than to facilitate UHC's compliance with applicable State Medicaid contractual requirements. Any other use or disclosure is strictly prohibited and requires the express written consent of UHC.



Prostate Surgeries and Interventions (for Louisiana Only)

Policy Number: CS334LA.**AB**

Effective Date: ~~June 1, 2022~~**TBD**

[➔ Instructions for Use](#)

Table of Contents	Page
Application	1
Coverage Rationale	1
Applicable Codes	3
Description of Services	4
Clinical Evidence	5
U.S. Food and Drug Administration	42
References	42
Policy History/Revision Information	51
Instructions for Use	54

Application

This Medical Policy only applies to the state of Louisiana.

Coverage Rationale

Transurethral ablation of the prostate is proven and medically necessary in certain circumstances. ~~Transurethral ablation of the prostate is unproven and not medically necessary for all other indications due to insufficient evidence of safety and/or efficacy.~~ For medical necessity clinical coverage criteria, refer to the InterQual® ~~2022, Apr. 2022 Release~~, CP: Procedures, Prostatectomy, Transurethral Ablation.

Click [here](#) to view the InterQual® criteria.

Transurethral ablation of the prostate is unproven and not medically necessary for all other indications due to insufficient evidence of safety and/or efficacy.

~~Cryoablation of the prostate is proven and medically necessary in certain circumstances.~~ For medical necessity clinical coverage criteria, refer to the InterQual® 2022, Apr. 2022 Release, CP: Procedures, Cryoablation, Prostate. Cryoablation of the prostate is proven and medically necessary for recurrent prostate cancer diagnosed by biopsy.

Cryoablation of the prostate is proven and medically necessary for recurrent prostate cancer diagnosed by biopsy. ~~Cryoablation of the prostate is unproven and not medically necessary for initial treatment of prostate cancer and for all other indications due to insufficient evidence of safety and/or efficacy.~~ For medical necessity clinical coverage criteria, refer to the InterQual® 2022, Apr. 2022 Release, CP: Procedures, Cryoablation, Prostate.

UnitedHealthcare, Inc. ("UHC") Proprietary and Confidential Information: The information contained in this document is confidential, proprietary and the sole property of UHC. The recipient of this information agrees not to disclose or use it for any purpose other than to facilitate UHC's compliance with applicable State Medicaid contractual requirements. Any other use or disclosure is strictly prohibited and requires the express written consent of UHC.

Click [here](#) to view the InterQual® criteria.

Cryoablation of the prostate is unproven and not medically necessary for initial treatment of prostate cancer and for all other indications due to insufficient evidence of safety and/or efficacy.

Surgical prostatectomy is proven and medically necessary in certain circumstances, including for some individuals with very high risk or recurrent prostate cancer. ~~Surgical prostatectomy is unproven and not medically necessary for all other indications due to insufficient evidence of safety and/or efficacy.~~ For medical necessity clinical coverage criteria, refer to the InterQual® ~~2022, Apr. 2022 Release~~, CP: Procedures, Prostatectomy, Radical.

Click [here](#) to view the InterQual® criteria.

Surgical prostatectomy is unproven and not medically necessary for all other indications due to insufficient evidence of safety and/or efficacy.

Prostatic urethral lift (PUL) is proven and medically necessary when performed according to the following U.S. Food and Drug Administration (FDA) labeled indications, contraindications, warning and precautions:

- Treating symptoms due to urinary outflow obstruction secondary to benign prostatic hyperplasia (BPH), including lateral and median lobe hyperplasia; in men 45 years of age or older, and
- The following are not present:
 - Prostate volume of > 100 cc
 - A urinary tract infection
 - Urethra conditions that may prevent insertion of delivery system into bladder
 - Urinary incontinence due to incompetent sphincter
 - Current gross hematuria

Prostatic urethral lift (PUL) is unproven and not medically necessary for all other indications due to insufficient evidence of safety and/or efficacy.

High-energy water vapor thermotherapy for the treatment of benign prostatic hyperplasia (BPH) is proven and medically necessary when performed according to the following U.S. Food and Drug Administration (FDA) labeled indications, contraindications, warning and precautions:

- To relieve symptoms, obstructions and reduce prostate tissue in men 50 years of age or older with a prostate volume ≥ 30 cm and ≤ 80 cm, or
- Treatment of the prostate with hyperplasia of the central zone and/or a median lobe

High-energy water vapor thermotherapy for the treatment of BPH ~~benign prostatic hyperplasia in circumstances not listed above or for~~ the treatment of malignant prostate tissue is unproven and not medically necessary due to insufficient evidence of safety and/or efficacy.

The transperineal placement of biodegradable material, peri-prostatic (via needle) is proven and medically necessary for use with radiotherapy for treating prostate cancer.

UnitedHealthcare, Inc. ("UHC") Proprietary and Confidential Information: The information contained in this document is confidential, proprietary and the sole property of UHC. The recipient of this information agrees not to disclose or use it for any purpose other than to facilitate UHC's compliance with applicable State Medicaid contractual requirements. Any other use or disclosure is strictly prohibited and requires the express written consent of UHC.

The transperineal placement of biodegradable material, peri-prostatic (via needle) is unproven and not medically necessary for all other indications due to insufficient evidence of safety and/or efficacy.

The following procedures are unproven and not medically necessary due to insufficient evidence of safety and/or efficacy:

- ~~Transurethral waterjet ablation of the prostate (aquablation)~~
- Transperineal Focal focal laser ablation
- Insertion of a temporary prostatic urethral stent
- Transperineal laser ablation (TPLA)
- Transurethral waterjet ablation of the prostate (aquablation)
- Vascular embolization

Applicable Codes

The following list(s) of procedure and/or diagnosis codes is provided for reference purposes only and may not be all inclusive. Listing of a code in this policy does not imply that the service described by the code is a covered or non-covered health service. Benefit coverage for health services is determined by federal, state, or contractual requirements and applicable laws that may require coverage for a specific service. The inclusion of a code does not imply any right to reimbursement or guarantee claim payment. Other Policies and Guidelines may apply.

CPT Code	Description
<u>*0421T</u>	Transurethral waterjet ablation of prostate, including control of post-operative bleeding, including ultrasound guidance, complete (vasectomy, meatotomy, cystourethroscopy, urethral calibration and/or dilation, and internal urethrotomy are included when performed)
<u>*0582T</u>	Transurethral ablation of malignant prostate tissue by high-energy water vapor thermotherapy, including intraoperative imaging and needle guidance
<u>*0655T</u>	Transperineal focal laser ablation of malignant prostate tissue, including transrectal imaging guidance, with MR-fused images or other enhanced ultrasound
<u>*0714T</u>	<u>Transperineal laser ablation of benign prostatic hyperplasia, including imaging guidance</u>
37243	Vascular embolization or occlusion, inclusive of all radiological supervision and interpretation, intraprocedural roadmapping, and imaging guidance necessary to complete the intervention: for tumors, organ ischemia, or infarction (when performed on prostate tissue)
52441	Cystourethroscopy, with insertion of permanent adjustable transprostatic implant; single implant
52442	Cystourethroscopy, with insertion of permanent adjustable transprostatic implant; each additional permanent adjustable transprostatic implant (List separately in addition to code for primary procedure)
53850	Transurethral destruction of prostate tissue; by microwave thermotherapy
<u>*53852</u>	Transurethral destruction of prostate tissue; by radiofrequency thermotherapy

UnitedHealthcare, Inc. ("UHC") Proprietary and Confidential Information: The information contained in this document is confidential, proprietary and the sole property of UHC. The recipient of this information agrees not to disclose or use it for any purpose other than to facilitate UHC's compliance with applicable State Medicaid contractual requirements. Any other use or disclosure is strictly prohibited and requires the express written consent of UHC.

CPT Code	Description
53854	Transurethral destruction of prostate tissue; by radiofrequency generated water vapor thermotherapy
53855	Insertion of a temporary prostatic urethral stent, including urethral measurement
55866	Laparoscopy, surgical prostatectomy, retropubic radical, including nerve sparing, includes robotic assistance, when performed
55873	Cryosurgical ablation of the prostate (includes ultrasonic guidance and monitoring)
55874	Transperineal placement of biodegradable material, peri-prostatic, single or multiple injection(s), including image guidance, when performed

CPT® is a registered trademark of the American Medical Association

Codes labeled with an asterisk(*) are not on the state of Louisiana Fee Schedule and therefore not covered by the State of Louisiana Medicaid Program.

Description of Services

Benign prostatic hyperplasia (BPH) is an enlarged prostate and occurs most often during the second growth phase of the prostate (around age 25 and up). As the prostate enlarges, it presses against the urethra, which can result in the thickening of the bladder wall, the inability to empty the bladder fully, trouble starting urination, a weak flow, urgency and needing to push or strain to urinate. In most men, BPH gets worse with age and can lead to bladder and kidney damage and infection.

Several procedures have been proposed for treatment of BPH including transurethral resection of the prostate (TURP), laser vaporization or enucleation, transurethral microwave therapy, transurethral needle ablation, waterjet ablation, thermotherapy, prostatic arterial embolization, prostatectomy, prosthetic stents, transurethral incision of the prostate transurethral microwave therapy (TUMT), transurethral holmium laser ablation (HoLAP), and prostatic urethral lift (PUL) (Hayes, Inc., 2020).

Minimally invasive surgical treatments (MIST⁺s) are options to relieve symptoms of BPH while minimizing hospitalizations and complications of other treatments. One of these MIST⁺s is the Rezūm[™] System which uses thermal water vapor to reduce prostate volume associated with BPH, including hyperplasia of the central zone, and/or a middle lobe (McVary et al., 2021).

Another, the Aquabeam[®] Robotic System which uses a heat-free water jet for the ablation of benign prostate tissue.

Transperineal laser ablation (TPLA) is a minimally invasive procedure intended to ablate prostate tissue using heat from a low-powered laser which is delivered via an optical fiber inserted through the patient's perineal skin and into the prostate. The intent of TPLA is to relieve lower urinary tract symptoms in patients with BPH. A surgeon performs the procedure under local or general anesthesia using a transrectal ultrasound for guidance and inserts thin needles into the targeted area of the prostate. An optical fiber is then inserted and delivers heat which destroys the prostate tissue (ECRI, 2022).

Aquablation is performed by a medical device that allows rapid removal of prostate tissue without leaving a zone of thermal damage on the treated tissue. It utilizes a waterjet

UnitedHealthcare, Inc. ("UHC") Proprietary and Confidential Information: The information contained in this document is confidential, proprietary and the sole property of UHC. The recipient of this information agrees not to disclose or use it for any purpose other than to facilitate UHC's compliance with applicable State Medicaid contractual requirements. Any other use or disclosure is strictly prohibited and requires the express written consent of UHC.

for automated tissue resection as well as for optical energy delivery for cauterization in the treatment of BPH (Hayes, 2018).

In PUL, permanent UroLift® implants are placed to hold open the lateral lobes of the prostate in a minimally invasive procedure to reduce urinary obstruction (Roerborn et al., 2017).

Embolization places medications or synthetic materials through a catheter into a blood vessel to block blood flow to an area of the body. It may be used to control or prevent abnormal bleeding, close off vessels supplying blood to a tumor, eliminate abnormal connections between arteries and veins, or to treat aneurysms. When performed for the prostate in patients with benign prostatic hyperplasia (BPH), microspheres are injected into the prostate blood vessels, occluding the vessels which results in the gradual shrinking of the prostate tissue which widens the urethra, alleviating urinary difficulties.

Prostate cancer can be treated by surgery, medications, and/or radiotherapy. Transperineal placement of biodegradable material is sometimes used to protect other pelvic structures during radiotherapy.

Clinical Evidence

Cryoablation

In a systematic review by Hopstaken et al. (2022), the authors ~~evaluated~~evaluated the effectiveness of focal therapy in patients with localized prostate cancer. A PubMed, Embase, and The Cochrane Library were searched for studies between October 2015 and December 31, 2020. Seventy-two studies were found which included the following: 27 studies on high-intensity focused ultrasound (HIFU), 9 studies on irreversible electroporation, 11 on cryoablation, 8 on focal laser ablation and focal brachytherapy, 7 on photodynamic therapy (PDT), 2 on radiofrequency ablation, and one on prostatic artery embolization.

Of the 11 studies on cryoablation, six were retrospective studies, one of which compared HIFU with cryoablation, and five were prospective studies. No randomized controlled trials (RCT) were identified for cryotherapy. ~~The studies were mostly prospective and included a total of 5,827 patients.~~ The authors concluded primary focal therapy has potential, but continues to remain in its early stages when used for localized prostate cancer. While evidence shows improvement in ~~functional~~functional outcomes and minimal adverse effects, additional research is needed to show its oncological effectiveness. For cryotherapy, the findings are limited by the observational nature of the studies and lack of comparison groups for many of the included studies.

In a Cochrane review, Jung et al. (2018) evaluated the evidence comparing cryotherapy to standard treatment options for primary treatment of localized or locally advanced prostate cancer. A search was conducted using multiple databases (CENTRAL, MEDLINE, EMBASE), clinical trial registries and a grey literature repository (Grey Literature Report). The search resulted in two RCTs which included 307 men that were randomized into either a group for cryotherapy or radiation. The authors found uncertainty with regards to the effects of freezing the prostate when compared to radiation treatment. The evidence was of low quality and validated by study limitations which included selection bias, lack of blinding, violation of inclusion criteria and inadequate trial completion; further research is ~~needed~~needed to validate the findings.

Surgical Prostatectomy

Emam et al (2021) retrospectively compared the oncologic outcomes of radical prostatectomy (RP) versus external beam radiotherapy (EBRT) and androgen deprivation therapy for patients who had high or very high risk localized prostate cancer. There were 335 patients treated of which 291 had RP and 44 had EBRT. Outcomes that were measured were clinical characteristics, biochemical progression-free survival (BPFS), metastasis-free survival (MFS), cancer-specific survival (CSS) and overall survival (OS). The median follow-up for RP was 5.1 years versus 3.3 years for EBRT. The mean age for treatment in EBRT was older (mean 71 years vs. 61 years) and had longer PSA doubling time. Race, pre-treatment PSA and biopsy Gleason scores were similar. Three- and five-year BPFS for patients that underwent RP were 42% and 36% versus 86% and 75% after EBRT. Three- and five-year MFS for RP was 80% and 77% vs 91% and 91% after EBRT but the authors note that this was not statistically significant. Adjuvant/salvage therapy rates were 58% after RP and 29% after EBRT. Three-year CSS was 98% in both groups and OS for RP was 97% vs 94% for EBRT. The authors concluded that RP had higher rates of biochemical failure and adjuvant or salvage treatment than patients who had EBRT. MFS also trended toward benefit after EBRT, however, CSS and OS remained high in both groups. Limitations of the study included the wide range of median follow-up between the two groups which may have led to fewer mortality events for EBRT and may limit the differences seen in MFS. Additional limitations include observational design, imbalance between EBRT and RP cases, inability to adjust for confounding factors which have been shown to be unbalanced between EBRT and RP, and that this was a single center retrospective analysis.

Varca et al. (2018) evaluated the functional safety of laparoscopic radical prostatectomy (LRP) for 1,114 high-risk prostate cancer (PCa) patients. The average age of the patient was 62 years and mean follow up was 74 ± 50 months. The overall complication rate was 5.2% with most of these needing blood transfusions which aligned with other studies. An overall patient survival rate of 96.6% and a disease-free survival of 66.2% was reported. The authors concluded that radical prostatectomy represents an effective treatment for patients with high risk PCa and that the laparoscopy approach is a safe way to offer a mini-invasive alternative to open surgery. The study is however limited by lack of comparison group.

Petrelli et al. (2014) conducted a systematic review and meta-analysis and compared the outcomes of patients treated with ~~radical prostatectomy (RP)~~ versus external beam radiotherapy (RT). A search was conducted using MEDLINE, EMBASE, Web of Science, SCOPUS, and The Cochrane Central Register of Controlled Trials (CENTRAL) and returned one RCT and 16 retrospective studies. The primary end points were overall survival and PC-specific mortality (PCSM), and the secondary end points were non-PCSM and the risk of biochemical relapse (b-RFS). Seven studies (1 prospective and 6 retrospective cohort studies) reported overall survival data, ten studies presented data for PCSM (1 prospective and 9 retrospective cohort studies), seven studies permitted the calculation of non-PCSM and data for b-RFS were available in 8 studies. The authors found RT combined with androgen deprivation therapy (ADT) continues to remain the treatment of choice for high-risk prostate cancer patients, ~~however the~~; it is believed that radical prostatectomy should be considered in selected high-risk prostate cancer patients that have minimal or no comorbidities and an average life expectancy. Limitations included physician bias in choice of treatment in each cohort, the RT group consisted of patients who were likely older than the RP patients, and an imbalance in the use of adjuvant therapies amongst the studies.

UnitedHealthcare, Inc. ("UHC") Proprietary and Confidential Information: The information contained in this document is confidential, proprietary and the sole property of UHC. The recipient of this information agrees not to disclose or use it for any purpose other than to facilitate UHC's compliance with applicable State Medicaid contractual requirements. Any other use or disclosure is strictly prohibited and requires the express written consent of UHC.

Prostatic Urethral Lift (PUL)

Kaplan-Marans et al. (2022) searched the FDA MAUDE database for reports of device malfunction and procedural complications for novel treatments for BPH. The results showed a total of 391 adverse events reported, 132 were related to PUL, and 69 were severe or life threatening. There were 25 cases in which the PUL implants required removal, or caused a problem with a later procedure, and 21 cases requiring blood transfusion. The most common device malfunctions were 27 cases of device misfire and 21 cases of needle fragmentation. The authors concluded these complications and device related malfunctions are significant, and patients and providers should be aware of the potential risks when choosing this treatment for BPH.

In 2017, Roehrborn et al. published five-year outcomes of the prospective, multi-center, randomized, blinded sham control trial of the Prostatic Urethral Lift (PUL) in men with bothersome lower urinary tract symptoms (LUTS) due to benign prostatic hyperplasia (BPH). In this 19-center study, 206 subjects ≥ 50 years old with an International Prostate Symptom Score (IPSS) > 12 , peak flow rate (Qmax) ≤ 12 mL/s, and prostate volume 30 cc-80 cc were randomized 2:1 to the PUL procedure or blinded sham control. IPSS improvement after PUL was 88% greater than that of sham at 3 months. LUTS and QOL were significantly improved by 2 weeks with return to preoperative physical activity within 8.6 days. Improvement in international prostate symptom score (IPSS), QOL, BPH Impact Index (BPHII), and maximum flow rate (Qmax) were durable through 5 years with improvements of 36%, 50%, 52%, and 44% respectively. Symptom improvement was commensurate with patient satisfaction. The authors conclude that PUL offers a durable, minimally invasive option in the treatment of LUTS due to BPH.

Two-year outcomes were reported by Gratzke et al. (2017) for the BPH6 prospective, multicenter, non-blinded randomized study (n=80) which compared PUL to transurethral resection of the prostate (TURP). Inclusion criteria were aged ≥ 50 years and a candidate for TURP, with IPSS > 12 , maximum urinary flow rate (Q max) ≤ 15 mL/s, and prostate volume ≤ 60 cc on ultrasonography. Parallel 1:1 randomization was performed using permuted blocks of random sizes, stratified by study site. Patients were followed up with visits at 2 weeks, 1 month, 3 months, 6 months, 1 year and 2 years. Significant improvements in IPSS, IPSS QoL, BPH Impact Index (BPHII) and Q-max were observed in both arms through 2-year follow-up. IPSS change with TURP was superior to that with PUL at 1 and 2 years, and TURP was superior with regard to Q max at all time points (Table 1). HRQoL and BPHII improvements were not statistically different. Quality of recovery, as defined by at least a score of 70 on the QoR VAS (0-100 scale), was superior for PUL compared with TURP, with 82% of patients in the PUL arm achieving the recovery endpoint by 1 month compared with 53% of patients in the TURP arm (P = 0.008). The results demonstrate that both the PUL and TURP procedures offered significant improvement in symptoms, Q max and HRQoL. The modest patient number may not have provided sufficient statistical power to detect differences in some of the secondary outcome variables.

High Energy Water Vapor Thermotherapy of Malignant Prostate Tissue

A search of the literature did not identify relevant peer reviewed original data publications.

High Energy Water Vapor Thermotherapy of Benign Prostate Hypertrophy (BPH)

Kaplan-Marans et al. (2022) searched the FDA MAUDE database for reports of device malfunction and procedural complications for novel treatments for BPH. The results showed

UnitedHealthcare, Inc. ("UHC") Proprietary and Confidential Information: The information contained in this document is confidential, proprietary and the sole property of UHC. The recipient of this information agrees not to disclose or use it for any purpose other than to facilitate UHC's compliance with applicable State Medicaid contractual requirements. Any other use or disclosure is strictly prohibited and requires the express written consent of UHC.

a total of 391 adverse events reported, 157 were related to transurethral water vapor therapy, and 24 were severe or life threatening. Blood transfusion frequency was 1%. The most common device malfunction in 19 cases was "unresolvable error message". The authors concluded these complications and device related malfunctions may be significant, and patients and providers should be aware of the potential risks when choosing this treatment for BPH.

McVary et al. (2021, included ~~—reviewed~~ in the Hayes report below) presented the final five-year outcomes of the multicenter, randomized, sham-controlled trial of water vapor therapy (Rezūm™) for treatment of moderate to severe lower urinary tract symptoms due to benign prostatic hyperplasia. A total of 197 men >50 years of age with an International Prostate Symptom Score (IPSS) ≥13, maximum flow rate ≤15 ml/second and prostate volume 30 to 80 cc were randomized and followed for 5 years. One hundred thirty-six subjects were randomized to the Rezūm™ treatment-arm and 61 subjects were randomized to the control-arm. Fifty-three control subjects requalified for crossover, received treatment and participated in follow-up through 5 years. All treatments were performed in the physician's office or an ambulatory surgery center. No participants withdrew due to any procedure or device-related adverse events (AEs). In the treatment group, eighteen participants were lost to follow-up and 13 withdrew consent. Five subjects were censored for protocol non-compliance, and 2 were withdrawn for prostate cancer treatment. This resulted in the data of 77 participants being analyzed and reported. Within the crossover group, 21 subjects completed 5 years of follow-up. Ten subjects withdrew consent and an additional 5 were lost to follow-up. One subject was censored for non-compliance, one due to cancer diagnosis, and one subject died due to unrelated causes. The results showed there was significant improvement of lower urinary tract symptoms observed at <3 months post treatment and that it remained sustained through 5 years in the treatment group, and the results were similar in the crossover group. IPSS reduced 48%, quality of life increased 45%, maximum flow rate improved 44%, and Benign Prostatic Hyperplasia Impact Index decreased 48%. Surgical re-treatment rate was 4.4% with no reports of device or procedure related sexual dysfunction or sustained de novo erectile dysfunction. The authors concluded that treatment of BPH with minimally invasive water vapor thermal therapy results in significant and durable relief of symptoms, results in low resurgery rates and does not impact sexual function. Also, it was demonstrated that the versatility of water vapor thermal therapy especially among MISTs, as the Rezūm System is capable of ablating tissue from lateral lobes as well as tissue of an enlarged central zone (i.e., middle lobe or median bar) without advanced training or learning a complex technique. The authors acknowledged that participant attrition is a limitation of this long-term study, but this is typical for many studies of prostate treatments. There was also a lack of urodynamic testing to assess the impact of these results on bladder function. The same author (McVary et al., reviewed in the Hayes and ECRI reports as well as the Kang, et al systematic review below) reported on the early results of this trial in 2016, in which the results showed baseline IPSS score of 22.0 for subjects receiving thermal therapy decreased at 3months to 10.8, while in the sham control group, it decreased from 21.9 to 17.5 (p<0.0001 for group differences). The peak flow rate increased by 6.2 ml per second at 3 months and sustained throughout the 12-month follow up period, while it only increased by 0.5 in the sham control group at 3 months (p<0.0001 for group differences). Adverse events were mild to moderate severity and most resolved within 3 weeks, and no de novo erectile dysfunction was reported. In 2018, McVary and Roehrborn (reviewed in the Hayes and ECRI reports below) reported the three-year outcomes which showed IPSS improvement of 160% at 3 months compared to control, and at least a 50% improvement in IPSS, quality of life, Qmax and BPH Impact Index over the three-year period with no adverse events, and no de novo erectile dysfunction reported.⁷ The authors noted a surgical retreatment rate of 4.4% over the three-year period.

UnitedHealthcare, Inc. ("UHC") Proprietary and Confidential Information: The information contained in this document is confidential, proprietary and the sole property of UHC. The recipient of this information agrees not to disclose or use it for any purpose other than to facilitate UHC's compliance with applicable State Medicaid contractual requirements. Any other use or disclosure is strictly prohibited and requires the express written consent of UHC.

A 2021 Hayes health technology assessment concluded that based on a low quality of evidence from one randomized controlled trial, two comparative cohort studies and mostly single arm observational studies, treatment with the Rezūm water vapor therapy system may alleviate lower urinary tract symptoms (LUTS) related to BPH in the short and medium term without an impact on sexual function or serious side effects. Substantial uncertainty remains due to the lack of high-quality studies that compare Rezūm with the surgical alternatives such as TURP.

Miller et al (2020) performed a systematic review and meta-analysis of published studies regarding water vapor thermal therapy (WVTT) for symptomatic BPH. Five cohorts comprised of 514 patients (including 40% with median lobe obstruction) were included in this meta-analysis. The mean prostate volume ranged from 45 to 53 cc, and 57% of patients presented with severe BPH symptoms and urinary flow obstruction was common. Patients received an average of 5 water vapor injections per procedure. Patient follow-up ranged from 6 to 48 months. The results showed minimal clinically important difference units (MCID) in the IPSS from 3.3 to 3.8, IPSS-QOL from 3.9 to 4.6, BPHII from 6.8 to 8.2, and 1.5 to 3.0 for Qmax scores at all follow up intervals. These were significant improvements. Surgical retreatment rate was 7 % at 4--year follow up. The most common adverse events were nonserious and transient and included dysuria, urinary retention, and urinary tract infection. There were no cases of de novo erectile dysfunction. The authors concluded that WVTT --improves BPH symptoms that exceeds MCID thresholds, preserves sexual function and has a low ~~resurgicatre~~ surgical rate, suggesting it may be a valuable treatment option for men with LUTS due to BPH.

In a 2020 Cochrane Database Systematic Review, Kang et al. assessed the effects of convective radiofrequency water vapor thermal therapy for the treatment of lower urinary tract symptoms in men with benign prostatic hyperplasia. Included was a single industry sponsored randomized controlled trial that included 197 men that compared this therapy to a sham procedure. In this review, the authors only found short term data measured up to 3 months urologic symptom scores and quality of life appear to improve with convective radiofrequency water vapor thermal therapy, but therapy but are very uncertain about major adverse events. The certainty of evidence ranged from moderate to very low, with study limitations and imprecision being the most common reasons for rating down. These findings are based on a single industry sponsored study, with three-month short-term follow-up. The authors did not find any studies comparing convective radiofrequency water vapor thermal therapy to any other active treatment.

~~A 2020 NICE medical technology guideline for the use of Rezum for treating lower urinary tract symptoms secondary to benign prostatic hyperplasia states that the evidence supports adopting Rezum for treating lower urinary tract symptoms (LUTS) caused by benign prostatic hyperplasia (BPH) in the NHS. It should be considered as a treatment option for men with moderate to severe LUTS (International Prostate Symptoms Score [IPSS] typically 13 or over), and a moderately enlarged prostate (typically between 30 cm and 80 cm).~~

In a 2016 ECRI clinical evidence assessment, updated in January 2022, regarding the Rezūm System (Boston Scientific Corp.) for Treating Benign Prostatic Hyperplasia, it was concluded that the evidence from one randomized controlled trial (RCT) and three pre-post studies suggests the Rezūm System decreases lower urinary tract symptoms (LUTS) for up to four years compared with pretreatment symptoms. Studies reported nonserious adverse events (AEs) that resolved with usual care, and patients maintained sexual function. There were no studies that compared Rezūm to any other BPH treatment, and ongoing studies will not address this evidence gap.

UnitedHealthcare, Inc. ("UHC") Proprietary and Confidential Information: The information contained in this document is confidential, proprietary and the sole property of UHC. The recipient of this information agrees not to disclose or use it for any purpose other than to facilitate UHC's compliance with applicable State Medicaid contractual requirements. Any other use or disclosure is strictly prohibited and requires the express written consent of UHC.

Transperineal Placement of Biodegradable Material

The SpaceOAR is used to temporarily position the anterior rectal wall away from the prostate during radiotherapy for prostate cancer and in creating this space it is the intent to reduce the radiation dose delivered to the anterior rectum. The absorbable spacer maintains space for the entire course of prostate radiotherapy treatment and is completely absorbed by the patient's body over time.

A Hayes health technology assessment (2021) summarized that while published evidence suggests a potential benefit of an absorbable perirectal spacer (APS) during radiation therapy for prostate cancer, there is substantial uncertainty regarding its safety and efficacy; future studies are needed to assess the APS clinical usefulness and cost-effectiveness.

In a custom product brief, ECRI (2020) concludes that SpaceOAR hydrogel is well tolerated and works as intended to reduce rectal irradiation long-term, but not acute, rectal toxicity, and it improves bowel quality of life (QOL), based on one randomized controlled trial and four prospective nonrandomized comparative studies.

Afkhami Ardekani and Ghaffari (2020) evaluated the effect of dosimetry and procedure toxicity of polyethylene glycol (PEG)-based hydrogel spacers during prostate brachytherapy. There were twelve studies included in the systematic review involving 615 patients. The approach used to place the hydrogel spacers was hydrodissection and considered one of the most common techniques. Ultrasonography is used to insert a large gauge needle where saline water is injected to create potential space between the prostate and anterior rectal wall; PEG hydrogel is then injected into the created space. The DuraSeal and SpaceOar then polymerize within 3 and 10 seconds after injection. The authors found the data of several studies revealed the rectal dosimetry was significantly reduced with the use of the PEG hydrogel spacers and that the procedure was safe. The authors concluded the implantation of PEG hydrogel spacers is practical and safe with well tolerance of the procedure. The use of PEG hydrogels for prostate brachytherapy has a very high success rate, however the advantages of these spacers should be weighed against possible risks of complications. Additional RCTs should be done to further clarify rectal dose reduction on toxicity and quality of life.

A systematic review was conducted by Vaggers et al. (2020) from nine full text articles reviewing polyethylene glycol-based hydrogel rectal spacers for prostate brachytherapy. Four studies used the DuraSeal Spinal Sealant and five studies used SpaceOar. Primary outcomes included procedure complications, failures, prostate-rectum separation, rectal dosimetry and GI toxicities for hydrogel insertion. There was little variation in technique used throughout the articles reviewed. The authors found the studies demonstrated a significant reduction in rectal dosimetry and concluded that the polyethylene glycol-based hydrogel rectal spacers appear to be safe and easy. Even though the spaces appear to reduce rectal toxicity, further studies are needed to confirm these findings. Limitations include the review as retrospective and non-randomization along with small sample size.

Paetkau et al. (2019) retrospectively evaluated 13 patients with SpaceOAR implant to determine future planning needs for patients with prostate cancer undergoing radiation therapy. Computerized tomography (CT) scans were taken pre- and post-implant. A prescription of 60 Gy in 20 fractions was planned on both scans. Six treatment plans were produced per anonymized dataset using either a structure of rectum plus the hydrogel,

UnitedHealthcare, Inc. ("UHC") Proprietary and Confidential Information: The information contained in this document is confidential, proprietary and the sole property of UHC. The recipient of this information agrees not to disclose or use it for any purpose other than to facilitate UHC's compliance with applicable State Medicaid contractual requirements. Any other use or disclosure is strictly prohibited and requires the express written consent of UHC.

termed composite rectum wall (CRW), or rectal wall (RW) as an inverse optimization structure and intensity modulated radiotherapy (IMRT) or volumetric modulated arc therapy (VMAT) as a treatment technique. Dose-volume histogram metrics were compared between plans to determine which optimization structure and treatment technique offered the maximum rectal dose sparing. RW structures offered a statistically significant decrease in rectal dose over CRW structures, whereas the treatment technique (IMRT vs VMAT) did not significantly affect the rectal dose. There was improvement seen in bladder and penile bulb dose when VMAT was used as a treatment technique. The authors concluded that overall, treatment plans using the RW optimization structure offered the lowest rectal dose while VMAT treatment technique offered the lowest bladder and penile bulb dose.

Wu et al. (2018, included in Afkhami Ardekani and Ghaffari (2020) and Vaggers et al. (2020) systematic reviews above) evaluated 18 consecutive patients underwent transperineal ultrasound-guided placement of 10 cc of SpaceOAR hydrogel prior to HDR brachytherapy in the treatment of prostate cancer. Treatment plans were generated using an inverse planning simulated annealing algorithm. Rectal dosimetry for these 18 patients was compared with the 36 preceding patients treated with HDR brachytherapy without SpaceOAR. There was no difference in age, pretreatment prostate-specific antigen, Gleason score, clinical stage, prostate volume, or contoured rectal volume between those who received SpaceOAR and those who did not. Patients who received SpaceOAR hydrogel had significantly lower dose to the rectum as measured by percent of contoured organ at risk (median, V80 < 0.005% vs. 0.010%, $p = 0.003$; V75 < 0.005% vs. 0.14%, $p < 0.0005$; V70 0.09% vs. 0.88%, $p < 0.0005$; V60 = 1.16% vs. 3.08%, $p < 0.0005$); similar results were seen for rectal volume in cubic centimeters. One patient who received SpaceOAR developed a perineal abscess 1 month after treatment. The authors concluded that transperineal insertion of SpaceOAR hydrogel at the time of HDR brachytherapy is feasible and decreases rectal radiation dose. Further investigation is needed with well-designed clinical trials and larger patient populations to further assess the clinical impact.

Chao et al. (2018) conducted a prospective case series analysis to report on the dosimetric benefits and late toxicity outcomes following injection of a hydrogel spacer between the prostate and rectum in 76 patients with T1-T3a prostate cancer treated with radiotherapy. There were no postoperative complications reported. Mean prostate size were 66.0cc (25.0cc - 187.0cc). Rectal dose volume parameters were observed with volume of rectum receiving 70Gy (rV70), 75Gy (rV75) and 78Gy (rV78) were 7.8%, 3.6% and 0.4%. 21% (16/76) developed acute grade 1 GI toxicities but all were resolved completely by 3 months post-treatment. 3% (2/76) developed late grade 1 GI toxicities. No patients experienced acute or late grade 2+ GI toxicities. The authors concluded that injection of hydrogel spacer resulted in a reduction of irradiated rectal dose volumes along with minimal GI toxicities, irrespective of prostate size. Additional studies with longer-term outcomes are needed to evaluate long-term toxicities. The findings are limited by lack of comparison group.

Taggar et al. (2018, included in Afkhami Ardekani and Ghaffari (2020) and Vaggers et al. (2020) systematic review above) conducted a prospective cohort study to evaluate placement of an absorbable rectal hydrogel spacer in 74 patients with prostate cancer undergoing low-dose-rate brachytherapy with palladium-103. Rectal dosimetry was compared with a consecutive cohort of 136 patients treated with seed implantation without a spacer. On average, 11.2-mm (SD 3.3) separation was achieved between the prostate and the rectum. The resultant mean rectal volume receiving 100% of prescribed dose (V100%), dose to 1 cc of rectum (D1cc), and dose to 2 cc of rectum (D2cc) were 0 (SD 0.05 cc), 25.3% (SD 12.7), and 20.5% (SD 9.9), respectively. All rectal dosimetric parameters improved significantly for the cohort with spacer placement as compared with the non-spacer

UnitedHealthcare, Inc. ("UHC") Proprietary and Confidential Information: The information contained in this document is confidential, proprietary and the sole property of UHC. The recipient of this information agrees not to disclose or use it for any purpose other than to facilitate UHC's compliance with applicable State Medicaid contractual requirements. Any other use or disclosure is strictly prohibited and requires the express written consent of UHC.

cohort. Injection of rectal spacer is feasible in the post-LDR brachytherapy setting and reduces dose to the rectum with minimal toxicity. Prostate and urethral dosimetries do not appear to be affected by the placement of a spacer.

Pinkawa et al. (2017a) reported 5-year outcomes of a cohort study after prostate cancer radiation therapy with and without the use of a hydrogel spacer. Fifty-four patients were selected to receive a hydrogel spacer. Patients were surveyed before RT; at the last day of RT; and a median time of 2 months, 17 months, and 63 months after RT. For patients treated with a hydrogel spacer, mean bowel function and bother score changes of >5 points in comparison with baseline levels were found only at the end of RT (10-15 points; $P<.01$). No spacer patient reported moderate or big problems with his bowel habits overall. Mean bother score changes of 21 points at the end of RT, 8 points at 2 months, 7 points at 17 months, and 6 points at 63 months after RT were found for patients treated without a spacer. A bowel bother score change >10 points was found in 6% versus 32% ($P<.01$) at 17 months and in 5% versus 14% ($P=.2$) at 63 months with versus without a spacer. The authors conclude that hydrogel spacer application demonstrates excellent treatment tolerability, in particular regarding bowel problems. They encourage further studies with dose-escalated or re-irradiation concepts.

Pinkawa et al. (2017b) evaluated in a cohort study of 167 consecutive patients who received prostate RT with 2 Gy fractions up to 76 Gy (without hydrogel, $n = 66$) or 76-80 Gy (with hydrogel, $n = 101$). The numbers of interventions resulting from bowel problems during the first 2 years after RT were compared. Patients were surveyed prospectively before RT, at the last day of RT, and at a median of 2 and 17 months after RT using a validated questionnaire (Expanded Prostate Cancer Index Composite). Treatment for bowel symptoms (0 vs. 11%; $p < 0.01$) and endoscopic examinations (3 vs. 19%; $p < 0.01$) were performed less frequently with a spacer. Mean bowel function scores did not change for patients with a spacer in contrast to patients without a spacer (mean decrease of 5 points) >1 year after RT in comparison to baseline, with 0 vs. 12% reporting a new moderate/big problem with passing stools ($p < 0.01$). It was noted that statistically significant differences were found for the items "loose stools", "bloody stools", "painful bowel movements" and "frequency of bowel movements". The authors concluded that spacer injection is associated with a significant benefit for patients after prostate cancer RT.

Hamstra et al. (2017) reported the final outcomes from their single-blind phase III trial of image guided intensity modulated radiation therapy ($n=222$). The 3-year incidence of grade ≥ 1 (9.2% vs 2.0%; $P=.028$) and grade ≥ 2 (5.7% vs 0%; $P=.012$) rectal toxicity favored the spacer arm. Grade ≥ 1 urinary incontinence was also lower in the spacer arm (15% vs 4%; $P=.046$), with no difference in grade ≥ 2 urinary toxicity (7% vs 7%; $P=0.7$). From 6 months onward, bowel QOL consistently favored the spacer group ($P=.002$), with the difference at 3 years (5.8 points; $P<.05$) meeting the threshold for a MID. The control group the authors reported that the benefit of a hydrogel spacer in reducing the rectal dose, toxicity, and QOL declines after image guided intensity modulated radiation therapy for prostate cancer was maintained or increased with a longer follow-up period, providing stronger evidence for the benefit of hydrogel spacer use in prostate radiation therapy. Additional long-term outcomes are needed to determine the benefits of hydrogel spacers.

In a prospective, randomized patient-blinded clinical study, Karsh et al. (2017) compared image-guided intensity modulated prostate radiotherapy (79.2Gy in 44 fractions) in men with or without insertion of prostate-rectum hydrogel spacer (SpaceOar). The mean additional space created between the prostate and the rectum was just over 1cm, which allowed significant rectum and penile bulb radiation dose reduction resulting in less

UnitedHealthcare, Inc. ("UHC") Proprietary and Confidential Information: The information contained in this document is confidential, proprietary and the sole property of UHC. The recipient of this information agrees not to disclose or use it for any purpose other than to facilitate UHC's compliance with applicable State Medicaid contractual requirements. Any other use or disclosure is strictly prohibited and requires the express written consent of UHC.

acute pain, lower rates of late rectal toxicity, and improved bowel and urinary QOL scores from 6 months through the 3-year follow-up period as compared to the control group. The authors concluded that spacer application significantly reduced rectal radiation dose, resulting in long-term reductions in rectal toxicity, as well as improvements in bowel, urinary, and sexual QOL. Patient sample volumes were not reported.

Yeh et al. (2016, included in Afkhami Ardekani and Ghaffari (2020) and Vaggers et al. (2020) systematic reviews above) studied rectal toxicity rates in 326 patients administered a polyethylene glycol (PEG) hydrogel rectal spacer in conjunction with combination high-dose-rate brachytherapy at 16 Gy (average dose 15.5 Gy; standard deviation [SD] = 1.6 Gy) and external beam radiotherapy of 59.4 Gy (average dose 60.2 Gy; SD = 2.9 Gy). Clinical efficacy was determined by measuring acute and chronic rectal toxicity using the National Cancer Center Institute Common Terminology Criteria for Adverse Events v4.0 grading scheme. Median follow-up was 16 months. The mean anterior-posterior separation achieved was 1.6 cm (SD = 0.4 cm). Rates of acute Grade 1 and 2 rectal toxicity were 37.4% and 2.8%, respectively. There were no acute Grade 3/4 toxicities. Rates of late Grade 1, 2, and 3 rectal toxicity were 12.7%, 1.4%, and 0.7%, respectively. There were no late Grade 4 toxicities. The authors concluded that acute and chronic rectal toxicities are low despite aggressive dose escalation. Longer term outcomes are needed to evaluate impact.

Mariados et al. (2015) conducted a prospective multicenter randomized controlled pivotal trial to assess outcomes following absorbable spacer (SpaceOAR system) implantation. The study included 222 patients with clinical stage T1 or T2 prostate cancer who underwent computed tomography (CT) and magnetic resonance imaging (MRI) scans for treatment planning, followed with fiducial marker placement. Patients were randomized to receive spacer injection or no injection (control). Spacer safety and impact on rectal irradiation, toxicity, and QOL were assessed throughout 15 months. Spacer application had a 99% hydrogel placement success rate. The authors reported that there were no device-related AEs, rectal perforations, serious bleeding, or infections within either group. Overall acute rectal adverse event rates were similar between groups, with fewer spacer patients experiencing rectal pain (P=.02). There was no late rectal toxicity greater than grade 1 in the spacer group. At 15 months 11.6% and 21.4% of spacer and control patients, respectively, experienced 10-point declines in bowel QOL. MRI scans at 12 months verified spacer absorption. The authors concluded that spacer application was well tolerated. Increased perirectal space reduced rectal irradiation, reduced rectal toxicity severity, and decreased rates of patients experiencing declines in bowel QOL. The spacer appears to be an effective tool, potentially enabling advanced prostate radiation therapy protocols. However, the short follow-up period is a study limitation, as researchers have published the median time to late gastrointestinal grade >2 toxicity onset was 17 months. The study was also limited by the exclusion of patients with prostate volumes >80 mL, patients with extracapsular extension, and those with prior radiation or surgery. Patients with extracapsular extension have the theoretical risk of pushing posterior extracapsular disease farther from the prostate during radiation therapy, whereas patients with prior radiation or surgery may have perirectal scar formation, limiting space creation. The authors noted that the use of spacers in these populations should proceed cautiously in separate clinical trials.

Prostatic Urethral Lift (PUL)

Kaplan Marans et al. (2022) searched the FDA MAUDE database for reports of device malfunction and procedural complications for novel treatments for BPH. The results showed a total of 391 adverse events reported, 132 were related to PUL, and 69 were severe or life threatening. There were 25 cases in which the PUL implants required removal, or caused a problem with a later procedure, and 21 cases requiring blood transfusion. The

UnitedHealthcare, Inc. ("UHC") Proprietary and Confidential Information: The information contained in this document is confidential, proprietary and the sole property of UHC. The recipient of this information agrees not to disclose or use it for any purpose other than to facilitate UHC's compliance with applicable State Medicaid contractual requirements. Any other use or disclosure is strictly prohibited and requires the express written consent of UHC.

~~most common device malfunctions were 27 cases of device misfire and 21 cases of needle fragmentation. The authors concluded these complications and device related malfunctions are significant, and patients and providers should be aware of the potential risks when choosing this treatment for BPH.~~

~~In 2017, Roehrborn et al. published five year outcomes of the prospective, multi-center, randomized, blinded sham control trial of the Prostatic Urethral Lift (PUL) in men with bothersome lower urinary tract symptoms (LUTS) due to benign prostatic hyperplasia (BPH). In this 19-center study, 206 subjects ≥ 50 years old with an International Prostate Symptom Score (IPSS) > 12 , peak flow rate (Q_{max}) ≤ 12 mL/s, and prostate volume 30 cc ≤ 80 cc were randomized 2:1 to the PUL procedure or blinded sham control. IPSS improvement after PUL was 88% greater than that of sham at 3 months. LUTS and QOL were significantly improved by 2 weeks with return to preoperative physical activity within 8.6 days. Improvement in IPSS, QOL, BPHII, and Q_{max} were durable through 5 years with improvements of 36%, 50%, 52%, and 44% respectively. Symptom improvement was commensurate with patient satisfaction. The authors conclude that PUL offers a durable, minimally invasive option in the treatment of LUTS due to BPH.~~

~~Two year outcomes were reported by Gratzke et al. (2017) for the BPH6 prospective, multicenter, non-blinded randomized study (n=80) which compared PUL to transurethral resection of the prostate (TURP). Inclusion criteria were aged ≥ 50 years and a candidate for TURP, with IPSS > 12 , maximum urinary flow rate (Q_{max}) ≤ 15 mL/s, and prostate volume ≤ 60 cc on ultrasonography. Parallel 1:1 randomization was performed using permuted blocks of random sizes, stratified by study site. Patients were followed up with visits at 2 weeks, 1 month, 3 months, 6 months, 1 year and 2 years. Significant improvements in IPSS, IPSS QoL, BPH Impact Index (BPHII) and Q_{max} were observed in both arms through 2-year follow-up. IPSS change with TURP was superior to that with PUL at 1 and 2 years, and TURP was superior with regard to Q_{max} at all time points (Table 1). HRQoL and BPHII improvements were not statistically different. Quality of recovery, as defined by at least a score of 70 on the QoR VAS (0-100 scale), was superior for PUL compared with TURP, with 82% of patients in the PUL arm achieving the recovery endpoint by 1 month compared with 53% of patients in the TURP arm (P = 0.008). The results demonstrate that both the PUL and TURP procedures offered significant improvement in symptoms, Q_{max} and HRQoL. The modest patient number may not have provided sufficient statistical power to detect differences in some of the secondary outcome variables.~~

Transurethral Waterjet Ablation (Aquablation)

~~Initial treatment for BPH is usually medical therapy, but this often provides only modest relief. Up to 30% of patients require surgical intervention. Aquablation is performed by a medical device that allows rapid removal of prostate tissue without leaving a zone of thermal damage on the treated tissue. It utilizes a waterjet for automated tissue resection as well as for optical energy delivery for cauterization in the treatment of BPH (Hayes, 2018).~~

~~Kaplan-Marans et al. (2022) searched the FDA MAUDE database for reports of device malfunction and procedural complications for novel treatments for BPH. The results showed a total of 391 adverse events reported, 102 were related to aquablation, and 79 were severe or life threatening. There were 4 cases of rectal perforation, and 31% were cases requiring blood transfusion. The most common device malfunctions (8 instances of each) were motion error, and handpiece attachment fracture/ attachment malfunction. The authors concluded these complications and device related malfunctions are significant, and patients and providers should be aware of the potential risks when choosing this treatment for BPH.~~

UnitedHealthcare, Inc. ("UHC") Proprietary and Confidential Information: The information contained in this document is confidential, proprietary and the sole property of UHC. The recipient of this information agrees not to disclose or use it for any purpose other than to facilitate UHC's compliance with applicable State Medicaid contractual requirements. Any other use or disclosure is strictly prohibited and requires the express written consent of UHC.

Gilling et al. (2022)

Elterman et al. (2021-reviewed in the 2020 ECRI clinical evidence assessment below) conducted a meta-analysis of individual patient data from patients undergoing aquablation treatment for BPH from four prospective global clinical trials; WATER, WATER II, FRANCAIS WATER and OPEN WATER.

~~In a 2021 Hayes technology assessment regarding aquablation for treating benign prostatic hyperplasia, it was concluded that a low-quality body of evidence suggests it may improve LUTS associated with BPH in the short to intermediate term without impacting sexual or function, and without serious safety concerns. However substantial uncertainty remains due to the scarcity of evidence comparing aquablation to TURP, which is the primary surgical modality for treating BPH in small to moderate sized prostates, as well as limited long term evidence. Furthermore, clarity is lacking as to which patient populations are likely to benefit the most from aquablation therapy. The report concludes that this technology has a potential but unproven benefit.~~

~~A 2020 2018 ECRI clinical evidence assessment, updated in 2021, of the Aquabeam Robotic System for treating benign prostatic hyperplasia (BPH), reports this is safe, and reduces BPH-related LUTS for up to three years based on evidence from one randomized controlled trial (RCT) and four prospective case series, and this evidence is somewhat favorable. The RCT reported that Aquablation works as well as TURP for improving LUTS at three-year follow up, but findings need confirmation in additional RCTs. RCTs of sufficient size are needed that compare AquaBeam to other minimally invasive treatments for LUTS due to BPH, as well as report patient-oriented outcomes.~~

~~Gilling et al. (2020) reported the results of participants from the Water I clinical trial to report 3-year outcomes for aquablation compared to TURP for the treatment of LUTS related to BPH. Assessments included IPAA, MSHQ-EjD, IIEF and uroflow. Over 3 years of treatment, improvements in IPSS scores were statistically similar across groups. Mean 3-year improvements were 14.4 and 13.9 points in the Aquablation and TURP groups, respectively (difference of 0.6 points, 95% CI -3.3-2.2, p = .6848). Similarly, 3-year improvements in Qmax were 11.6 and 8.2 cc/sec (difference of 3.3 [95% CI -0.5-7.1] cc/sec, p = .0848). At 3 years, PSA was reduced significantly in both groups by 0.9 and 1.1 ng/mL, respectively; the reduction was similar across groups (p = .5983). There were no surgical retreatments for BPH beyond 20 months for either Aquablation or TURP. It was concluded that three-year BPH symptom reduction and urinary flow rate improvement were similar after TURP and Aquablation therapy. No subjects required surgical retreatment beyond 20 months postoperatively. This study is limited by a maximum prostate size of 80cc (however Desai reported on results with larger prostates below), and whether the rigor of clinical trial data can be applied in real world settings. Furthermore, the study may have been too small to detect clinically significant differences at three years, as it was powered for non-inferiority at six months.~~

~~Desai et al. (2020) reported the 2-year safety and effectiveness of aquablation in men with larger prostate volumes of 80-150cc in a prospective, multicenter international case series (WATER II). Participants had a mean prostate volume of 107 cc and the results showed IPSS and IPSS quality of life improved from 23.2 to 1.1, and 4.6 to 1.1 from baseline to 2 years respectively. Maximum urinary flow increased from 8.7 to 18.2 cc/sec. By the end of the 2-year study timeframe, all but 2 of the 74 participants stopped taking alpha blockers and all but 32 stopped taking 5 α -reductase inhibitors. During the 2-year~~

~~study time frame, adverse urological events were low and included 2 subjects with recurrent BPH symptoms that required retreatment with TURP and HOLEP. The authors concluded that the aquablation procedure is a safe and effective treatment for men with LUTS due to BPH with larger prostate volumes and has an acceptable safety profile and a low retreatment rate. This trial is limited by a lack of a control group which prevented direct comparison to other treatments.~~

~~Bach et al. (2020) conducted an international prospective, multicenter, single-arm, open-label, international clinical trial of the efficacy of the Aquablation procedure for the treatment of lower urinary tract symptoms (LUTS) due to BPH in 177 men enrolled at five treatment centers between September 2017 and December 2018. The primary endpoint was the change in total IPSS from baseline to 3 months. Secondary endpoints included the following: (1) Proportion of subjects who were sexually active at the baseline and experienced either ejaculatory or erectile dysfunction at 3 months, change from the baseline to 3 months in maximal flow rate (Qmax), prostate specific antigen (PSA) level, post void residual (PVR), total MSHQ score, and selected IIEF 5 score. The degree of dysuria was collected on a 0 (not at all) to 5 (almost always) scale. Inclusion criteria was a diagnosis of LUTS due to BPH and a prostate size between 20 and 150 cc. Men were excluded if they were unable to stop anticoagulants and antiplatelet agents perioperatively or had a bleeding disorder, had a history of gross hematuria, were using systemic immune suppressants, had a contraindication to both general and spinal anesthesia, were unwilling to accept transfusion if required, or had any severe illness that could prevent complete follow-up. Patients with prior BPH surgery were not excluded. At baseline and 3 and 12 month follow up, participants completed the International Prostate Symptom Score (IPSS), Incontinence Severity Index, Pain Intensity Scale, Quality of Recovery Visual Analog Scale, International Index of Erectile Function (IIEF-15), the Male Sexual Health Questionnaire (MSHQ-EjD), uroflowmetry and post void residual volume (PVR) measurements. The results showed of the original 177 participants enrolled and had the procedure completed, by month 12, 30 were lost to follow up, three voluntarily withdrew, and one died of an unrelated cause. Mean IPSS improved from 21.7 (7.1) at baseline to 7.1 (5.8) at 3-month follow-up, and 6.4 (4.8) at 12-month follow-up. IPSS QOL scores improved from 4.7 (1.1) at baseline to 1.5 (1.4) at 3-month follow-up, and 1.4 (1.4) at 12-month follow-up. IPSS storage and voiding scales also improved significantly ($p < 0.0001$) at 3 and 12 months (Baseline IPSS scores were unavailable in nine men; of these, seven were using a urinary catheter at the baseline and two cases had incomplete questionnaire responses). Maximum urinary flow rate increased from 9.9 (5.3) cc/sec at baseline to 20.3 (11.4) cc/sec at month 3 and 20.8 (11.2) cc/s at month 12. Postvoid residual improved from 108 (108) to 47 (77) cc at three months and 61 (74) cc at 12 months. Of the 92 men that were sexually active at baseline and 12 months, the MSHQ-EjD score changed by -1 at 3 months, and -1.1 points at 12 months. MSHQ bother/satisfaction changed by -0.3 and -0.7 points at 3 and 12 months respectively. IIEF-15 scores remained stable through month 3. 141 patients had transrectal ultrasound at baseline and after 3 months which showed a decrease in prostate size of 36%. Leakage of urine was reported by 68% of participants at baseline and had reduced to 55% at 12 months, and ISI improved non-significantly. Dysuria of any frequency was reported by 51% at baseline and 29% at 3-month follow-up, and associated pain decreased from 3.5 to 2.4. General pelvic pain decreased from 1.3 at baseline to 0.4 at 3 month follow up. 82 of the participants were taking medication for BPH preoperatively and by month 3, all but 8 had discontinued the medication. There were 69 adverse events reported in 56 participants; 33 grade 1 events, 15 grade 2 events, five grade 3a events and 16 grade 3b events. The authors concluded that Aquablation is safe and effective for men with LUTS due to BPH and replicate results previously seen in a trial setting. This study is limited by a lack of a concurrent control group and a relatively short-term efficacy follow-up.~~

A 2019 Cochrane review on Aquablation (Hwang et al.) identified only one RCT, the Gilling study described below. The authors concluded that based on short-term (up to 12 months) follow-up, the effect of Aquablation on urological symptoms is probably similar to that of TURP (moderate-certainty evidence). The effect on quality of life may also be similar (low-certainty evidence). There is uncertainty whether patients undergoing Aquablation are at higher or lower risk for major adverse events (very low-certainty evidence). Aquablation may result in little to no difference in erectile function but offer a small improvement in preservation of ejaculatory function (both very low certainty evidence). These conclusions are based on a single study of men with a prostate volume up to 80 mL in size. Longer-term data and comparisons with other modalities appear critical to a more thorough assessment of the role of Aquablation for the treatment of LUTS in men with BPH.

Gilling et al. (2019) compared 2-year safety and efficacy outcomes after Aquablation or TURP for the treatment of LUTS related to BPH. A total of 181 patients with BPH were randomly assigned (2:1 ratio) to either Aquablation or TURP. Patients and follow up assessors were blinded to treatment. Assessments included the IPSS, MSHQ, IIEF and uroflow. At 2 years, IPSS scores improved by 14.7 points in the Aquablation group and 14.9 points in TURP ($p = 0.8304$, 95 % CI: - 2.1 to 2.6 points). Two-year improvements in Qmax were 11.2 and 8.6 cc/s for Aquablation and TURP, respectively ($p = 0.1880$, 95 % CI: - 1.3 to 6.4). Sexual function as assessed by MSHQ was stable in the Aquablation group and decreased slightly in the TURP group. At 2 years, PSA was reduced in both groups by 0.7 and 1.2 points, respectively; the reduction was similar across groups ($p = 0.1816$). Surgical re-treatment rates after 12 months for Aquablation were 1.7 % and 0 % for TURP. Over 2 years, surgical BPH retreatment rates were 4.3 % and 1.5 % ($p = 0.4219$), respectively. The authors concluded that 2-year efficacy outcomes after TURP and Aquablation were similar, and the rate of surgical re-treatment was low and similar to TURP; Aquablation may be an alternative for men who strongly prefer maintenance of ejaculatory function. The sample size may however have been too small to detect clinically important differences.

Reale et al. (2019) performed a systematic review or case series and comparison studies, to evaluate functional outcomes (Qmax, QoL, IPSS, PVR), sexual outcome (erectile dysfunction and anejaculation rate), and adverse events evaluated according to the Clavien-Dindo classification. The functional outcomes, evaluated after water jet dissection, have shown improvement with respect to the baseline in all the selected articles. In the comparison papers with the TURP, the Aquablation has been statistically not inferior regarding functional outcomes. The sexual outcomes have highlighted a better ejaculation rate for water jet dissection than TURP. Regarding the adverse events, water jet dissection documented low rates of adverse events and, in comparison studies, were not statistically superior to TURP. Multicenter randomized trials with larger cohorts and longer follow-up are still needed.

A study to compare urodynamic outcomes between aquablation vs transurethral resection of the prostate (TURP) was performed (Pimentel et al., 2019). Patients (n=66) were randomized 2:1 (aquablation: TURP) in the Waterjet Ablation Therapy for Endoscopic Resection of prostate tissue study. Urodynamics were measured at baseline and 6 months. At mean baseline pDet@qmax was 71 and 73cm H2O in the aquablation and TURP groups, respectively. At 6-month follow-up, pDet@qmax decreased by 35 and 34cm H2O, respectively. A large negative shift in bladder outlet obstruction index was observed, consistent with a large reduction in the proportion of subjects with obstruction at follow-up compared to baseline (79% to 22% in aquablation and 96% to 22% in TURP). The authors concluded that

~~in this trial, improvements after aquablation in objective measures of bladder outlet obstruction were similar to those observed after TURP.~~

~~Plante et al. (2018) conducted prespecified post hoc exploratory subgroup analyses from a double-blind, multicenter prospective randomized controlled trial that compared transurethral resection of the prostate (TURP) using either standard electrocautery vs surgery using robotic waterjet (aquablation) to determine whether certain baseline factors predicted more marked responses after aquablation as compared with TURP. The primary efficacy endpoint was reduction in International Prostate Symptom Score (IPSS) at 6 months. The primary safety endpoint was the occurrence of Clavien-Dindo persistent grade 1 or grade ≥ 2 surgical complications. For men with larger prostates (50–80 g), the mean IPSS reduction was four points greater after aquablation than after TURP, a larger difference than the overall result. The primary safety endpoint difference was greater for men with large prostate compared with the overall result. Postoperative anejaculation was also less common after aquablation compared with TURP in sexually active men with large prostates vs the overall results. Exploratory analysis showed larger IPSS changes after aquablation in men with enlarged middle lobes, men with severe middle lobe obstruction, men with a low baseline maximum urinary flow rate, and men with elevated post-void residual urine volume. The authors concluded that in men with moderate-to-severe lower urinary tract symptoms attributable to BPH and larger, more complex prostates, aquablation was associated with both superior symptom score improvements and a superior safety profile, with a significantly lower rate of postoperative anejaculation. The authors noted that the standardized, robotically executed, surgical approach with aquablation may overcome the increased outcome variability in more complex anatomy, resulting in superior symptom score reduction. The RCT reported short term outcomes and included patients with a prostate size 30 to 80 cc. Therefore, results may not be generalizable for all prostate sizes.~~

~~Gilling et al. (2018) conducted a double-blind, multicenter, prospective, randomized, controlled trial (WATER I) to compare safety and efficacy of Aquablation and transurethral prostate resection for the treatment of lower urinary tract symptoms related to benign prostatic hyperplasia. One hundred and eighty-one patients with moderate to severe lower urinary tract symptoms related to benign prostatic hyperplasia underwent transurethral prostate resection or Aquablation. The primary efficacy end point was the reduction in International Prostate Symptom Score at 6 months. The primary safety end point was the development of Clavien-Dindo persistent grade 1, or 2 or higher operative complications. The results showed the mean total operative time was similar for Aquablation and transurethral prostate resection, but resection time was lower for Aquablation. At month 6 patients treated with Aquablation and transurethral prostate resection experienced large IPSS improvements. The prespecified study noninferiority hypothesis was satisfied. Of the patients who underwent Aquablation and transurethral prostate resection 26% and 42%, respectively, experienced a primary safety end point, which met the study primary noninferiority safety hypothesis and subsequently demonstrated superiority. Among sexually active men the rate of anejaculation was lower in those treated with Aquablation (10% vs 36%) The authors concluded that surgical prostate resection using Aquablation showed noninferior symptom relief compared to transurethral prostate resection but with a lower risk of sexual dysfunction. Larger prostates (50 to 80 ml) demonstrated a more pronounced superior safety and efficacy benefit. Longer term follow-up would help assess the clinical value of Aquablation. This study was supported by PROCEPT Bio Robotics, the manufacturer of the AquaBeam® device. Several of the authors indicate a financial interest and/or other relationship with PROCEPT BioRobotics. These conflicts of interest may limit the conclusions that can be drawn from the study.~~

Gilling et al. (2017) performed a prospective, single arm, multicenter trial at a total of 3 centers in Australia and New Zealand with 1-year follow-up to establish the safety and effectiveness of aquablation, an image guided, robotic assisted, water jet tissue ablation technology, for the treatment of benign prostatic hyperplasia. A total of 21 men with moderate to severe lower urinary tract symptoms (LUTS) were included in the study with in-clinic follow up visits at 1, 3, 6 and 12 months. The visits included a review of AEs, uroflow measurements prostate specific antigen (PSA) measurement (at 6 and 12 months only), completion of study questionnaires, and (at 6 months only) urodynamics and transrectal ultrasound (TRUS). Symptoms related to LUTS had significantly improved from baseline at 1 month and were sustained through month 12. At 12 months, the mean international prostatic symptom score (I-PSS) score had improved by 16.2 points. The I-PSS QOL component improved by 3.3 points. Mean maximum urinary flow improved from 8.7 ml per second at baseline to 18.3 ml per second and post-void residual volume (PVR) improved from 136 to 54 ml. Prostate volume decreased from 57 ml at baseline to 35 ml. The bladder outlet obstruction index decreased from 48 at baseline to 13 at month 6. Mean serum PSA, which was measured in 20 subjects, showed no significant change from 3.15 ng/ml at baseline to 2.56 ng/ml at 12 months. No urinary incontinence developed, and sexual function was preserved postoperatively. The authors concluded that this study provides early evidence to support the safety and effectiveness of aquablation for symptomatic benign prostatic hyperplasia by improved symptom scores and other measures of obstruction. The study is of small sample size and lacks a concurrent control group.

Temporary Urethral Stents

Temporary urethral stents are used to maintain urine flow and for short-term use; they are commonly used in males with BPH. These temporary devices can be either removable or absorbable.

Ahn et al. (2020) retrospectively investigated the clinical effectiveness between two temporary urethral stents in the treatment of traumatic bulbar urethral strictures. 30 patients diagnosed with complete bulbar urethral rupture following blunt trauma underwent temporary urethral stent placement. Fifteen patients were treated with a thermo-expandable nickel-titanium alloy urethral stent (Memokath) and the other fifteen with the Allium Bulbar Urethral Stent (BUS). After placement, all stents were removed at 6 months with participant follow up at 1, 3, 6 and 12 months. The follow-up visits included patient assessment with uroflowmetry and ureteroscopy. While the BUS had a lower incidence of stent-related complications than Memokaths, the authors concluded both stents were effective for managing traumatic complete bulbar urethral rupture. This review is limited by lack of randomization, lack of comparison group undergoing traditional open urethroplasty, small sample size and short duration of follow-up; further investigation is warranted.

Chughai et al. (2020) conducted a RCT that compared a temporarily implanted nitinol device (iTind; aka ITIND or Tind) to that of a sham on 175 males with lower urinary tract symptoms due to benign prostatic hyperplasia (BPH). Inclusion criteria for the participants were males 50 years of age or older, an International Prostate Symptoms Score (IPSS) of ≥ 10 , peak urinary flow rate (PFR) of ≤ 12 mL/sec with a 125 mL voided volume, and prostate volume between 25 and 75 cc. Subjects were randomized into either insertion of the iTIND or a sham control group; the sham group received the insertion of a foley catheter to simulate both implantation and retrieval of a temporary implanted device. The a priori primary outcome was changes in IPSS score at three months post procedure. In the intention to treat patient population, the iTind arm improved IPSS by -9.0 ± 8.5 (22.1-13.0) while the sham arm improved -6.6 ± 9.5 (22.8-15.8) ($p = 0.063$) at 3

months. A total of 78.6% of patients in the iTind arm showed a reduction of ≥ 3 points in IPSS, vs 60% of patients in the control arm at 3 months ($p = .029$). Adverse events occurred in 38.1% of patients in the iTind arm and 17.5% in the control arm. The study failed to identify significant differences between groups in peak urinary flow rate, quality of life, or sexual function. The authors found iTIND to be durable for twelve months with only 4.7% of participants having undergone another surgical intervention for BPH. 78.6% of the patients receiving the iTIND had improvement of their IPSS score. Limitations included mixed results, loss to follow-up of almost 30% of participants, and specific inclusion criteria that could or could not be applied to all males with BPH.

Porpiglia et al. (2018) reported 3-year outcomes from a prospective case series study involving the temporary implantable nitinol device (iTIND) implantation for the treatment of BPH. Thirty-two patients with LUTS were enrolled. Follow-up assessments were made at 3 and 6 weeks, and 3, 6, 12, 24 and 36 months after the implantation. The change from baseline in IPSS, QoL score and Qmax was significant at every follow-up time point. After 36 months of follow up, a 41% rise in Qmax was achieved (mean 10.1 mL/s), the median (IQR) IPSS was 12 (6-24) and the IPSS QoL was 2 (1-4). Four early complications (12.5%) were recorded, including one case of urinary retention (3.1%), one case of transient incontinence due to device displacement (3.1%), and two cases of infection (6.2%). No further complications were recorded during the 36-month follow-up. In the authors' opinion, the extended follow up period supports the temporary stent to be safe, effective, and well-tolerated. Lack of comparison group or randomization and small patient population are limitations to this study.

Kimata et al. (2015) conducted a small prospective case series ($n=37$ elderly male patients) to evaluate the use of the Memokath in patients who required long-term urination management with Foley catheters. Patients were followed for a mean of approximately 33 months. A total of 21 patients (56.7%) were able to urinate without assistance after insertion of the Memokath stent. This study was hampered by several limitations, including lack of randomization and appropriate control group.

Kim et al. (2014) conducted a small cohort study ($n=27$) to compare patients who received treatment with a Memokath stent and a self-expandable covered metallic stent (UVENTA) for managing ureteral obstructions. Study results showed no significant differences between the two types of stents for benign and malignant ureteral obstructions. However, the clinical success rate was higher for the UVENTA stent (82.4%) compared with the Memokath stent (42.9%) ($P=0.031$). Patients who received the Memokath stent experienced tumor progression ($n=2$), stent migration ($n=6$), flank pain ($n=1$), and acute pyelonephritis ($n=1$). The study is limited by lack of comparison with other established treatments or randomization between the two stent types.

Jordan et al. (2013) investigated the ability of the Memokath[®] 044TW stent to maintain urethral patency after dilation or internal urethrotomy for recurrent urethral stricture. A total of 92 patients with recurrent bulbar urethral strictures were treated with dilation or internal urethrotomy and randomized to short-term urethral catheter diversion ($n=29$) or insertion of a Memokath 044TW stent ($n=63$). The primary end point was urethral patency, as assessed by passage of a calibrated endoscope. Secondary end points included urinary symptoms and uroflowmetry parameters. Stents were scheduled to remain in situ for 12 months. The rate of successful stent insertion was 93.6%. In stented patients, patency was maintained significantly longer than controls (median 292 vs 84 days). Patency was reflected in significantly improved uroflowmetry and symptom scores. The stent was removed in 100% of patients. The most frequently noted side effects in stented patients were bacteriuria, hematuria and penile pain, which were usually mild and transient. Stent

UnitedHealthcare, Inc. ("UHC") Proprietary and Confidential Information: The information contained in this document is confidential, proprietary and the sole property of UHC. The recipient of this information agrees not to disclose or use it for any purpose other than to facilitate UHC's compliance with applicable State Medicaid contractual requirements. Any other use or disclosure is strictly prohibited and requires the express written consent of UHC.

dislocation and occlusion were observed in 8 and 3 patients, respectively. The authors concluded that patients with recurrent bulbar urethral strictures treated with dilation or urethrotomy and a Memokath 044TW stent maintained urethral patency significantly longer than those treated with dilation or urethrotomy alone. The Memokath stent is not FDA approved and should be considered investigational. Based on differences in stent design, these findings may not be generalizable to FDA-approved devices.

Goh et al. (2013) assessed the ease of insertion and removal of a temporary prostatic stent (the Spanner) following the use of a prostatic urethral measuring device (the Surveyor[™]) in patients with bladder outflow obstruction or urinary retention awaiting definitive surgery. 16 patients had the Spanner inserted following use of the Surveyor. All insertions were uncomplicated. No symptomatic infection was reported. The stents stayed in situ for a median of 10 days. 12 stents were removed prematurely due to severe symptoms or retention. A total of 12 stents had to be removed endoscopically. The authors concluded that the Spanner is easy to insert. Stent removal via the retrieval suture has been difficult necessitating the use of endoscopy in the majority of cases. Possible causes of stent failure include underestimation of the prostatic urethral length by the Surveyor leading to obstruction by apical prostatic tissue, excessive suture length between the stent and distal anchor permitting proximal migration or inadequate suture length leading to urinary incontinence. According to the authors, further design modifications are suggested.

Following transurethral microwave thermotherapy, 186 patients were randomized to receive a Spanner (n=100) or the standard of care (n=86). The stent group reported significantly superior improvement in symptoms at the one week follow up visit. Thereafter, there was no significant difference between the stent and control groups. The investigators concluded that the Spanner is a safe, effective and well tolerated temporary stent for severe prostatic obstruction resulting from therapy induced edema after transurethral microwave thermotherapy (Dineen et al., 2008). Shore et al. published the same study in 2007. The study results are limited in demonstrating meaningful improvement in clinical outcomes in the group that received the temporary prostatic stent compared to the patients in the control group.

Egilmez et al. (2006) evaluated the efficacy of intraurethral metal stents in preventing or eradicating urinary tract infections (UTI) during the management of bladder outlet obstruction (BOO) by comparing the frequency and nature of the infections with indwelling catheter-associated UTI. The SAS relative-risk test was used to compare the risks of UTI in 76 patients with temporary urethral stents, 60 patients with BOO who had never been catheterized nor stented, and 34 patients with a permanent indwelling urethral catheter (PIUC). Infection was assessed 1 month after placement of the devices. After insertion of the catheter, UTI developed in 79.4% of the patients who originally had sterile urine. However, after insertion of the stent, UTI developed in only 40.9% of the patients with sterile urine. In 21 (44.6%) of the catheterized patients who had infected urine, UTI was eradicated after stent insertion. The investigators concluded that urinary infection is a significant problem in patients with PIUC but is significantly less frequent and less severe in patients with urethral stents. These findings require confirmation in larger randomized controlled trials.

A series of 43 consecutive patients were stented with the Spanner temporary prostatic stent and reviewed retrospectively. Stents were removed and replaced every 3 months if tolerated. More than half of the patients (63%) had an unsatisfactory outcome, namely, immediate or delayed retention or elective removal because of unbearable symptoms. The remaining 37% of patients had a satisfactory outcome and either continued to have the

UnitedHealthcare, Inc. ("UHC") Proprietary and Confidential Information: The information contained in this document is confidential, proprietary and the sole property of UHC. The recipient of this information agrees not to disclose or use it for any purpose other than to facilitate UHC's compliance with applicable State Medicaid contractual requirements. Any other use or disclosure is strictly prohibited and requires the express written consent of UHC.

~~stent in situ after a mean of five changes or are stent free after a successful voiding trial (Grimsley et al., 2007).~~

~~The National Institute for Health and Care Excellence (NICE) 2018 medical technology guidance on use of the Memokath-051 stent for ureteric obstruction concludes that the quality of reporting across all the studies was generally poor. None of the studies provided adequate details on patient characteristics, stent insertion procedures, follow-up, statistical analyses and uncertainty around the results. Migration rates and clinical success were the most commonly reported outcomes, but definitions of clinical success varied, so statistical pooling could not be done.~~

~~For information on current and completed trials studying the use of temporary prostatic urethral stents refer to [ClinicalTrials.gov](https://clinicaltrials.gov). (Accessed May 5, 2021)~~

Transperineal Focal Laser Ablation

Standard treatments for prostate cancer such as surgery and radiation involve the whole gland, even if the tumor is small and localized. These treatment modalities are associated with significant urinary and sexual dysfunction. Focal laser ablation (FLA) has been proposed as an alternative, as it allows the treatment of only the tumor, sparing the rest of the gland. The quality of the evidence is however insufficient to support the efficacy and safety of this technology.

Bates et al. (2021) conducted a systematic review (SR) to compare the clinical effectiveness of primary focal ablative therapy (FT) to standard current treatment options for clinically localized prostate cancer (PCa) to make clinical practice recommendations, and identify gaps, providing recommendations for further research. Four primary studies (1 randomized controlled trial [RCT] and 3 retrospective studies) including 3,961 patients, (and ten eligible SRs were identified) reporting on different types of FT. The results showed the following: The RCT compared photodynamic therapy (PDT) with active surveillance and found PDT was associated with a significantly lower rate of treatment failure at 2 years, no difference in functional outcomes, and was associated with worse transient adverse events. A retrospective matched-pair study comparing focal high-intensity focused ultrasound (HIFU) with robotic radical prostatectomy (RP) found no significant differences in treatment failure at 3 years, while the focal HIFU group had better recovery of continence and erectile function. Two retrospective SEER-based, propensity-matched cohort studies compared focal laser ablation (FLA) against radical prostatectomy (RP) and external beam radiotherapy (EBRT), reporting significantly worse overall survival with FLA on adjusted analysis. Overall, the evidence in support of FT as an alternative to either AS or radical interventions for localized PCa is was limited. Data regarding the oncological effectiveness were mixed and inconsistent. For FLA specifically, limited quality data suggest harm, as compared to alternative, established therapies. Overall, for FT, the vast majority of primary studies were small and uncontrolled; others were comparative studies with serious methodological flaws with extremely low internal and external validity. Most studies had significant clinical heterogeneity, with poorly defined populations, interventions (e.g., intermingling of whole-gland and FT as a single index intervention), different definitions of retreatments with different intervals, different imaging and follow-up schedules, different comparators, outcome measures with different definitions of treatment failure measured at different time points, and a lack of long-term data. The overview of SRs confirmed these findings, and none showed high-certainty evidence. The authors concluded that the routine use of FT in clinical practice is currently not recommended and should ideally be restricted to a clinical trial or prospective

comparative study involving comprehensive data capture using standardized definitions and appropriate outcome measures.

In a 2019 Delphi consensus project following a systematic review of the literature, van Luijckelaar et al. presented the evidence-based consensus of 37 international experts in the field of focal therapy for prostate cancer (PCa). Consensus was agreed upon in 39/43 topics. Clinically significant PCa (csPCa) was defined as any volume Grade Group 2 [Gleason score (GS) 3+4]. Focal therapy was specified as treatment of all csPCa and can be considered primary treatment as an alternative to radical treatment in carefully selected patients. In patients with intermediate-risk PCa (GS 3+4) as well as patients with MRI-visible and biopsy-confirmed local recurrence, the experts felt that FLA is optimal for targeted ablation of a specific magnetic resonance imaging (MRI)-visible focus. However, FLA should not be applied to candidates for active surveillance and close follow-up is required. Suitability for FLA is based on tumor volume, location to vital structures, GS, MRI-visibility, and biopsy confirmation. The expert consensus concluded that FLA is a promising technique for treatment of clinically localized PCa and should ideally be performed within approved clinical trials. They noted that there are only a few studies have reported on FLA and further validation with longer follow-up is mandatory before widespread clinical implementation is justified.

Valerio et al. (2017) completed a systematic review summarizing the evidence regarding the specific sources of energy used in focal ablative therapy for prostate cancer. Thirty-seven articles reporting on 3230 patients undergoing focal therapy were selected. Thirteen reported on high-intensity focused ultrasound, 11 on cryotherapy, three on photodynamic therapy, four on laser interstitial thermotherapy, two on brachytherapy, three on irreversible electroporation, and one on radiofrequency. Laser interstitial thermotherapy has been evaluated in up to Stage 2a studies. Median follow-up varied between 4 months and 61 months, and the median rate of serious adverse events ranged between 0% and 10.6%. Padfree leak-free continence and potency were obtained in 83.3-100% and 81.5-100%, respectively. In series with intention to treat, the median rate of significant and insignificant disease at control biopsy varied between 0% and 13.4% and 5.1% and 45.9%, respectively. The authors concluded that while focal therapy seems to have a minor impact on quality of life and genito-urinary function, the oncological effectiveness has not been defined against the current standard of care. The author identified limitations of this SR include the length of follow-up, the absence of a comparator arm, and study heterogeneity.

Transperineal laser ablation (TPLA)

Transperineal laser ablation (TPLA) is a minimally invasive procedure that focuses on lower urinary tract symptoms (LUTS) in patients with BPH. Currently there is insufficient evidence regarding the long-term effectiveness and safety for the use of TPLA; additional well designed RCTs and comparative analyses are warranted.

An ECRI clinical evidence assessment focused on TPLA's safety and effectiveness and compared it to TURP and other minimally invasive BPH treatments (2022). The report included 4 prospective and 2 retrospective before and after studies. The 4 prospective studies compared patients with BPH before and after undergoing TPLA. The results reported on hospital length of stay (LOS), catheterization duration, medication usage, symptoms and QOL (measured on the International Prostate Symptom Score (IPSS)), sexual health, and adverse effects (AEs); the data was measured at 1-, 3-, 6-, and/or 12-month follow-up. The single center ~~retrospective~~retrospective study containedincluded 20 participants with BPH and also reported symptoms before and after undergoing TPLA. Data measured included patient reported symptoms, QOL and AEs at 6-month follow-up. A multicenter

UnitedHealthcare, Inc. ("UHC") Proprietary and Confidential Information: The information contained in this document is confidential, proprietary and the sole property of UHC. The recipient of this information agrees not to disclose or use it for any purpose other than to facilitate UHC's compliance with applicable State Medicaid contractual requirements. Any other use or disclosure is strictly prohibited and requires the express written consent of UHC.

before and after study of 160 participants measured hospital LOS, catheterization duration, QOL and AEs at 6- and 12-month follow-up. The results appear to show TPLA as promising, safe and effective. However, limitations included small sample sizes, no comparative studies and a high risk of bias due to two or more of the following: retrospective design, single-center focus, and lack of control groups and randomization. Further large, multicenter RCTs are needed to validate the studies' findings and to compare TPLA with other treatments. The overall conclusion of the report is that the evidence is inconclusive.

De Reinzo et al. (2021), conducted a case series which prospectively analyzed patients that underwent TPLA of the prostate. Inclusion criteria were presence of moderate to severe lower urinary tract symptoms (LUTS) due to benign prostatic obstruction (BPO) with an International Prostate Symptom Score (IPSS) > 12, aged 40 - 90 years, with prostate volume up to 100 ml and lack of efficacy, and intolerance or poor compliance to medical therapies. Twenty-one patients were included in this study sixteen of which had received at least one oral medication and five had refused medical treatment. Patients were discharged within 24 hours of procedure and kept the transurethral catheter for 8.7 + 2.5d. The authors found all but one patient was able to discontinue medical therapy within one month showing improvement in Qmax and IPSS scores which continued to improve at 6 months. Ejaculatory function was preserved and only one patient complication (prostatic abscess) was noted. Limitations included the absence of a control group, small sample size, and heterogeneity among patients with previous medical therapy.

Pacella et al. (2020, included in the ECRI report cite above) conducted a multicenter retrospective study case series to assess the effectiveness and safety of TPLA in the treatment of patients with symptomatic BPH. The primary endpoints were evaluation of IPSS, QoL, Qmax, PVR, and prostatic volume at 6- and 12-months post TPLA treatment with secondary end-point evaluation of complications. Prior to treatment, patients completed the IPSS and QoL questionnaires, completed urodynamics to evaluate Qmax, had an abdominal ultrasound to determine post voiding residual (PVR), and transrectal ultrasound to evaluate volume of the prostate. There were 160 patients included with a follow-up of at least 6 months and 83 patients with a follow-up of at least 12 months. At both 6 and 12 months, the authors noted improvement of IPSS, PVR, Qmax, QoL and volume. Out of 160 cases, there were 7 grade I complications including hematuria, acute urinary retention and one orchitis, along with 1 grade III complication which was a prostatic abscess. Limitations included retrospective analysis with no other treatment comparison, treatment in separate centers with potential effect to the results related to operator experience with a novel technique, only a select subgroup of patients with minimum of 6 months of follow-up data were included, and ejaculatory function was not evaluated in the questionnaire and might be underestimated in this study.

Clinical trials for TPLA are currently ongoing. See the following website for more information: <https://clinicaltrials.gov/ct2/home> (Accessed May 20, 2022).

Temporary Urethral Stents

Temporary urethral stents are used to maintain urine flow and for short-term use; they are commonly used in males with BPH. These temporary devices can be either removable or absorbable. The quality of the evidence is however insufficient to support the efficacy and safety of this technology.

A Hayes (2022) evolving evidence report identifies limited evidence and weak support for the use of the iTind system; it is currently under investigation for its use in treatment of lower urinary tract symptoms for BPH.

UnitedHealthcare, Inc. ("UHC") Proprietary and Confidential Information: The information contained in this document is confidential, proprietary and the sole property of UHC. The recipient of this information agrees not to disclose or use it for any purpose other than to facilitate UHC's compliance with applicable State Medicaid contractual requirements. Any other use or disclosure is strictly prohibited and requires the express written consent of UHC.

Ahn et al. (2020) retrospectively investigated the clinical effectiveness between two temporary urethral stents in the treatment of traumatic bulbar urethral strictures. 30 patients diagnosed with complete bulbar urethral rupture following blunt trauma underwent temporary urethral stent placement. Fifteen patients were treated with a thermo-expandable nickel-titanium alloy urethral stent (Memokath) and the other fifteen with the Allium Bulbar Urethral Stent (BUS). After placement, all stents were removed at 6 months with participant follow up at 1, 3, 6 and 12 months. The follow-up visits included patient assessment with uroflowmetry and ureteroscopy. While the BUS had a lower incidence of stent-related complications than Memokaths, the authors concluded both stents were effective for managing traumatic complete bulbar urethral rupture. This review is limited by lack of randomization, lack of comparison group undergoing traditional open urethroplasty, small sample size and short duration of follow-up; further investigation is warranted.

Chughai et al. (2020) conducted a RCT that compared a temporarily implanted nitinol device (iTind; aka ITIND or Tind) to that of a sham on 175 males with lower urinary tract symptoms due to benign prostatic hyperplasia (BPH). Inclusion criteria for the participants were males 50 years of age or older, an International Prostate Symptoms Score (IPSS) of ≥ 10 , peak urinary flow rate (PFR) of ≤ 12 mL/sec with a 125 mL voided volume, and prostate volume between 25 and 75 cc. Subjects were randomized into either insertion of the iTIND or a sham control group; the sham group received the insertion of a foley catheter to simulate both implantation and retrieval of a temporary implanted device. The a priori primary outcome was changes in IPSS score at three months post procedure. In the intention to treat patient population, the iTind arm improved IPSS by -9.0 ± 8.5 (22.1-13.0) while the sham arm improved -6.6 ± 9.5 (22.8-15.8) ($p = 0.063$) at 3 months. A total of 78.6% of patients in the iTind arm showed a reduction of ≥ 3 points in IPSS, vs 60% of patients in the control arm at 3 months ($p = .029$). Adverse events occurred in 38.1% of patients in the iTind arm and 17.5% in the control arm. The study failed to identify significant differences between groups in peak urinary flow rate, quality of life, or sexual function. The authors found iTIND to be durable for twelve months with only 4.7% of participants having undergone another surgical intervention for BPH. 78.6% of the patients receiving the iTIND had improvement of their IPSS score. Limitations included mixed results, loss to follow-up of almost 30% of participants, and specific inclusion criteria that could or could not be applied to all males with BPH.

Porpiglia et al. (2018) reported 3-year outcomes from a prospective case series study involving the temporary implantable nitinol device (iTIND) implantation for the treatment of BPH. Thirty-two patients with LUTS were enrolled. Follow-up assessments were made at 3 and 6 weeks, and 3, 6, 12, 24 and 36 months after the implantation. The change from baseline in IPSS, QoL score and Qmax was significant at every follow-up time point. After 36 months of follow-up, a 41% rise in Qmax was achieved (mean 10.1 mL/s), the median (IQR) IPSS was 12 (6-24) and the IPSS QoL was 2 (1-4). Four early complications (12.5%) were recorded, including one case of urinary retention (3.1%), one case of transient incontinence due to device displacement (3.1%), and two cases of infection (6.2%). No further complications were recorded during the 36-month follow-up. In the authors' opinion, the extended follow-up period supports the temporary stent to be safe, effective, and well-tolerated. Lack of comparison group or randomization and small patient population are limitations to this study.

Kimata et al. (2015) conducted a small prospective case series (n=37 elderly male patients) to evaluate the use of the Memokath in patients who required long-term urination management with Foley catheters. Patients were followed for a mean of

UnitedHealthcare, Inc. ("UHC") Proprietary and Confidential Information: The information contained in this document is confidential, proprietary and the sole property of UHC. The recipient of this information agrees not to disclose or use it for any purpose other than to facilitate UHC's compliance with applicable State Medicaid contractual requirements. Any other use or disclosure is strictly prohibited and requires the express written consent of UHC.

approximately 33 months. A total of 21 patients (56.7%) were able to urinate without assistance after insertion of the Memokath stent. This study was hampered by several limitations, including lack of randomization and appropriate control group.

Kim et al. (2014) conducted a small cohort study (n=27) to compare patients who received treatment with a Memokath stent and a self-expandable covered metallic stent (UVENTA) for managing ureteral obstructions. Study results showed no significant differences between the two types of stents for benign and malignant ureteral obstructions. However, the clinical success rate was higher for the UVENTA stent (82.4%) compared with the Memokath stent (42.9%) (P=0.031). Patients who received the Memokath stent experienced tumor progression (n=2), stent migration (n=6), flank pain (n=1), and acute pyelonephritis (n=1). The study is limited by lack of comparison with other established treatments or randomization between the two stent types.

Jordan et al. (2013) investigated the ability of the Memokath™ 044TW stent to maintain urethral patency after dilation or internal urethrotomy for recurrent urethral stricture. A total of 92 patients with recurrent bulbar urethral strictures were treated with dilation or internal urethrotomy and randomized to short-term urethral catheter diversion (n=29) or insertion of a Memokath 044TW stent (n=63). The primary end point was urethral patency, as assessed by passage of a calibrated endoscope. Secondary end points included urinary symptoms and uroflowmetry parameters. Stents were scheduled to remain in situ for 12 months. The rate of successful stent insertion was 93.6%. In stented patients, patency was maintained significantly longer than controls (median 292 vs 84 days). Patency was reflected in significantly improved uroflowmetry and symptom scores. The stent was removed in 100% of patients. The most frequently noted side effects in stented patients were bacteriuria, hematuria and penile pain, which were usually mild and transient. Stent dislocation and occlusion were observed in 8 and 3 patients, respectively. The authors concluded that patients with recurrent bulbar urethral strictures treated with dilation or urethrotomy and a Memokath 044TW stent maintained urethral patency significantly longer than those treated with dilation or urethrotomy alone. The Memokath stent is not FDA approved and should be considered investigational. Based on differences in stent design, these findings may not be generalizable to FDA-approved devices.

Goh et al. (2013) assessed the ease of insertion and removal of a temporary prostatic stent (the Spanner) following the use of a prostatic urethral measuring device (the Surveyor™) in patients with bladder outflow obstruction or urinary retention awaiting definitive surgery. 16 patients had the Spanner inserted following use of the Surveyor. All insertions were uncomplicated. No symptomatic infection was reported. The stents stayed in situ for a median of 10 days. 12 stents were removed prematurely due to severe symptoms or retention. A total of 12 stents had to be removed endoscopically. The authors concluded that the Spanner is easy to insert. Stent removal via the retrieval suture has been difficult necessitating the use of endoscopy in the majority of cases. Possible causes of stent failure include underestimation of the prostatic urethral length by the Surveyor leading to obstruction by apical prostatic tissue, excessive suture length between the stent and distal anchor permitting proximal migration or inadequate suture length leading to urinary incontinence. According to the authors, further design modifications are suggested.

Following transurethral microwave thermotherapy, 186 patients were randomized to receive a Spanner (n=100) or the standard of care (n=86). The stent group reported significantly superior improvement in symptoms at the one-week follow-up visit. Thereafter, there was no significant difference between the stent and control groups. The investigators concluded that the Spanner is a safe, effective and well tolerated temporary stent for

UnitedHealthcare, Inc. ("UHC") Proprietary and Confidential Information: The information contained in this document is confidential, proprietary and the sole property of UHC. The recipient of this information agrees not to disclose or use it for any purpose other than to facilitate UHC's compliance with applicable State Medicaid contractual requirements. Any other use or disclosure is strictly prohibited and requires the express written consent of UHC.

severe prostatic obstruction resulting from therapy induced edema after transurethral microwave thermotherapy (Dineen et al., 2008). Shore et al. published the same study in 2007. The study results are limited in demonstrating meaningful improvement in clinical outcomes in the group that received the temporary prostatic stent compared to the patients in the control group.

Egilmez et al. (2006) evaluated the efficacy of intraurethral metal stents in preventing or eradicating urinary-tract infections (UTI) during the management of bladder outlet obstruction (BOO) by comparing the frequency and nature of the infections with indwelling-catheter-associated UTI. The SAS relative-risk test was used to compare the risks of UTI in 76 patients with temporary urethral stents, 60 patients with BOO who had never been catheterized nor stented, and 34 patients with a permanent indwelling urethral catheter (PIUC). Infection was assessed 1 month after placement of the devices. After insertion of the catheter, UTI developed in 79.4% of the patients who originally had sterile urine. However, after insertion of the stent, UTI developed in only 40.9% of the patients with sterile urine. In 21 (44.6%) of the catheterized patients who had infected urine, UTI was eradicated after stent insertion. The investigators concluded that urinary infection is a significant problem in patients with PIUC but is significantly less frequent and less severe in patients with urethral stents. These findings require confirmation in larger randomized controlled trials.

A series of 43 consecutive patients were stented with the Spanner temporary prostatic stent and reviewed retrospectively. Stents were removed and replaced every 3 months if tolerated. More than half of the patients (63%) had an unsatisfactory outcome, namely, immediate or delayed retention or elective removal because of unbearable symptoms. The remaining 37% of patients had a satisfactory outcome and either continued to have the stent in situ after a mean of five changes or are stent free after a successful voiding trial (Grimsley et al., 2020). For information on current and completed trials studying the use of temporary prostatic urethral stents refer to ClinicalTrials.gov. (Accessed May 20, 2022)

Transperineal Placement of Biodegradable Material

The SpaceOAR is used to temporarily position the anterior rectal wall away from the prostate during radiotherapy for prostate cancer and in creating this space it is the intent to reduce the radiation dose delivered to the anterior rectum. The absorbable spacer maintains space for the entire course of prostate radiotherapy treatment and is completely absorbed by the patient's body over time.

A Hayes health technology assessment report (2020/2021) summarized that while published evidence suggests a potential benefit of an absorbable perirectal spacer (APS) during radiation therapy for prostate cancer, there is substantial uncertainty regarding its safety and efficacy; future studies are needed to assess the APS clinical usefulness and cost-effectiveness.

In a custom product brief, ECRI (2020) concludes that SpaceOAR hydrogel is well tolerated and works as intended to reduce rectal irradiation long-term, but not acute, rectal toxicity, and it improves bowel quality of life (QOL), based on one randomized controlled trial and four prospective nonrandomized comparative studies.

Afkhami Ardekani and Chaffari (2020) evaluated the effect of dosimetry and procedure toxicity of polyethylene glycol (PEG)-based hydrogel spacers during prostate brachytherapy. There were twelve studies included in the systematic review involving 615 patients. The approach used to place the hydrogel spacers was hydrodissection and considered one of the most common techniques. Ultrasonography is used to insert a large

gauge needle where saline water is injected to create potential space between the prostate and anterior rectal wall; PEG hydrogel is then injected into the created space. The DuraSeal and SpaceOar then polymerize within 3 and 10 seconds after injection. The authors found the data of several studies revealed the rectal dosimetry was significantly reduced with the use of the PEG hydrogel spacers and that the procedure was safe. The authors concluded the implantation of PEG hydrogel spacers is practical and safe with well tolerance of the procedure. The use of PEG hydrogels for prostate brachytherapy has a very high success rate, however the advantages of these spacers should be weighed against possible risks of complications. Additional RCTs should be done to further clarify rectal dose reduction on toxicity and quality of life.

A systematic review was conducted by Vaggers et al. (2020) from nine full text articles reviewing polyethylene glycol-based hydrogel rectal spacers for prostate brachytherapy. Four studies used the DuraSeal Spinal Sealant and five studies used SpaceOar. Primary outcomes included procedure complications, failures, prostate-rectum separation, rectal dosimetry and GI toxicities for hydrogel insertion. There was little variation in technique used throughout the articles reviewed. The authors found the studies demonstrated a significant reduction in rectal dosimetry and concluded that the polyethylene glycol-based hydrogel rectal spacers appear to be safe and easy. Even though the spaces appear to reduce rectal toxicity, further studies are needed to confirm these findings. Limitations include the review as retrospective and non-randomization along with small sample size.

Paetkau et al. (2019) retrospectively evaluated 13 patients with SpaceOAR implant to determine future planning needs for patients with prostate cancer undergoing radiation therapy. Computerized tomography (CT) scans were taken pre- and post-implant. A prescription of 60 Gy in 20 fractions was planned on both scans. Six treatment plans were produced per anonymized dataset using either a structure of rectum plus the hydrogel, termed composite rectum wall (CRW), or rectal wall (RW) as an inverse optimization structure and intensity modulated radiotherapy (IMRT) or volumetric modulated arc therapy (VMAT) as a treatment technique. Dose-volume histogram metrics were compared between plans to determine which optimization structure and treatment technique offered the maximum rectal dose sparing. RW structures offered a statistically significant decrease in rectal dose over CRW structures, whereas the treatment technique (IMRT vs VMAT) did not significantly affect the rectal dose. There was improvement seen in bladder and penile bulb dose when VMAT was used as a treatment technique. The authors concluded that overall, treatment plans using the RW optimization structure offered the lowest rectal dose while VMAT treatment technique offered the lowest bladder and penile bulb dose.

Wu et al. (2018, included in Afkhami Ardekani and Chaffari (2020) and Vaggers et al. (2020) systematic reviews above) evaluated 18 consecutive patients underwent transperineal ultrasound-guided placement of 10 cc of SpaceOAR hydrogel prior to HDR brachytherapy in the treatment of prostate cancer. Treatment plans were generated using an inverse planning simulated annealing algorithm. Rectal dosimetry for these 18 patients was compared with the 36 preceding patients treated with HDR brachytherapy without SpaceOAR. There was no difference in age, pretreatment prostate-specific antigen, Gleason score, clinical stage, prostate volume, or contoured rectal volume between those who received SpaceOAR and those who did not. Patients who received SpaceOAR hydrogel had significantly lower dose to the rectum as measured by percent of contoured organ at risk (median, V80 < 0.005% vs. 0.010%, $p = 0.003$; V75 < 0.005% vs. 0.14%, $p < 0.0005$; V70 0.09% vs. 0.88%, $p < 0.0005$; V60 = 1.16% vs. 3.08%, $p < 0.0005$); similar results were seen for rectal volume in cubic centimeters. One patient who received SpaceOAR developed a perineal abscess 1 month after treatment. The authors concluded that transperineal

~~insertion of SpaceOAR hydrogel at the time of HDR brachytherapy is feasible and decreases rectal radiation dose. Further investigation is needed with well-designed clinical trials and larger patient populations to further assess the clinical impact.~~

~~Chao et al. (2018) conducted a prospective case series analysis to report on the dosimetric benefits and late toxicity outcomes following injection of a hydrogel spacer between the prostate and rectum in 76 patients with T1-T3a prostate cancer treated with radiotherapy. There were no postoperative complications reported. Mean prostate size were 66.0cc (25.0cc – 187.0cc). Rectal dose volume parameters were observed with volume of rectum receiving 70Gy (rV70), 75Gy (rV75) and 78Gy (rV78) were 7.8%, 3.6% and 0.4%. 21% (16/76) developed acute grade 1 GI toxicities but all were resolved completely by 3 months post treatment. 3% (2/76) developed late grade 1 GI toxicities. No patients experienced acute or late grade 2+ GI toxicities. The authors concluded that injection of hydrogel spacer resulted in a reduction of irradiated rectal dose volumes along with minimal GI toxicities, irrespective of prostate size. Additional studies with longer-term outcomes are needed to evaluate long term toxicities. The findings are limited by lack of comparison group.~~

~~Taggar et al. (2018, included in Afkhami Ardekani and Chaffari (2020) and Vaggers et al. (2020) systematic review above) conducted a prospective cohort study to evaluate placement of an absorbable rectal hydrogel spacer in 74 patients with prostate cancer undergoing low-dose-rate brachytherapy with palladium-103. Rectal dosimetry was compared with a consecutive cohort of 136 patients treated with seed implantation without a spacer. On average, 11.2-mm (SD 3.3) separation was achieved between the prostate and the rectum. The resultant mean rectal volume receiving 100% of prescribed dose (V100%), dose to 1 cc of rectum (D1cc), and dose to 2 cc of rectum (D2cc) were 0 (SD 0.05 cc), 25.3% (SD 12.7), and 20.5% (SD 9.9), respectively. All rectal dosimetric parameters improved significantly for the cohort with spacer placement as compared with the non-spacer cohort. Injection of rectal spacer is feasible in the post-LDR brachytherapy setting and reduces dose to the rectum with minimal toxicity. Prostate and urethral dosimetries do not appear to be affected by the placement of a spacer.~~

~~Pinkawa et al. (2017a) reported 5-year outcomes of a cohort study after prostate cancer radiation therapy with and without the use of a hydrogel spacer. Fifty-four patients were selected to receive a hydrogel spacer. Patients were surveyed before RT; at the last day of RT; and a median time of 2 months, 17 months, and 63 months after RT. For patients treated with a hydrogel spacer, mean bowel function and bother score changes of >5 points in comparison with baseline levels were found only at the end of RT (10–15 points; $P<.01$). No spacer patient reported moderate or big problems with his bowel habits overall. Mean bother score changes of 21 points at the end of RT, 8 points at 2 months, 7 points at 17 months, and 6 points at 63 months after RT were found for patients treated without a spacer. A bowel bother score change >10 points was found in 6% versus 32% ($P<.01$) at 17 months and in 5% versus 14% ($P=.2$) at 63 months with versus without a spacer. The authors conclude that hydrogel spacer application demonstrates excellent treatment tolerability, in particular regarding bowel problems. They encourage further studies with dose-escalated or re-irradiation concepts.~~

~~Pinkawa et al. (2017b) evaluated in a cohort study of 167 consecutive patients who received prostate RT with 2 Gy fractions up to 76 Gy (without hydrogel, $n = 66$) or 76–80 Gy (with hydrogel, $n = 101$). The numbers of interventions resulting from bowel problems during the first 2 years after RT were compared. Patients were surveyed prospectively before RT, at the last day of RT, and at a median of 2 and 17 months after RT using a validated questionnaire (Expanded Prostate Cancer Index Composite). Treatment for bowel~~

symptoms (0 vs. 11%; $p < 0.01$) and endoscopic examinations (3 vs. 19%; $p < 0.01$) were performed less frequently with a spacer. Mean bowel function scores did not change for patients with a spacer in contrast to patients without a spacer (mean decrease of 5 points) >1 year after RT in comparison to baseline, with 0 vs. 12% reporting a new moderate/big problem with passing stools ($p < 0.01$). It was noted that statistically significant differences were found for the items "loose stools", "bloody stools", "painful bowel movements" and "frequency of bowel movements". The authors concluded that spacer injection is associated with a significant benefit for patients after prostate cancer RT.

Hamstra et al. (2017) reported the final outcomes from their single-blind phase III trial of image guided intensity modulated radiation therapy (n=222). The 3-year incidence of grade ≥ 1 (9.2% vs 2.0%; $P=0.028$) and grade ≥ 2 (5.7% vs 0%; $P=0.012$) rectal toxicity favored the spacer arm. Grade ≥ 1 urinary incontinence was also lower in the spacer arm (15% vs 4%; $P=0.046$), with no difference in grade ≥ 2 urinary toxicity (7% vs 7%; $P=0.7$). From 6 months onward, bowel QOL consistently favored the spacer group ($P=0.002$), with the difference at 3 years (5.8 points; $P<0.05$) meeting the threshold for a MID. The control group the authors reported that the benefit of a hydrogel spacer in reducing the rectal dose, toxicity, and QOL declines after image guided intensity modulated radiation therapy for prostate cancer was maintained or increased with a longer follow up period, providing stronger evidence for the benefit of hydrogel spacer use in prostate radiation therapy. Additional long-term outcomes are needed to determine the benefits of hydrogel spacers.

In a prospective, randomized patient blinded clinical study, Karsh et al. (2017) compared image-guided intensity modulated prostate radiotherapy (79.2Gy in 44 fractions) in men with or without insertion of prostate-rectum hydrogel spacer (SpaceOar). The mean additional space created between the prostate and the rectum was just over 1cm, which allowed significant rectum and penile bulb radiation dose reduction resulting in less acute pain, lower rates of late rectal toxicity, and improved bowel and urinary QOL scores from 6 months through the 3-year follow-up period as compared to the control group. The authors concluded that spacer application significantly reduced rectal radiation dose, resulting in long-term reductions in rectal toxicity, as well as improvements in bowel, urinary, and sexual QOL. Patient sample volumes were not reported.

Yeh et al. (2016, included in Afkhami Ardekani and Chaffari (2020) and Vaggers et al. (2020) systematic reviews above) studied rectal toxicity rates in 326 patients administered a polyethylene glycol (PEG) hydrogel rectal spacer in conjunction with combination high-dose-rate brachytherapy at 16 Gy (average dose 15.5 Gy; standard deviation [SD] = 1.6 Gy) and external beam radiotherapy of 59.4 Gy (average dose 60.2 Gy; SD = 2.9 Gy). Clinical efficacy was determined by measuring acute and chronic rectal toxicity using the National Cancer Center Institute Common Terminology Criteria for Adverse Events v4.0 grading scheme. Median follow-up was 16 months. The mean anterior-posterior separation achieved was 1.6 cm (SD = 0.4 cm). Rates of acute Grade 1 and 2 rectal toxicity were 37.4% and 2.8%, respectively. There were no acute Grade 3/4 toxicities. Rates of late Grade 1, 2, and 3 rectal toxicity were 12.7%, 1.4%, and 0.7%, respectively. There were no late Grade 4 toxicities. The authors concluded that acute and chronic rectal toxicities are low despite aggressive dose escalation. Longer term outcomes are needed to evaluate impact.

Mariados et al. (2015) conducted a prospective multicenter randomized controlled pivotal trial to assess outcomes following absorbable spacer (SpaceOAR system) implantation. The study included 222 patients with clinical stage T1 or T2 prostate cancer who underwent computed tomography (CT) and magnetic resonance imaging (MRI) scans for treatment

UnitedHealthcare, Inc. ("UHC") Proprietary and Confidential Information: The information contained in this document is confidential, proprietary and the sole property of UHC. The recipient of this information agrees not to disclose or use it for any purpose other than to facilitate UHC's compliance with applicable State Medicaid contractual requirements. Any other use or disclosure is strictly prohibited and requires the express written consent of UHC.

planning, followed with fiducial marker placement. Patients were randomized to receive spacer injection or no injection (control). Spacer safety and impact on rectal irradiation, toxicity, and QOL were assessed throughout 15 months. Spacer application had a 99% hydrogel placement success rate. The authors reported that there were no device-related AEs, rectal perforations, serious bleeding, or infections within either group. Overall acute rectal adverse event rates were similar between groups, with fewer spacer patients experiencing rectal pain ($P=.02$). There was no late rectal toxicity greater than grade 1 in the spacer group. At 15 months 11.6% and 21.4% of spacer and control patients, respectively, experienced 10-point declines in bowel QOL. MRI scans at 12 months verified spacer absorption. The authors concluded that spacer application was well tolerated. Increased perirectal space reduced rectal irradiation, reduced rectal toxicity severity, and decreased rates of patients experiencing declines in bowel QOL. The spacer appears to be an effective tool, potentially enabling advanced prostate radiation therapy protocols. However, the short follow-up period is a study limitation, as researchers have published the median time to late gastrointestinal grade ≥ 2 toxicity onset was 17 months. The study was also limited by the exclusion of patients with prostate volumes >80 mL, patients with extracapsular extension, and those with prior radiation or surgery. Patients with extracapsular extension have the theoretical risk of pushing posterior extracapsular disease farther from the prostate during radiation therapy, whereas patients with prior radiation or surgery may have perirectal scar formation, limiting space creation. The authors noted that the use of spacers in these populations should proceed cautiously in separate clinical trials.

Transurethral Waterjet Ablation (Aquablation)

Aquablation is performed by a medical device that allows rapid removal of prostate tissue without leaving a zone of thermal damage on the treated tissue. It utilizes a waterjet for automated tissue resection as well as for optical energy delivery for cauterization in the treatment of BPH (Hayes, 2018). The evidence for aquablation is insufficient to demonstrate the efficacy of this technology and safety concerns have emerged.

Kaplan-Marans et al. (2022) searched the FDA MAUDE database for reports of device malfunction and procedural complications for novel treatments for BPH. The results showed a total of 391 adverse events reported, 102 were related to aquablation, and 79 were severe or life threatening. There were 4 cases of rectal perforation, and 31% were cases requiring blood transfusion. The most common device malfunctions (8 instances of each) were motion error, and handpiece attachment fracture/ attachment malfunction. The authors concluded these complications and device related malfunctions are significant, and patients and providers should be aware of the potential risks when choosing this treatment for BPH.

In a multicenter double-blinded RCT, Gilling et al. (2022) compared the safety and efficacy of Aquablation to that of a TURP, the gold standard for BPH. 181 men aged 45-80 with BPH were randomized into either receiving Aquablation or the control group (TURP). The Aquablation was performed using the AquaBeam Robotic System. The patients were followed for 5 years and staff performing assessments were blinded for 3 years; year 4 and 5 occurred during the COVID-19 pandemic. The primary efficacy endpoint was the change in IPSS from baseline to 6 months and was successfully achieved; at 6 months the aquablation group showed slightly better numbers with an IPSS decrease of 16.9 points from baseline whereas the TURP group had a decrease of 15.1 points. At 5 years, the median IPSS score was 5.5 for the aquablation group and 6 for the TURP group. The MSHQ-EjD-SF (MSHQ-EjD) score averaged 2.7 points lower (or worse) for the TURP group compared

UnitedHealthcare, Inc. ("UHC") Proprietary and Confidential Information: The information contained in this document is confidential, proprietary and the sole property of UHC. The recipient of this information agrees not to disclose or use it for any purpose other than to facilitate UHC's compliance with applicable State Medicaid contractual requirements. Any other use or disclosure is strictly prohibited and requires the express written consent of UHC.

to the aquablation group. After 5 years, the QoL was no different between the two groups, but 12.3% of the TURP group needed additional BPH therapy while only 6% of the aquablation participants did. The authors found the health outcomes from aquablation therapy outweigh those when compared to a TURP and at 5 years, uroflow improvement continues to show durability and consistency. Limitations included the loss to follow up rate unequal number of patients in the treatment group to that of the control group, the impact the pandemic had on the overall follow up percentages at year 4 and 5 and the sole funding of the study came from the device manufacturer.

Elterman et al. (2021, ~~included in the 2018 ECRI report below~~) conducted a meta-analysis of individual patient data from patients undergoing aquablation treatment for BPH from four selected prospective global clinical trials; WATER, WATER II, FRANCAIS WATER and OPEN WATER. 425 men with BPH were evaluated with a one year follow-up. The following were items of focus: symptom scores, components of IPSS, uroflow and incontinence. In each study, participants were evaluated using transrectal ultrasound (TRUS), serum prostate specific antigen (PSA), uroflow measures and completion of the IPSS18 and Incontinence Severity Index (ISI). The authors found the IPSS scores improved significantly in all studies; and at 1-year improvement of 16 points from baseline was noted. While this study was a meta-analysis of selected study, not based on a systematic review of the literature, it was not governed by a registered protocol; further limitations include lack of comparison group, lack of long-term efficacy and a variation in patient population.

In a 2021 Hayes technology assessment regarding aquablation for treating benign prostatic hyperplasia, it was concluded that a low-quality body of evidence suggests it may improve LUTS associated with BPH in the short to intermediate term without impacting sexual or function, and without serious safety concerns. However substantial uncertainty remains due to the scarcity of evidence comparing aquablation to TURP, which is the primary surgical modality for treating BPH in small to moderate sized prostates, as well as limited long-term evidence. Furthermore, clarity is lacking as to which patient populations are likely to benefit the most from aquablation therapy. The report concludes that this technology has a potential but unproven benefit.

A 2018 ECRI clinical evidence assessment, updated in 2021, of the Aquabeam Robotic System for treating benign prostatic hyperplasia (BPH), reports this is safe, and reduces BPH-related LUTS for up to three years based on evidence from one randomized controlled trial (RCT) and four prospective case series, and this evidence is somewhat favorable. The RCT reported that Aquablation works as well as TURP for improving LUTS at three-year follow-up, but findings need confirmation in additional RCTs. RCTs of sufficient size are needed that compare AquaBeam to other minimally invasive treatments for LUTS due to BPH, as well as report patient-oriented outcomes.

Gilling et al. (2020) reported the results of participants from the Water I clinical trial to report 3-year outcomes for aquablation compared to TURP for the treatment of LUTS related to BPH. Assessments included IPAA, MSHQ-EjD, IIEF and uroflow. Over 3 years of treatment, improvements in IPSS scores were statistically similar across groups. Mean 3-year improvements were 14.4 and 13.9 points in the aquablation and TURP groups, respectively (difference of 0.6 points, 95% CI -3.3-2.2, p = .6848). Similarly, 3-year improvements in Qmax were 11.6 and 8.2 cc/sec (difference of 3.3 [95% CI -0.5-7.1] cc/sec, p = .0848). At 3 years, PSA was reduced significantly in both groups by 0.9 and 1.1 ng/mL, respectively; the reduction was similar across groups (p = .5983). There were no surgical retreatments for BPH beyond 20 months for either aquablation or TURP.

UnitedHealthcare, Inc. ("UHC") Proprietary and Confidential Information: The information contained in this document is confidential, proprietary and the sole property of UHC. The recipient of this information agrees not to disclose or use it for any purpose other than to facilitate UHC's compliance with applicable State Medicaid contractual requirements. Any other use or disclosure is strictly prohibited and requires the express written consent of UHC.

It was concluded that three-year BPH symptom reduction and urinary flow rate improvement were similar after TURP and aquablation therapy. No subjects required surgical retreatment beyond 20 months postoperatively. This study is limited by a maximum prostate size of 80cc (however Desai reported on results with larger prostates below), and whether the rigor of clinical trial data can be applied in real world settings. Furthermore, the study may have been too small to detect clinically significant differences at three years, as it was powered for non-inferiority at six months.

Desai et al. (2020) reported the 2-year safety and effectiveness of aquablation in men with larger prostate volumes of 80-150cc in a prospective, multicenter international case series (WATER II). Participants had a mean prostate volume of 107 cc and the results showed IPSS and IPSS quality of life improved from 23.2 to 1.1, and 4.6 to 1.1 from baseline to 2 years respectively. Maximum urinary flow increased from 8.7 to 18.2 cc/sec. By the end of the 2-year study timeframe, all but 2 of the 74 participants stopped taking alpha blockers and all but 32 stopped taking 5 α -reductase inhibitors. During the 2-year study time frame, adverse urological events were low and included 2 subjects with recurrent BPH symptoms that required retreatment with TURP and HOLEP. The authors concluded that the aquablation procedure is a safe and effective treatment for men with LUTS due to BPH with larger prostate volumes and has an acceptable safety profile and a low retreatment rate. This trial is limited by a lack of a control group which prevented direct comparison to other treatments.

Bach et al. (2020) conducted an international prospective, multicenter, single-arm, open-label, international clinical trial of the efficacy of the aquablation procedure for the treatment of LUTS due to BPH in 177 men enrolled at five treatment centers between September 2017 and December 2018. The primary endpoint was the change in total IPSS from baseline to 3 months. Secondary endpoints included the following: (1) Proportion of subjects who were sexually active at the baseline and experienced either ejaculatory or erectile dysfunction at 3 months, change from the baseline to 3 months in maximal flow rate (Q_{max}), prostate specific antigen (PSA) level, post-void residual (PVR), total MSHQ score, and selected IIEF-5 score. The degree of dysuria was collected on a 0 (not at all) to 5 (almost always) scale. Inclusion criteria was a diagnosis of LUTS due to BPH and a prostate size between 20 and 150 cc. Men were excluded if they were unable to stop anticoagulants and antiplatelet agents perioperatively or had a bleeding disorder, had a history of gross hematuria, were using systemic immune suppressants, had a contraindication to both general and spinal anesthesia, were unwilling to accept transfusion if required, or had any severe illness that could prevent complete follow-up. Patients with prior BPH surgery were not excluded. At baseline and 3 and 12 month follow up, participants completed the International Prostate Symptom Score (IPSS), Incontinence Severity Index, Pain Intensity Scale, Quality of Recovery Visual Analog Scale, International Index of Erectile Function (IIEF-15), the Male Sexual Health Questionnaire (MSHQ-EjD), uroflowmetry and post void residual volume (PVR) measurements. The results showed of the original 177 participants enrolled and had the procedure completed, by month 12, 30 were lost to follow up, three voluntarily withdrew, and one died of an unrelated cause. Mean IPSS improved from 21.7 (7.1) at baseline to 7.1 (5.8) at 3-month follow-up, and 6.4 (4.8) at 12-month follow-up. IPSS QOL scores improved from 4.7 (1.1) at baseline to 1.5 (1.4) at 3-month follow-up, and 1.4 (1.4) at 12-month follow-up. IPSS storage and voiding scales also improved significantly (p < 0.0001) at 3 and 12 months (Baseline IPSS scores were unavailable in nine men; of these, seven were using a urinary catheter at the baseline and two cases had incomplete questionnaire responses). Maximum urinary flow rate increased from 9.9 (5.3) cc/sec at baseline to 20.3 (11.4) cc/sec at month 3 and 20.8 (11.2) cc/s at month 12. Postvoid residual improved from 108 (108) to 47 (77) cc at three months and 61 (74) cc at 12 months. Of the 92 men that were sexually

active at baseline and 12 months, the MSHQ-EjD score changed by -1 at 3 months, and -1.1 points at 12 months. MSHQ bother/satisfaction changed by -0.3 and -0.7 points at 3 and 12 months respectively. IIEF-15 scores remained stable through month 3. 141 patients had transrectal ultrasound at baseline and after 3 months which showed a decrease in prostate size of 36%. Leakage of urine was reported by 68% of participants at baseline and had reduced to 55% at 12 months, and ISI improved non-significantly. Dysuria of any frequency was reported by 51% at baseline and 29% at 3-month follow-up, and associated pain decreased from 3.5 to 2.4. General pelvic pain decreased from 1.3 at baseline to 0.4 at 3 month follow up. 82 of the participants were taking medication for BPH preoperatively and by month 3, all but 8 had discontinued the medication. There were 69 adverse events reported in 56 participants; 33 grade 1 events, 15 grade 2 events, five grade 3a events and 16 grade 3b events. The authors concluded that aquablation is safe and effective for men with LUTS due to BPH and replicate results previously seen in a trial setting. This study is limited by a lack of a concurrent control group and a relatively short-term efficacy and follow-up.

A 2019 Cochrane review on aquablation (Hwang et al.) identified only one RCT, the Gilling study described below. The authors concluded that based on short-term (up to 12 months) follow-up, the effect of aquablation on urological symptoms is probably similar to that of TURP (moderate-certainty evidence). The effect on quality of life may also be similar (low-certainty evidence). There is uncertainty whether patients undergoing aquablation are at higher or lower risk for major adverse events (very low-certainty evidence). aquablation may result in little to no difference in erectile function but offer a small improvement in preservation of ejaculatory function (both very low certainty evidence). These conclusions are based on a single study of men with a prostate volume up to 80 mL in size. Longer-term data and comparisons with other modalities appear critical to a more thorough assessment of the role of aquablation for the treatment of LUTS in men with BPH.

Gilling et al. (2019) compared 2-year safety and efficacy outcomes after aquablation or TURP for the treatment of LUTS related to BPH. A total of 181 patients with BPH were randomly assigned (2:1 ratio) to either aquablation or TURP. Patients and follow-up assessors were blinded to treatment. Assessments included the IPSS, MSHQ, IIEF and uroflow. At 2 years, IPSS scores improved by 14.7 points in the aquablation group and 14.9 points in TURP (p = 0.8304, 95 % CI: - 2.1 to 2.6 points). Two-year improvements in Qmax were 11.2 and 8.6 cc/s for aquablation and TURP, respectively (p = 0.1880, 95 % CI: - 1.3 to 6.4). Sexual function as assessed by MSHQ was stable in the aquablation group and decreased slightly in the TURP group. At 2 years, PSA was reduced in both groups by 0.7 and 1.2 points, respectively; the reduction was similar across groups (p = 0.1816). Surgical re-treatment rates after 12 months for aquablation were 1.7 % and 0 % for TURP. Over 2 years, surgical BPH retreatment rates were 4.3 % and 1.5 % (p = 0.4219), respectively. The authors concluded that 2-year efficacy outcomes after TURP and aquablation were similar, and the rate of surgical re-treatment was low and similar to TURP; aquablation may be an alternative for men who strongly prefer maintenance of ejaculatory function. The sample size may however have been too small to detect clinically important differences.

Reale et al. (2019) performed a systematic review of case series and comparison studies, to evaluate functional outcomes (Qmax, QoL, IPSS, PVR), sexual outcome (erectile dysfunction and anejaculation rate), and adverse events evaluated according to the Clavien-Dindo classification. The functional outcomes, evaluated after water jet dissection, have shown improvement with respect to the baseline in all the selected articles. In the comparison papers with the TURP, the aquablation has been statistically not inferior regarding functional outcomes. The sexual outcomes have highlighted a better

UnitedHealthcare, Inc. ("UHC") Proprietary and Confidential Information: The information contained in this document is confidential, proprietary and the sole property of UHC. The recipient of this information agrees not to disclose or use it for any purpose other than to facilitate UHC's compliance with applicable State Medicaid contractual requirements. Any other use or disclosure is strictly prohibited and requires the express written consent of UHC.

ejaculation rate for water jet dissection than TURP. Regarding the adverse events, water jet dissection documented low rates of adverse events and, in comparison studies, were not statistically superior to TURP. Multicenter randomized trials with larger cohorts and longer follow-up are still needed.

A study to compare urodynamic outcomes between aquablation vs transurethral resection of the prostate (TURP) was performed (Pimentel et al., 2019). Patients (n=66) were randomized 2:1 (aquablation:TURP) in the Waterjet Ablation Therapy for Endoscopic Resection of prostate tissue study. Urodynamics were measured at baseline and 6 months. At mean baseline pDet@qmax was 71 and 73cm H2O in the aquablation and TURP groups, respectively. At 6-month follow-up, pDet@qmax decreased by 35 and 34cm H2O, respectively. A large negative shift in bladder outlet obstruction index was observed, consistent with a large reduction in the proportion of subjects with obstruction at follow-up compared to baseline (79% to 22% in aquablation and 96% to 22% in TURP). The authors concluded that in this trial, improvements after aquablation in objective measures of bladder outlet obstruction were similar to those observed after TURP.

Plante et al. (2018) conducted prespecified post hoc exploratory subgroup analyses from a double-blind, multicenter prospective randomized controlled trial that compared transurethral resection of the prostate (TURP) using either standard electrocautery vs surgery using robotic waterjet (aquablation) to determine whether certain baseline factors predicted more marked responses after aquablation as compared with TURP. The primary efficacy endpoint was reduction in International Prostate Symptom Score (IPSS) at 6 months. The primary safety endpoint was the occurrence of Clavien-Dindo persistent grade 1 or grade ≥ 2 surgical complications. For men with larger prostates (50-80 g), the mean IPSS reduction was four points greater after aquablation than after TURP, a larger difference than the overall result. The primary safety endpoint difference was greater for men with large prostate compared with the overall result. Postoperative anejaculation was also less common after aquablation compared with TURP in sexually active men with large prostates vs the overall results. Exploratory analysis showed larger IPSS changes after aquablation in men with enlarged middle lobes, men with severe middle lobe obstruction, men with a low baseline maximum urinary flow rate, and men with elevated post-void residual urine volume. The authors concluded that in men with moderate-to-severe lower urinary tract symptoms attributable to BPH and larger, more complex prostates, aquablation was associated with both superior symptom score improvements and a superior safety profile, with a significantly lower rate of postoperative anejaculation. The authors noted that the standardized, robotically executed, surgical approach with aquablation may overcome the increased outcome variability in more complex anatomy, resulting in superior symptom score reduction. The RCT reported short-term outcomes and included patients with a prostate size 30 to 80 cc. Therefore, results may not be generalizable for all prostate sizes.

Gilling et al. (2018) conducted a double-blind, multicenter, prospective, randomized, controlled trial (WATER I) to compare safety and efficacy of aquablation and transurethral prostate resection for the treatment of lower urinary tract symptoms related to benign prostatic hyperplasia. One hundred and eighty-one patients with moderate to severe lower urinary tract symptoms related to benign prostatic hyperplasia underwent transurethral prostate resection or aquablation. The primary efficacy end point was the reduction in International Prostate Symptom Score at 6 months. The primary safety end point was the development of Clavien-Dindo persistent grade 1, or 2 or higher operative complications. The results showed the mean total operative time was similar for aquablation and transurethral prostate resection, but resection time was lower for aquablation. At month 6 patients treated with aquablation and transurethral prostate

resection experienced large I-PSS improvements. The prespecified study noninferiority hypothesis was satisfied. Of the patients who underwent aquablation and transurethral prostate resection 26% and 42%, respectively, experienced a primary safety end point, which met the study primary noninferiority safety hypothesis and subsequently demonstrated superiority. Among sexually active men the rate of anejaculation was lower in those treated with aquablation (10% vs 36%) The authors concluded that surgical prostate resection using aquablation showed noninferior symptom relief compared to transurethral prostate resection but with a lower risk of sexual dysfunction. Larger prostates (50 to 80 ml) demonstrated a more pronounced superior safety and efficacy benefit. Longer term follow-up would help assess the clinical value of aquablation. This study was supported by PROCEPT Bio Robotics, the manufacturer of the AquaBeam® device. Several of the authors indicate a financial interest and/or other relationship with PROCEPT BioRobotics. These conflicts of interest may limit the conclusions that can be drawn from the study.

Gilling et al. (2017) performed a prospective, single arm, multicenter trial at a total of 3 centers in Australia and New Zealand with 1-year follow-up to establish the safety and effectiveness of aquablation, an image guided, robotic assisted, water jet tissue ablation technology, for the treatment of benign prostatic hyperplasia. A total of 21 men with moderate to severe lower urinary tract symptoms (LUTS) were included in the study with in-clinic follow up visits at 1, 3, 6 and 12 months. The visits included a review of AEs, uroflow measurements prostate specific antigen (PSA) measurement (at 6 and 12 months only), completion of study questionnaires, and (at 6 months only) urodynamics and transrectal ultrasound (TRUS). Symptoms related to LUTS had significantly improved from baseline at 1 month and were sustained through month 12. At 12 months, the mean international prostatic symptom score (I-PSS) score had improved by 16.2 points. The I-PSS QOL component improved by 3.3 points. Mean maximum urinary flow improved from 8.7 ml per second at baseline to 18.3 ml per second and post-void residual volume (PVR) improved from 136 to 54 ml. Prostate volume decreased from 57 ml at baseline to 35 ml. The bladder outlet obstruction index decreased from 48 at baseline to 13 a month 6. Mean serum PSA, which was measured in 20 subjects, showed no significant change from 3.15 ng/ml at baseline to 2.56 ng/ml at 12 months. No urinary incontinence developed, and sexual function was preserved postoperatively. The authors concluded that this study provides early evidence to support the safety and effectiveness of aquablation for symptomatic benign prostatic hyperplasia by improved symptom scores and other measures of obstruction. The study is of small sample size and lacks a concurrent control group.

Prostate Artery Embolization (PAE)

~~Embolization places medications or synthetic materials through a catheter into a blood vessel to block blood flow to an area of the body. It may be used to control or prevent abnormal bleeding, close off vessels supplying blood to a tumor, eliminate abnormal connections between arteries and veins, or to treat aneurysms. When performed for the prostate in patients with benign prostatic hyperplasia (BPH), microspheres are injected into the prostate blood vessels, occluding the vessels which results in the gradual shrinking of the prostate tissue which widens the urethra, alleviating urinary difficulties. Evidence from several randomized controlled trials show inconclusive findings or inferiority of prostate artery embolization as compared to established treatments for effectiveness, with some apparent benefit for adverse events. Therefore, the evidence is currently insufficient to consider this technology to be proven as non-inferior or superior to established approaches.~~

In a Cochrane review, Jung et al. (2022) completed a systematic review of literature to assess the effects of prostatic arterial embolization (PAE) compared to other procedures

UnitedHealthcare, Inc. ("UHC") Proprietary and Confidential Information: The information contained in this document is confidential, proprietary and the sole property of UHC. The recipient of this information agrees not to disclose or use it for any purpose other than to facilitate UHC's compliance with applicable State Medicaid contractual requirements. Any other use or disclosure is strictly prohibited and requires the express written consent of UHC.

for treatment of lower urinary tract symptoms in men diagnosed with benign prostatic hypertrophy. They authors focused on PAE versus transurethral resection of the prostate (TURP) which included 6 RCTs and 2 non-randomized studies (NRSs) evaluating short-term follow-up and 2 RCTs and 1 NRS evaluating long-term follow-up. The evidence suggests that PAE may provide similar improvement in urologic symptom scores and quality of life when compared to TURP, but there is high uncertainty regarding major adverse events and PAE likely increases retreatment rates. While erectile function was similar for both groups, PAE may reduce ejaculatory disorders. The authors noted that the certainty of evidence for the outcomes measured in this review was low or very low except for retreatment which was moderate-certainty evidence indicating that confidence in the reported effect size is limited to very limited and should be better informed by future research.

Sajan et al. (2022) conducted a systematic review and network analysis on the outcomes of minimally invasive therapies for LUTS secondary to BPH. Nine studies were included which contained 1,034 patients. The following comparisons were identified: 4 studies focused on PAE versus TURP and then the following individual studies: PAE versus sham, Urolift versus TURP, Urolift versus sham, Rezum versus Sham, and aquablation versus TURP. Data for IPSS, QoL, QMax, PVR, and prostate volume were all obtained presurgical for baseline values and then again at 3-, 6-, and 12-months; primary outcome measured was the IPSS scores. Four RCTs compared PAE to TURP and one RCT compared PAE versus sham. No major IPSS differences were noted but for PAE, the IPSS mean difference was one of the lowest at 12 months. No significant differences were found in Qmax, QoL, and PVR. The sham group (Rezum vs sham, Urolift vs sham and PAE vs sham) found significant differences favoring the TURP for Qmax, PVR, and QoL with no other substantial differences noted. The authors found the main strength of PAE were the 5 RCTs studies with four direct comparisons to TURP and the findings of lower in hospital costs. The disadvantages were a longer procedural time, exposure to radiation and potential for nontarget embolization. The authors concluded there were clinical benefits for PAE with minimal adverse effects. The analysis is limited by the indirectness of network meta-analyses and inclusions of studies not specifically designed to test non-inferiority of PAE compared to established approaches.

In a 2021 systematic review and meta-analysis, Xiang et al. investigated the efficacy and safety of PAE versus TURP in patients with BPH. Eleven randomized controlled trials (RCTs) met the selection criteria, and ten independent patient series were included in the final analysis. Pooled estimates were inconclusive for the difference between TURP and PAE for patient-reported outcomes including International Prostate Symptom Score (2.32 (- 0.44 to 5.09)) and quality of life (0.18 (- 0.41 to 0.77)) at 12 months. PAE was less effective regarding improvements in most functional outcomes such as maximum flow rate, prostate volume, and prostate-specific antigen. PAE may however be associated with relatively fewer complications, lower cost, and shorter hospitalization. After the PAE procedure, the overall weighted mean differences for all outcomes except sexual health scores were significantly improved from baseline during follow-up to 24 months. The authors concluded that PAE is non-inferior to TURP with regard to improving patient-reported outcomes, though most functional parameters undergo more improvement after TURP than after PAE. They also concluded that PAE can significantly continue to relieve symptoms for 24 months without causing serious complications. The findings are limited by the overall sample size that may have been too small to demonstrate non-inferiority. For example, the upper limit of the pooled estimate for the International Prostate Symptom Score was 5 on a scale from 0 to 35. Furthermore, inferiority of PEA, compared to TURP was shown on other outcomes, with the exception of adverse events.

UnitedHealthcare, Inc. ("UHC") Proprietary and Confidential Information: The information contained in this document is confidential, proprietary and the sole property of UHC. The recipient of this information agrees not to disclose or use it for any purpose other than to facilitate UHC's compliance with applicable State Medicaid contractual requirements. Any other use or disclosure is strictly prohibited and requires the express written consent of UHC.

Xu et al. (2021) conducted a small case series to assess the safety and efficacy of PAE for large BPH and severe LUTS in 28 patients over the age of 80 who were not suitable candidates for open or endoscopic surgical procedures. PAE was performed using microspheres and functional outcomes including International Prostate Symptom Score (IPSS), quality of life (QoL), maximum urine flow rate (Qmax), post-void residual urine volume, prostate volume and total prostate-specific antigen level were evaluated at 1, 3, 6, and 12 months postoperatively. Safety was evaluated using perioperative data and included operative time, fluoroscopy time, changes in hemoglobin within 24 hours postoperatively, hospitalization days, postoperative duration, as well as complications. Bilateral PAE was performed in 25 patients, and 2 received unilateral PAE. The results showed technical success with PAE in 27 of the 28 participants. All of the functional outcomes results were significantly improved at 12 months postoperatively compared to baseline. The overall complication rate was 46.4%, and included post-embolization syndrome, hematuria, urinary tract infection, and acute urinary retention. The authors concluded that PAE may be an effective treatment option for patients with BPH that are not suitable candidates for open or endoscopic procedures following failed treatments. This study is limited by a lack of comparison group, a small number of participants and a short follow up period. Furthermore, radiation doses and fluoroscopy time were not examined.

In 2021, Abt et al. reported the two-year safety and efficacy outcomes of the open label, randomized non-inferiority trial they conducted in 2018 for which 12-week outcomes were reported previously. In the 2018 trial (included in the Xiang systematic review), 103 participants aged 40 or greater with refractory LUTS secondary to benign prostatic obstruction (BPO) were treated with either PAE using 250-400 µm microspheres under local anesthesia, or monopolar transurethral resection of the prostate (TURP) under spinal or general anesthesia. International Prostate Symptoms Score (IPSS) and other patient reported outcomes, functional measures, prostate volume, and adverse events were evaluated. Changes from baseline to 2 years were tested for differences between the two interventions with standard two-sided tests. For the participants that received PAE, the results showed the mean reduction in IPSS was 9.21 points, and 12.09 points after TURP (difference of 2.88 [95% confidence interval 0.04-5.72]; p = 0.047). TURP showed superiority for most other patient reported outcomes as well (except erectile dysfunction), including maximum urinary flow rate, reduction of postvoid residual urine, and reduction of prostate volume. Adverse events were less frequent after PAE than after TURP, but the severity was similar. 21% of participants who initially received PAE required TURP within 2 years due to unsatisfactory results. The authors concluded that PAE for the treatment of BPH remains investigational due to inferior functional outcomes and a relevant re-treatment rate found 2 years after PAE compared with TURP. These disadvantages should be considered for patient selection and counselling.

~~In 2020, a Cochrane systematic review was performed to assess the effects of prostatic arterial embolization (PAE) compared to transurethral resection of the prostate (TURP) for the treatment of lower urinary tract symptoms in men with benign prostatic hyperplasia (BPH). Mean age, IPSS, and prostate volume of participants were 66 years, 22.8, and 72.8 mL, respectively. Primary outcomes assessed were urological symptom scores, quality of life and major adverse events. Based on the 6 RCTs with up to 12 month follow up (short term), as measured by the International Prostatic Symptom Score (IPSS), the authors found little to no difference in urological symptom score, and quality of life, ((with improvement mean difference (MD) 1.55, 95% confidence interval (CI) -0.40 to 3.50 in 369 participants in 6 RCTs with 75% low certainty evidence on a scale from 0 to 35. Quality of Life (QOL) results also showed little to no difference with MD 0.16, 95% CI -0.37 to 0.68 in 309 participants in 5 RCTs with 56% low certainty evidence on a~~

UnitedHealthcare, Inc. ("UHC") Proprietary and Confidential Information: The information contained in this document is confidential, proprietary and the sole property of UHC. The recipient of this information agrees not to disclose or use it for any purpose other than to facilitate UHC's compliance with applicable State Medicaid contractual requirements. Any other use or disclosure is strictly prohibited and requires the express written consent of UHC.

~~scale from 0-6.)) There is low certainty evidence regarding the effects of PAE on major adverse events, with perineal pain, hematuria, and acute urinary retention commonly reported. Additionally, there was inconsistency in reporting and classifying the adverse events. PAE may increase the need for retreatment, with 47 or more (0 more to 214 more) per 1000 men receiving PAE compared to 18 re-treatments per 1000 men who received TURP. Regarding the effects of PAE on erectile dysfunction, the authors reported very low certainty evidence as measured by the International Index of Erectile Function (IIEF) that PAE may reduce ejaculatory disorders. Only one RCT reported 13-24 month (long term) follow up and the results showed in 95 participants very low certainty evidence on urological symptom scores and QOL compared to TURP, as well as major adverse events. There was no evidence regarding erectile function and ejaculation disorders. 56 re-treatments per 1000 men in the TURP group were identified, compared to 143 more (25 more to 430 more) per 1,000 in the PAE group. The authors concluded that the evidence for the main outcomes assessed is low to very low and this procedure needs future research. (Jung et al., 2020)~~

In a ~~May 2020~~**2019** Hayes ~~comparative effectiveness review~~**health technology assessment, updated in 2022, of** regarding prostatic artery embolization (PAE) for the treatment of benign prostatic hypertrophy (BPH), it was concluded that based on a moderately large body of evidence, ~~and albeit~~ low quality due to individual study limitations, PAE is reasonable safe, but long-term efficacy has not been adequately evaluated. **The overall conclusion of the report is that there is potential but unproven benefit of this technology.**

In a 2019 systematic review and meta-analysis, Zumstein et al. performed a systematic review and meta-analysis of clinical trials comparing the efficacy and safety of prostate artery embolization (PAE) to established surgical therapies. Functional parameters assessed included maximum urinary flow, post void residual, and reduction of prostate volume. There were 5 comparative studies consisting of 708 patients, some of which had an unclear risk of bias in patient selection, blinding, and incomplete outcome data. Reporting of complications varied widely and was poor in some. The results showed that compared to standard surgical therapies PAE showed less improvement in the International Prostate Symptom Score and was less efficient in a in all functional parameters assessed. Conversely, patient reported erectile function was better after PAE and there were significantly fewer adverse events overall. The authors concluded that PAE is safe and effective in the short term, particularly regarding safety and sexual function, but clear disadvantages for all other patient reported and functional outcomes assessed compared to established surgical therapies were identified. This suggests PAE is not as effective as established surgical therapies. The authors recommend large scale randomized controlled trials that include longer follow up, as well as defining ideal indications are mandatory before PAE can be considered a standard treatment option.

Abt et al. (2018) conducted a randomized, open label, non-inferiority trial in the urology and radiology departments of a Swiss tertiary care center. 103 patients aged ≥40 years with refractory lower urinary tract symptoms secondary to benign prostatic hyperplasia were randomized to receive prostatic artery embolization (PAE) with 250-400 µm microspheres under local anesthesia, or monopolar transurethral resection of the prostate (TURP) under spinal or general anesthesia. 48 and 51 patients reached the primary endpoint 12 weeks after PAE and TURP, respectively. Primary outcome was change in international prostate symptoms score (IPSS) from baseline to 12 weeks after surgery (a difference of less than 3 points between treatments was defined as non-inferiority for PAE and tested with a one-sided t test). Secondary outcomes included further questionnaires functional measures, magnetic resonance imaging findings and adverse

events. Changes from baseline to 12 weeks were compared between treatments with two sided tests for superiority. The authors failed to prove non-inferiority for the primary outcome (1.54 points in favor of TURP (95% confidence interval -1.45 to 4.52)), but fewer adverse events occurred after PAE than after TURP (36 v 70 events; P=0.003). (This trial was included in the systematic review by Xiang et al., 2021, and Sajan et.al., 2022 ~~Jung et al. 2020~~).

~~The 2018 NICE guidelines for prostate artery embolization for lower urinary tract symptoms caused by BPH states that the current evidence of the safety and efficacy is adequate to support the use of this procedure provided that standard arrangements are in place for clinical governance, consent and audit. Furthermore, patient selection should be done by a urologist and an interventional radiologist. This procedure is technically demanding and should only be done by an interventional radiologist with specific training and expertise in prostatic artery embolization.~~

Clinical Practice Guidelines

American Urological Association (AUA)

In ~~2020~~ **2021**, the AUA (~~Lerner Parsons et al.~~) ~~amended~~ **revised** their clinical guidelines on the surgical management of BPH/LUTS. Included in their guideline statements are the following:

- PUL may be offered as an option for patients with LUTS attributed to BPH provided prostate volume <80g and verified absence of an obstructive middle lobe (Moderate Recommendation; Evidence Level: Grade C).
- PUL may be offered to eligible patients who desire preservation of erectile and ejaculatory function (Conditional Recommendation; Evidence Level: Grade C).
- Aquablation may be offered to patients provided prostate volume >30/<80g. (Conditional Recommendation; Evidence Level: Grade C)
- Water vapor thermal therapy:
 - o May be offered to patients with LUTS/BPH with a prostate volume of <80g. patients should be informed that evidence of efficacy, including longer-term retreatment rates, remains limited. (Conditional Recommendation; Evidence Level: Grade C)
 - o Patients who desire preservation of erectile and ejaculatory function (Conditional Recommendation; Evidence Level: Grade C)
- Prostate artery embolization is not recommended for the treatment of LUTS/BPH outside the context of a clinical trial until sufficient evidence from rigorous studies is available that show benefit over well-established therapies (Expert Opinion)

American Urological Association (AUA) / American Society for Radiation Oncology (ASTRO) / ~~Society of Urologic Oncology (SUO)~~

~~In a 2017 AUA/ASTRO/SUO practice guideline for clinically localized prostate cancer states that if focal therapy is offered as an alternative treatment modality for localized prostate cancer, it should only be done within the context of a clinical trial. Initial studies with short term follow up suggest that effective disease eradication in the treated volume can be attained, however, it should be noted that long-term follow up data is lacking. The Panel recognizes that concern exists about the potential for undetected and, therefore, occult untreated clinically significant multifocal disease. The also state that confirmation of oncologic effectiveness is currently lacking and will require prospective studies with long-term follow up. Additionally, this practice guideline states that prostate cancer is often multifocal, clinicians should inform localized prostate cancer patients considering focal therapy that focal therapy may not be curative and that further treatment for prostate cancer may be necessary. e~~

UnitedHealthcare, Inc. ("UHC") Proprietary and Confidential Information: The information contained in this document is confidential, proprietary and the sole property of UHC. The recipient of this information agrees not to disclose or use it for any purpose other than to facilitate UHC's compliance with applicable State Medicaid contractual requirements. Any other use or disclosure is strictly prohibited and requires the express written consent of UHC.

The 2022 AUA/ASTRO guidelines for clinically localized prostate cancer from which are endorsed by the Society of Urologic Oncology (SUO) state the following:

- For patients with favorable intermediate-risk prostate cancer, clinicians should discuss active surveillance, radiation therapy, and radical prostatectomy. (Strong Recommendation; Evidence Level: Grade A)
- Clinicians should inform patients with intermediate-risk prostate cancer considering whole gland or focal ablation that there are a lack of high-quality data comparing ablation outcomes to radiation therapy, surgery, and active surveillance. (Expert Opinion)
- For patients with unfavorable intermediate- or high-risk prostate cancer and estimated life expectancy greater than 10 years, clinicians should offer a choice between radical prostatectomy or radiation therapy plus androgen deprivation therapy (ADT). (Strong Recommendation; Evidence Level: Grade A)

~~The clinical guidelines on advanced prostate cancer (Lowrance et al., 2020), and clinically localized prostate cancer (Sanda et al., 2017) do not include water vapor therapy as a treatment option.~~

The National Comprehensive Cancer Network (NCCN)

Clinical practice guidelines for the treatment of prostate cancer (~~v21~~v4.2021~~2021~~2022) states "biocompatible and biodegradable perirectal spacer materials may be implanted between the prostate and rectum in patients undergoing external radiotherapy with organ-confined prostate cancer in order to displace the rectum from high radiation dose regions. Perirectal spacer materials may be employed when the other techniques are insufficient to improve oncologic cure rates and/or reduce side effects due to anatomic geometry or other patient related factors, such as medication usage and/or comorbid conditions. Patients with obvious rectal invasion or visible T3 and posterior extension should not undergo perirectal spacer implantation.

For cryotherapy, the guidelines state that "cryotherapy or other local therapies are not recommended as routine primary therapy for localized prostate cancer due to lack of long-term data comparing these treatments to radiation or radical prostatectomy." Presently, the panel recommends cryosurgery and HIFU as the only local therapy options for radiation therapy recurrence in the absence of metastatic disease.

For radical prostatectomy, the guidelines state: "Radical prostatectomy is appropriate for any patient whose cancer appears clinically localized to the prostate. However, because of potential perioperative morbidity, radical prostatectomy should generally be reserved for patients whose life expectancy is 10 years or more." Radical prostatectomy is listed as one of the options for some high or very high-risk patients, and some patients with cancer recurrence.

NCCN Clinical Practice Guidelines in Oncology do not mention water vapor thermotherapy for the treatment of prostate cancer (~~2021~~ 2022).

National Institute for Health and Care Excellence

A 2020 NICE medical technology guideline for the use of Rezum for treating lower urinary tract symptoms secondary to benign prostatic hyperplasia states that the evidence supports adopting Rezum for treating lower urinary tract symptoms (LUTS) caused by benign prostatic hyperplasia (BPH) in the NHS. It should be considered as a treatment option for

UnitedHealthcare, Inc. ("UHC") Proprietary and Confidential Information: The information contained in this document is confidential, proprietary and the sole property of UHC. The recipient of this information agrees not to disclose or use it for any purpose other than to facilitate UHC's compliance with applicable State Medicaid contractual requirements. Any other use or disclosure is strictly prohibited and requires the express written consent of UHC.

men with moderate to severe LUTS (International Prostate Symptoms Score [IPSS] typically 13 or over), and a moderately enlarged prostate (typically between 30 cm and 80 cm).

The 2018 NICE medical technology guidance on use of the Memokath-051 stent for ureteric obstruction concludes that the quality of reporting across all the studies was generally poor. None of the studies provided adequate details on patient characteristics, stent insertion procedures, follow-up, statistical analyses and uncertainty around the results. Migration rates and clinical success were the most commonly reported outcomes, but definitions of clinical success varied, so statistical pooling could not be done.

The 2018 NICE guidelines for prostate artery embolization for lower urinary tract symptoms caused by BPH states that the current evidence of the safety and efficacy is adequate to support the use of this procedure provided that standard arrangements are in place for clinical governance, consent and audit. Furthermore, patient selection should be done by a urologist and an interventional radiologist. This procedure is technically demanding and should only be done by an interventional radiologist with specific training and expertise in prostatic artery embolization.

Society of Interventional Radiology (SIR)

In a ~~2020~~ **2019** (McWilliams et al.) multi-society, evidence-based position statement regarding PAE for the treatment of lower urinary tract symptoms due to BPH, the SIR states that PAE is a safe and effective treatment, has good short and intermediate term efficacy and is a treatment option for the following:

- For appropriately selected men with BPH and moderate to severe LUTS (strong recommendation)
- In patients with BPH and moderate to severe LUTS who have very large prostate glands (> 80 cm³), without an upper limit of prostate size (moderate recommendation)
- In patients with BPH and acute or chronic urinary retention in the setting of preserved bladder function as a method of achieving catheter independence (moderate recommendation)
- In patients with BPH and moderate to severe LUTS who wish to preserve erectile and/or ejaculatory function (weak recommendation)
- In patients with hematuria of prostatic origin as a method of achieving cessation of bleeding (strong recommendation)
- in patients with BPH and moderate to severe LUTS who are deemed not to be surgical candidates for any of the following reasons: advanced age, multiple comorbidities, coagulopathy, or inability to stop anticoagulation or antiplatelet therapy (moderate recommendation)
- PAE should be included in the individualized patient centered discussions regarding treatment options (strong recommendation)

SIR also gives a strong recommendation that Interventional radiologists, given their knowledge of arterial anatomy, advanced microcatheter techniques, and expertise in embolization procedures, are the specialists best suited for the performance of PAE. (McWilliams et al., 2019)

U.S. Food and Drug Administration (FDA)

This section is to be used for informational purposes only. FDA approval alone is not a basis for coverage.

UnitedHealthcare, Inc. ("UHC") Proprietary and Confidential Information: The information contained in this document is confidential, proprietary and the sole property of UHC. The recipient of this information agrees not to disclose or use it for any purpose other than to facilitate UHC's compliance with applicable State Medicaid contractual requirements. Any other use or disclosure is strictly prohibited and requires the express written consent of UHC.

Prostate surgeries are procedures and, therefore, not regulated by the FDA. However, devices and instruments used during the surgery may require FDA approval. See the following website for additional information:

<http://www.accessdata.fda.gov/scripts/cdrh/cfdocs/cfPMN/pmn.cfm>. (Accessed ~~June 3, 2021~~ **May 19, 2022**)

On September 13, 2013, the FDA approved the UroLift® System (Teleflex Inc., Pleasanton, CA) for marketing through a de novo classification as a class II device used as a permanent implant to relieve low or blocked urine flow in men with BPH. Since that time, additional FDA clearances have been granted. For additional information see the following, using the product code PEW:

<https://www.accessdata.fda.gov/scripts/cdrh/cfdocs/cfpmn/pmn.cfm?ID=K193269>. (Accessed ~~June 3, 2021~~ **May 19, 2022**)

On August 2, 2019, The U.S. Food and Drug Administration (FDA) cleared the Rezūm™ Water Vapor Therapy system (Boston Scientific Corp.) under 510(k) premarket notification for treatment of symptoms of benign prostatic hyperplasia (BPH), and treatment of the prostate with hyperplasia of the central zone and/or a median lobe. It is not approved for treatment of malignant prostate tissue. For additional information, see:

<https://www.accessdata.fda.gov/scripts/cdrh/cfdocs/cfpmn/pmn.cfm?ID=K191505>. (Accessed ~~April 16, 2021~~ **May 19, 2022**)

The U.S. Food and Drug Administration (FDA) has cleared powered laser devices under 510(k) Premarket Notification. For device specific information, search product code GEX here: <https://www.accessdata.fda.gov/scripts/cdrh/cfdocs/cfpmn/pmn.cfm>. (Accessed ~~April 15, 2021~~ **May 19, 2022**)

The U.S. Food and Drug Administration (FDA) cleared SpaceOAR Vue hydrogel (Boston Scientific Corporation) (K182971) under its 510(k) premarket notification process as substantially equivalent to predicate devices on June 19, 2019. For additional information see the following:

https://www.accessdata.fda.gov/cdrh_docs/pdf18/K182971.pdf. (Accessed May ~~20, 2021~~ **19, 2022**)

The U.S. Food and Drug Administration (FDA) approved the Spanner® Temporary Prostatic Stent (SRS Medical, North Billerica, MA) on December 14, 2006. Refer to the following website for additional information:

- https://www.accessdata.fda.gov/scripts/cdrh/cfdocs/cfpma/pma_template.cfm?id=p060010
 - https://www.accessdata.fda.gov/cdrh_docs/pdf6/p060010a.pdf
- (Accessed May ~~5, 2021~~ **19, 2022**)

In December 2017, the FDA granted a De Novo request for the ITind system (Olympus America, Center Valley, PA) (DEN190020), a temporarily-placed system for the urethra to treat urinary symptoms associated with BPH. Refer to the following website for additional information: <https://www.accessdata.fda.gov/scripts/cdrh/cfdocs/cfPMN/denovo.cfm>. (Accessed May ~~5, 2021~~ **19, 2022**)

On March 3, ~~2021~~ **2021**, the Aquabeam® Robotic System (Procept BioRobotics, Redwood City, CA) received 510(k) approval as a Class II device. Refer to the following for further information: <https://www.accessdata.fda.gov/scripts/cdrh/cfdocs/cfPMN/pmn.cfm?ID=K202961>. (Accessed May ~~21, 2021~~ **19, 2022**).

UnitedHealthcare, Inc. ("UHC") Proprietary and Confidential Information: The information contained in this document is confidential, proprietary and the sole property of UHC. The recipient of this information agrees not to disclose or use it for any purpose other than to facilitate UHC's compliance with applicable State Medicaid contractual requirements. Any other use or disclosure is strictly prohibited and requires the express written consent of UHC.

In June 2017, the FDA granted a De Novo request for Embosphere® Microspheres (Merit Medical Systems, Jordan, UT) for embolization of prostatic arteries for symptomatic benign prostatic hyperplasia. Refer to the following website for additional information: <https://www.accessdata.fda.gov/scripts/cdrh/cfdocs/cfpmn/denovo.cfm?id=DEN160040>. (Accessed ~~June 10, 2021~~ **May 19, 2022**).

For additional information on microsphere products with 510(k) premarket notification, refer to the following website and search by product code NOY: <https://www.accessdata.fda.gov/scripts/cdrh/cfdocs/cfPMN/pmn.cfm>. (Accessed ~~June 10, 2021~~ **May 19, 2022**)

The ECHOLASER X4 system received 510(k) Premarket Notification (K181510) from the FDA in September of 2018. The device is intended for use in cutting, vaporization, ablation and coagulation of soft tissue and in the treatment and/or removal of vascular lesions (tumors). For additional information, refer to the following website: <https://www.accessdata.fda.gov/scripts/cdrh/cfdocs/cfPMN/pmn.cfm>. (Accessed May 24, 2022)

The Memokath has not yet received FDA approval.

References

Abt D, Hechelhammer L, Müllhaupt G, et al. Comparison of prostatic artery embolization (PAE) versus transurethral resection of the prostate (TURP) for benign prostatic hyperplasia: randomized, open label, non-inferiority trial. BMJ. 2018 Jun 19;361:k2338.

Abt D, Müllhaupt G, Hechelhammer L, et al. Prostatic ~~Artery~~ artery ~~Embolization~~ embolization ~~Versus~~ versus ~~Transurethral~~ transurethral ~~Resection~~ resection of the Prostate ~~prostate~~ for Benign ~~benign~~ Prostatic ~~prostatic~~ Hyperplasia hyperplasia: 2-yr Outcomes ~~outcomes~~ of a ~~Randomized~~ randomized, Open ~~open~~-label, Single ~~single~~-center Trial trial. Eur Urol. 2021 Jul;80(1):34-42.

Afkhami Ardekani M, Ghaffari H. Optimization of prostate brachytherapy techniques with polyethylene glycol-based hydrogel spacers: A systematic review. Brachytherapy. 2020 Jan-Feb;19(1):13-23.

Ahn ST, Lee DH, Kim JW, et al. Novel treatment strategy for management of traumatic bulbar urethral rupture using temporary urethral stent after primary realignment; retrospective comparison between thermo-expandable urethral stent and self-expandable polymer-coated urethral stent. J Clin Med. 2020;9(5): E1274. Published 2020 Apr 28.

~~American Urological Association. Clinically localized prostate cancer: AUA/ASTRO/SUO guideline. 2017. Available at: <https://www.auanet.org/guidelines/prostate-cancer-clinically-localized-guideline> Accessed April 15, 2021~~ **May 10, 2022.**

American Urological Association. Clinically Localized Prostate Cancer: AUA/ASTRO Guideline 2022. Available at: <https://www.auanet.org/guidelines/guidelines/clinically-localized-prostate-cancer-aua/astro-guideline-2022>. Accessed May 23, 2022.

Bach T, Gilling P, El Hajj A, et al. First multi-center all-comers study for the aAquablation procedure. J Clin Med. 2020 Feb 24;9(2):603.

Bates AS, Ayers J, Kostakopoulos N, et al. A systematic review of focal ablative therapy for clinically localised prostate cancer in comparison with standard management options: limitations of the available evidence and recommendations for clinical practice and further research. Eur Urol Oncol. 2021 Jan 8: S2588-9311(20)30216-9.

Bhojani N, Bidair M, Zorn KC, et al. Aquablation for Bbenign Pprostatic Hhyperplasia in Llarge Pprostates (80-150 cc): 1-Yyear Rresults. Urology. 2019 Jul;129:1-7.

UnitedHealthcare, Inc. ("UHC") Proprietary and Confidential Information: The information contained in this document is confidential, proprietary and the sole property of UHC. The recipient of this information agrees not to disclose or use it for any purpose other than to facilitate UHC's compliance with applicable State Medicaid contractual requirements. Any other use or disclosure is strictly prohibited and requires the express written consent of UHC.

~~Chao B, Llukani E, Lepor H. Two-year outcomes following focal laser ablation of localized prostate cancer. Eur Urol Oncol. 2018 Jun;1(2):129-133.~~

Chao M, Ho H, Chan Y, et al. Prospective analysis of hydrogel spacer for patients with prostate cancer undergoing radiotherapy. BJU Int. 2018 Sep;122(3):427-433.

Chughtai B, Elterman D, Shore N, et al. The iTind temporarily implanted nitinol device for the treatment of lower urinary tract symptoms secondary to benign prostatic hyperplasia: a multicenter, randomized, controlled trial. Urology. 2020 Dec 26:S0090-4295(20)31520-X.

ClinicalTrials.gov. Available at: <https://clinicaltrials.gov/ct2/home>. Accessed May 19, 2021.

de Rienzo G, Lorusso A, Minafra P, et al. Transperineal interstitial laser ablation of the prostate, a novel option for minimally invasive treatment of benign prostatic obstruction. Eur Urol. 2021 Jul;80(1):95-103.

Desai M, Bidair M, Bhojani N, et al. Aquablation for benign prostatic hyperplasia in large prostates (80-150 cc): 2-year results. Can J Urol. 2020 Apr;27(2):10147-10153.

Dineen MK, Shore ND, Lumerman JH, et al. Use of a temporary prostatic stent after transurethral microwave thermotherapy reduced voiding symptoms and bother without exacerbating irritative symptoms. Urology. 2008 May;71(5):873-7.

ECRI Institute. Aquabeam Robotic System (Procept Biorobotics Corp.) for Treating Benign Prostatic Hyperplasia. Plymouth Meeting (PA): ECRI Institute; October 2018; uUpdated June 23, 2020 July 2021.

ECRI Institute. Product Brief. UroLift System (NeoTract, Inc.) for treating benign prostatic hyperplasia. July 2019.

ECRI Institute. Clinical Evidence Assessment. Rezūm System (Boston Scientific Corp.) for Treating Benign Prostatic Hyperplasia. November 2016; Updated January 2022.

ECRI Institute. Product Brief. Spanner Prostatic Stent (SRS Medical) for maintaining urine flow after treatment for benign prostatic hyperplasia. April 2018.

ECRI Institute. Custom Product Briefs Guidance. SpaceOAR System (Augmenix, Inc.) hydrogel spacer for reducing rectal exposure during radiation therapy for prostate cancer. February 2016. Updated February 2020.

ECRI Institute. Clinical Evidence Assessment. Transperineal Laser Ablation for Treating Benign Prostatic Hyperplasia. May 2022.

~~Egilmez T, Aridogan IA, Yachia D, et al. Comparison of nitinol urethral stent infections with indwelling catheter-associated urinary tract infections. J Endourol. 2006 Apr;20(4):272-7.~~

Elterman D, Gilling P, Roehrborn C, et al. Meta-analysis with individual data of functional outcomes following aAquablation for lower urinary tract symptoms due to BPH in various prostate anatomies. BMJ Surg Interv Health Technol. 2021 Jun 23

Emam A, Hermann G, Attwood K, et al. Oncologic outcome of radical prostatectomy versus radiotherapy as primary treatment for high and very high risk localized prostate cancer. Prostate. 2021 Mar;81(4):223-230.

Gilling P, Barber N, Bidair M, et al. Three-year outcomes after aAquablation therapy compared to TURP: results from a blinded randomized trial. Can J Urol. 2020 Feb;27(1):10072-10079.

UnitedHealthcare, Inc. ("UHC") Proprietary and Confidential Information: The information contained in this document is confidential, proprietary and the sole property of UHC. The recipient of this information agrees not to disclose or use it for any purpose other than to facilitate UHC's compliance with applicable State Medicaid contractual requirements. Any other use or disclosure is strictly prohibited and requires the express written consent of UHC.

Gilling PJ, Barber N, Bidair M, et al. Five-year outcomes for aquablation therapy compared to TURP: results from a double-blind, randomized trial in men with LUTS due to BPH. Can J Urol. 2022 Feb;29(1):10960-10968.

Gilling P, Anderson P and Tan A. Aquablation of the prostate for symptomatic benign prostatic hyperplasia: 1-year results. J Urol. 2017 Jun;197(6):1565-1572.

Gilling P, Barber N, Bidair M, et al. A double-blind, randomized controlled trial of ~~a~~Aquablation vs transurethral resection of the prostate in benign prostatic hyperplasia. J Urol. 2018 May;199(5):1252-1261.

Gilling PJ, Barber N, Bidair M, et al. Randomized Controlled Trial of ~~a~~Aquablation versus ~~t~~Transurethral ~~r~~Resection of the ~~p~~Prostate in ~~B~~benign ~~p~~Prostatic ~~h~~Hyperplasia: One-year Outcomes. Urology. 2019; 125:169-173.

Goh MH, Kastner C, Khan S, et al. First experiences with the Spanner™ temporary prostatic stent for prostatic urethral obstruction. Urol Int. 2013;91(4):384-90.

Gratzke C, Barber N, Speakman MJ, et al. Prostatic urethral lift vs transurethral resection of the prostate: 2-year results of the BPH 6 prospective, multicenter, randomized study. BJU Int 2017 May;119(5):767-775.

~~Grimsley SJ, Khan MH, Lennox E, et al. Experience with the spanner prostatic stent in patients unfit for surgery: an observational study. J Endourol. 2007 Sep;21(9):1093-6.~~

Hamstra DA, Mariados N, Sylvester J, et al. Continued benefit to rectal separation for prostate radiation therapy: final results of a phase III trial. Int J Radiat Oncol Biol Phys. 2017 Apr 1;97(5):976-985.

Hayes, Inc. Evolving Evidence Review. iTind (Olympus) for Benign Prostatic Hyperplasia. Lansdale, PA: Hayes Inc.; March 2022.

Hayes, Inc. Health Technology Assessment. Rezūm System (Boston Scientific Corp.) for Benign Prostatic Hyperplasia. Lansdale, PA: Hayes Inc.; October 2021.

Hayes, Inc. Health Technology Assessment. Prostatic urethral lift (UroLift System) for treatment of symptoms associated with benign prostatic hyperplasia. Lansdale, PA: Hayes Inc.; June 2020; amended August 2021. ~~±~~Updated July 2021.

Hayes, Inc. Aquablation for Treatment of Benign Prostatic Hyperplasia. Lansdale, PA: Hayes, Inc.; March 30, 2021; amended April 2021. Updated March 2022.

Hayes, Inc. Health Technology ~~Annual Review~~ Assessment. Comparative Effectiveness Review of Prostatic Artery Embolization (PAE) for Treatment of Benign Prostatic Hypertrophy (BPH). ~~May 15, 2020~~ February 2019; amended August 2019. Updated January 2022.

Hayes, Inc. Health Technology Brief Assessment. Absorbable perirectal spacer (SpaceOAR) System; Augmenix Inc.) during radiation therapy for prostate cancer. Lansdale, PA: Hayes, Inc., ~~April 2020~~ September 2021.

Hopstaken JS, Bomers JGR, Sedelaar MJP, et al. An updated systematic review on focal therapy in localized prostate cancer: what has changed over the past 5 years? Eur Urol. 2022 Jan;81(1):5-33.

Hwang EC, Jung JH, Borofsky M, et al. Aquablation of the prostate for the treatment of lower urinary tract symptoms in men with benign prostatic hyperplasia. Cochrane Database Syst Rev. 2019;2(2):CD013143. Published 2019 Feb 13.

iTIND. Available at: <https://www.itind.com/>. Accessed May 5, 2021.

UnitedHealthcare, Inc. ("UHC") Proprietary and Confidential Information: The information contained in this document is confidential, proprietary and the sole property of UHC. The recipient of this information agrees not to disclose or use it for any purpose other than to facilitate UHC's compliance with applicable State Medicaid contractual requirements. Any other use or disclosure is strictly prohibited and requires the express written consent of UHC.

Jordan GH, Wessells H, Secrest C, et al. United States Study Group. Effect of a temporary thermo-expandable stent on urethral patency after dilation or internal urethrotomy for recurrent bulbar urethral stricture: results from a 1-year randomized trial. J Urol. 2013 Jul;190(1):130-6.

~~Jung JH, McCutcheon KA, Borofsky M, et al. Prostatic arterial embolization for the treatment of lower urinary tract symptoms in men with benign prostatic hyperplasia. Cochrane Database Syst Rev. 2020 Dec 19;12:CD012867.~~

Jung JH, McCutcheon KA, Borofsky M, et al. Prostatic arterial embolization for the treatment of lower urinary tract symptoms in men with benign prostatic hyperplasia. Cochrane Database Syst Rev. 2022 Mar 29;3(3):CD012867.

Jung JH, Risk MC, Goldfarb R, et al. Primary cryotherapy for localised or locally advanced prostate cancer. Cochrane Database Syst Rev. 2018 May 30;5(5):CD005010.

Kang TW, Jung JH, Hwang EC, et al. Convective radiofrequency water vapour thermal therapy for lower urinary tract symptoms in men with benign prostatic hyperplasia. Cochrane Database Syst Rev. 2020 Mar 25;3(3):CD013251.

Kaplan-Marans E, Martinez M, Wood A, et al. Aquablation, Prostatic Urethral Lift, and Transurethral Water Vapor Therapy: A comparison of device-related adverse events in a national registry. J Endourol. 2022 Feb;36(2):231-235.

Karsh LI, Gross ET, Pieczonka CM, et al. Absorbable hydrogel spacer use in prostate radiotherapy: a comprehensive review of phase 3 clinical trial published data. Urology. 2018 May;115:39-44.

Kim KS, Choi S, Choi YS et al. Comparison of efficacy and safety between a segmental thermos-expandable metal alloy spiral stent (Memokath 051) and a self-expandable covered metallic stent (UVENTA) in the management of ureteral obstructions. J Laparoendosc Adv Surg Tech. 2014; 24(8):550-5.

Kimata R, Nemoto K, Tomita Y, et al. Efficacy of a thermo expandable metallic prostate stent (Memokath) in elderly patients with urethral obstruction requiring long-term management with urethral Foley catheters. Geriatr Gerontol. 2015; 15(5):553-558.

Lerner LB, McVary, KT, Barry MJ et al: Management of lower urinary tract symptoms attributed to benign prostatic hyperplasia: AUA Guideline part II, surgical evaluation and treatment . J Urol 2021; 206: 818.

~~Lowrance W, Breau R, Chou R, et al. Advanced prostate cancer: AUA/ASTRO/SUO guideline. 2020. Available at: <https://www.auanet.org/guidelines/advanced-prostate-cancer>. Accessed April 16, 2021May 10, 2022.~~

Mariados N, Sylvester J, Shah D, et al. Hydrogel spacer prospective multicenter randomized controlled pivotal trial: dosimetric and clinical effects of perirectal spacer application in men undergoing prostate image guided intensity modulated radiation therapy. Int J Radiat Oncol Biol Phys. 2015 Aug 1;92(5):971-7.

~~McVary KT, Roehrborn CG, Avins AL, et al. American Urological Association (AUA) Guideline: management of benign prostatic hyperplasia (BPH). Updated 2010. Reviewed and validity reconfirmed 2014.~~

McVary KT, Gittelman MC, Goldberg KA, et al. Final 5-year Outcomes of the Multicenter Randomized Sham-Controlled Trial of a Water Vapor Thermal Therapy for Treatment of Moderate to Severe Lower Urinary Tract Symptoms Secondary to Benign Prostatic Hyperplasia. J Urol. 2021 Sep;206(3):715-724.

UnitedHealthcare, Inc. ("UHC") Proprietary and Confidential Information: The information contained in this document is confidential, proprietary and the sole property of UHC. The recipient of this information agrees not to disclose or use it for any purpose other than to facilitate UHC's compliance with applicable State Medicaid contractual requirements. Any other use or disclosure is strictly prohibited and requires the express written consent of UHC.

McVary KT, Roehrborn CG. Three-Year Outcomes of the Prospective, Randomized Controlled Rezūm System Study: Convective Radiofrequency Thermal Therapy for Treatment of Lower Urinary Tract Symptoms Due to Benign Prostatic Hyperplasia. Urology. 2018 Jan;111:1-9.

McVary KT, Gange SN, Gittelman MC, et al. Minimally Invasive Prostate Convective Water Vapor Energy Ablation: A Multicenter, Randomized, Controlled Study for the Treatment of Lower Urinary Tract Symptoms Secondary to Benign Prostatic Hyperplasia. J Urol. 2016 May;195(5):1529-1538.

McWilliams JP, Bilhim TA, Carnevale FC, et al. Society of Interventional Radiology Multi-society Consensus Position Statement on Prostatic Artery Embolization for Treatment of Lower Urinary Tract Symptoms Attributed to Benign Prostatic Hyperplasia: From the Society of Interventional Radiology, the Cardiovascular and Interventional Radiological Society of Europe, Société Française de Radiologie, and the British Society of Interventional Radiology: Endorsed by the Asia Pacific Society of Cardiovascular and Interventional Radiology, Canadian Association for Interventional Radiology, Chinese College of Interventionalists, Interventional Radiology Society of Australasia, Japanese Society of Interventional Radiology, and Korean Society of Interventional Radiology. J Vasc Interv Radiol. 2019 May;30(5):627-637

Memokath™. Available at: <https://pnnmedical.com/memokath/>. Accessed May 10, 2021/2022.

Miller LE, Chughtai B, McVary K, et al. Water vapor thermal therapy for lower urinary tract symptoms secondary to benign prostatic hyperplasia: Systematic review and meta-analysis. Medicine (Baltimore). 2020 Jul 24;99(30):e21365.

National Comprehensive Cancer Network (NCCN) Clinical Practice Guidelines in Oncology. Prostate Cancer, v2v34.2021 2022. February 17, 2021/May/January 10, 2022.

National Institute for Health and Care Excellence (NICE). Medical technologies guidance [MTG35]. Memokath-051 stent for ureteric obstruction. February 2018.

National Institute for Health and Care Excellence (NICE). Medical technologies guidance [MTG49]. Rezūm for treating lower urinary tract symptoms secondary to benign prostatic hyperplasia. June 2020.

National Institute for Health and Care Excellence (NICE). Interventional procedures guidance [IPG611]. Prostate artery embolization for lower urinary tract symptoms caused by benign prostatic hyperplasia. April 2018.

Paetkau O, Gagne IM, Pai HH, et al. Maximizing rectal dose sparing with hydrogel: A retrospective planning study. J Appl Clin Med Phys. 2019 Mar 19.

~~Parsons JK, Barry MJ, Dahm P, et al. American Urological Association. Benign prostatic hyperplasia: surgical management of benign of benign prostatic hyperplasia/lower urinary tract symptoms. Published 2018, amended 2019, 2020.~~

Pacella CM, Patelli G, Iapicca G, et al. Transperineal laser ablation for percutaneous treatment of benign prostatic hyperplasia: a feasibility study. Results at 6 and 12 months from a retrospective multi-centric study. Prostate Cancer Prostatic Dis. 2020 Jun;23(2):356-363.

Petrelli F, Vavassori I, Coinu A, et al. Radical prostatectomy or radiotherapy in high-risk prostate cancer: a systematic review and metaanalysis. Clin Genitourin Cancer. 2014 Aug;12(4):215-24.

UnitedHealthcare, Inc. ("UHC") Proprietary and Confidential Information: The information contained in this document is confidential, proprietary and the sole property of UHC. The recipient of this information agrees not to disclose or use it for any purpose other than to facilitate UHC's compliance with applicable State Medicaid contractual requirements. Any other use or disclosure is strictly prohibited and requires the express written consent of UHC.

Pimentel MA, Yassaie O, Gilling P. Urodynamic outcomes after aquablation. Urology. 2019 Apr; 126:165-170.

Pinkawa M, Berneking V, Schlenter M, et al. Quality of life after radiation therapy for prostate cancer with a hydrogel spacer: 5-year results. Int J Radiat Oncol Biol Phys. 2017a;99(2):374-377.

Pinkawa M, Berneking V, Konig L, et al. Hydrogel injection reduces rectal toxicity after radiotherapy for localized prostate cancer. Strahlenther Onkol. 2017b;193(1):22-28.

Plante M, Gilling P, Barber N, et al. Symptom relief and anejaculation after aquablation or transurethral resection of the prostate: subgroup analysis from a blinded randomized trial. BJU Int. 2018.

Porpiglia F, Fiori C, Bertolo R, et al. 3-Year follow-up of temporary implantable nitinol device implantation for the treatment of benign prostatic obstruction. BJU Int. 2018 Jul;122(1):106-112.

Reale G, Cimino S, Bruno G, et al. "Aquabeam® System" for benign prostatic hyperplasia and LUTS: birth of a new era. A systematic review of functional and sexual outcome and adverse events of the technique. Int J Impot Res. 2019;31(6):392-399.

Roehrborn CG, Barkin J, Gange SN, et al. Five-year results of the prospective randomized controlled prostatic urethral L.I.F.T. study. Can J Urol. 2017 Jun;24(3):8802-8813.

Sajan A, Mehta T, Desai P, et al. Minimally invasive treatments for benign prostatic hyperplasia: systematic review and network meta-analysis. J Vasc Interv Radiol. 2022 Apr;33(4):359-367.e8.

~~Sanda MC, Chen RC, Crispino T, et al. Clinically localized prostate cancer: AUA/ASTRO/SUO guideline. 2017. Available at: https://www.astro.org/uploadedFiles/MAIN_SITE/Patient_Care/Clinical_Practice_Statements/Content_Pieces/ClinicallyLocalizedProstateCancer.pdf. Accessed May 24, 2022.~~

Shore ND, Dineen MK, Saslawsky MJ, et al. A temporary intraurethral prostatic stent relieves prostatic obstruction following transurethral microwave thermotherapy. J Urol. 2007 Mar;177(3):1040-6.

Taggar AS, Charas T, Cohen GN, et al. Placement of an absorbable rectal hydrogel spacer in patients undergoing low-dose-rate brachytherapy with palladium-103. Brachytherapy. 2018;17(2):251-258.

Vaggers S, Rai BP, Chedgy ECP, et al. Polyethylene glycol-based hydrogel rectal spacers for prostate brachytherapy: a systematic review with a focus on technique. World J Urol. 2020 Aug 25.

Valerio M, Cerantola Y, Eggner SE, et al. New and established technology in focal ablation of the prostate: A systematic review. Eur Urol. 2017 Jan;71(1):17-34.

van Luijtelaar A, Greenwood BM, Ahmed HU, et al. Focal laser ablation as clinical treatment of prostate cancer: report from a Delphi consensus project. World J Urol. 2019 Oct;37(10):2147-2153.

Varca V, Benelli A, Perri D, et al. Laparoscopic radical prostatectomy in patients with high-risk prostate cancer: feasibility and safety. Results of a Multicentric Study. J Endourol. 2018 Sep 12;32(9):843-851.

Wu SY, Boreta L, Wu A, et al. Improved rectal dosimetry with the use of SpaceOAR during high dose-rate brachytherapy. Brachytherapy. 2018;17(2):259-264.

UnitedHealthcare, Inc. ("UHC") Proprietary and Confidential Information: The information contained in this document is confidential, proprietary and the sole property of UHC. The recipient of this information agrees not to disclose or use it for any purpose other than to facilitate UHC's compliance with applicable State Medicaid contractual requirements. Any other use or disclosure is strictly prohibited and requires the express written consent of UHC.

Xiang P, Guan D, Du Z, et al. Efficacy and safety of prostatic artery embolization for benign prostatic hyperplasia: a systematic review and meta-analysis of randomized controlled trials. Eur Radiol. 2021 Jan 15.

Xu C, Zhang G, Wang JJ, et al. Safety and efficacy of prostatic artery embolization for large benign prostatic hyperplasia in elderly patients. J Int Med Res. 2021 Jan;49(1):300060520986284.

Yeh J, Lehrich B, Tran C, et al. Polyethylene glycol hydrogel rectal spacer implantation in patients with prostate cancer undergoing combination high-dose-rate brachytherapy and external beam radiotherapy. Brachytherapy. 2016 Feb 4. pii: S1538-4721(15) 00633-9.

Zumstein V, Betschart P, Vetterlein MW, et al. Prostatic Artery Embolization versus Standard Surgical Treatment for Lower Urinary Tract Symptoms Secondary to Benign Prostatic Hyperplasia: A Systematic Review and Meta-analysis. Eur Urol Focus. 2019 Nov;5(6):1091-1100.

DRAFT

UnitedHealthcare, Inc. ("UHC") Proprietary and Confidential Information: The information contained in this document is confidential, proprietary and the sole property of UHC. The recipient of this information agrees not to disclose or use it for any purpose other than to facilitate UHC's compliance with applicable State Medicaid contractual requirements. Any other use or disclosure is strictly prohibited and requires the express written consent of UHC.

Policy History/Revision Information

DRAFT

UnitedHealthcare, Inc. ("UHC") Proprietary and Confidential Information: The information contained in this document is confidential, proprietary and the sole property of UHC. The recipient of this information agrees not to disclose or use it for any purpose other than to facilitate UHC's compliance with applicable State Medicaid contractual requirements. Any other use or disclosure is strictly prohibited and requires the express written consent of UHC.

Date

Summary of Changes

DRAFT

TBD

Coverage Rationale

- Added language to indicate:
 - High-energy water vapor thermotherapy for the treatment of benign prostatic hyperplasia (BPH) is proven and medically necessary when performed according to the following U.S. FDA labeled indications, contraindications, warnings, and precautions:
 - To relieve symptoms, obstructions and reduce prostate tissue in men 50 years of age or older with a prostate volume \geq 30 cm and \leq 80 cm; or
 - Treatment of the prostate with hyperplasia of the central zone and/or a median lobe
 - The following procedures are unproven and not medically necessary:
 - Cryoablation of the prostate for initial treatment of prostate cancer
 - Transperineal laser ablation (TPLA)
 - The following procedures are unproven and not medically necessary for all other indications [not listed in the policy as proven and medically necessary] due to insufficient evidence of safety and/or efficacy:
 - Cryoablation of the prostate
 - High-energy water vapor thermotherapy
 - Prostatic urethral lift (PUL)
 - Surgical prostatectomy
 - Transurethral ablation of the prostate
- Replaced language indicating:
 - "Cryoablation of the prostate is proven and medically necessary in certain circumstances" with "cryoablation of the prostate is proven and medically necessary for recurrent prostate cancer diagnosed by biopsy"
 - "Surgical prostatectomy is proven and medically necessary in certain circumstances" with "surgical prostatectomy is proven and medically necessary in certain circumstances, including for some individuals with very high risk or recurrent prostate cancer"
 - "Prostatic urethral lift (PUL) is proven and medically necessary when performed according to the following U.S. FDA labeled indication" with "prostatic urethral lift (PUL) is proven and medically necessary when performed according to the following U.S. FDA labeled indications, contraindications, warnings, and precautions"
 - "Focal laser ablation is unproven and not medically necessary" with "transperineal focal laser ablation is unproven and not medically necessary"

Applicable Codes

- Added CPT code 0714T
- Added language to indicate CPT codes 0421T, 0582T, 0655T, 0714T, and 53852 are not on the State of Louisiana Fee Schedule and therefore are not covered by the State of Louisiana Medicaid Program

Supporting Information

- Updated Description of Services, Clinical Evidence, FDA, and References sections to reflect the most current information
- Archived previous policy version CS334LA.A

UnitedHealthcare, Inc. ("UHC") Proprietary and Confidential Information: The information contained in this document is confidential, proprietary and the sole property of UHC. The recipient of this information agrees not to disclose or use it for any purpose other than to facilitate UHC's compliance with applicable State Medicaid contractual requirements. Any other use or disclosure is strictly prohibited and requires the express written consent of UHC.

Instructions for Use

This Medical Policy provides assistance in interpreting UnitedHealthcare standard benefit plans. When deciding coverage, the federal, state or contractual requirements for benefit plan coverage must be referenced as the terms of the federal, state or contractual requirements for benefit plan coverage may differ from the standard benefit plan. In the event of a conflict, the federal, state or contractual requirements for benefit plan coverage govern. Before using this policy, please check the federal, state or contractual requirements for benefit plan coverage. UnitedHealthcare reserves the right to modify its Policies and Guidelines as necessary. This Medical Policy is provided for informational purposes. It does not constitute medical advice.

UnitedHealthcare may also use tools developed by third parties, such as the InterQual® criteria, to assist us in administering health benefits. The UnitedHealthcare Medical Policies are intended to be used in connection with the independent professional medical judgment of a qualified health care provider and do not constitute the practice of medicine or medical advice.