

National Imaging Associates, Inc.*	
Clinical guidelines PASSIVE TREATMENT	Original Date: November 2 , 2015
Physical Medicine – Clinical Decision Making	Last Revised Date: Novem October December 202 10
Guideline Number: NIA_CG_604	Implementation Date: July 202 21

Policy Statement

This organization does not support the use of multiple passive treatments for the care of musculoskeletal pain within the scope of network practitioners. Most passive treatments have similar physiological effects related to pain control and reduction of inflammation. The use of treatments with duplicative physiological effects is unnecessary and inappropriate. Multiple passive treatments have not been shown to improve or accelerate patient health outcomes.

Scope

Physical medicine participating network practitioners, including rendering chiropractors, physical therapists, occupational therapists, speech therapists, and therapist assistants as applicable. This policy also applies to out of network practitioners as dictated by the health plan.

Definitions

Modality

Modality is defined as any group of agents that may include thermal, acoustic, radiant, mechanical, or electrical energy to produce physiologic changes in tissues of therapeutic purposes. Modalities affect tissue at the cellular level.

Multiple Modalities

Multiple modalities are defined as the use of and/or billing of two or more physical medicine modalities each visit or during the same session to the same region.

Passive Treatment

Treatment that is applied by the provider or in a clinical setting and does not involve active participation by the patient.

* National Imaging Associates, Inc. (NIA) is a subsidiary of Magellan Healthcare, Inc.

Procedure

Procedure is a service provided to increase the functional abilities in self-care, mobility, or safety.

The following is a list of procedures and modalities considered to be passive treatment:

- Thermal and light therapy – Hot/cold (97010), diathermy (97024), microwave (97020), infrared (97026), ultraviolet (97028), ultrasound (US) (97035), paraffin bath (97018), and whirlpool (97022).
- Electrical therapy – High volt, low volt, interferential current, transcutaneous electrical nerve stimulation (TENS) (97014 and 97032).
- Mechanical – mechanically assisted and often a sustained pull of the spine or limb, such as traction (97012).
- Therapeutic massage and manual therapy (97124 and 97140) ~~—M—~~ Manual therapy includes Active Release Technique, trigger point therapy, myofascial release, mobilization/manipulation, manual lymphatic drainage, and manual traction.

Appropriate Use of Passive Treatment

- Passive treatment modalities may be utilized in the initial period of an episode of treatment or exacerbation of a sub-acute or chronic condition for pain control, reduction of inflammation, or reduction of muscle spasm. As a condition progresses, passive care should be replaced by active treatment modalities, such as therapeutic exercise. Insufficient evidence exists to support the continued use of passive treatment as a means for improved clinical outcomes.
- Passive treatment is considered to be clinically appropriate and/or necessary in the conservative management of neuromusculoskeletal conditions when:
 - There are no contraindications to the intervention
 - Self-administration is implausible or places the patient at risk of harm
 - Used primarily during the initial period of an episode of treatment
 - Used to support an active care approach (i.e., therapeutic exercise)
 - Used for a particular condition for which there is an evidence-basis of significant benefit
- Passive treatment is considered NOT to be clinically appropriate and/or necessary when:
 - Patient safety is jeopardized by the application of the modality

- The treatment can safely and effectively be administered by the patient or another individual
- Used during a course of treatment, which continues beyond the initial period
- Used as the primary or sole therapy
- Greater than one passive treatment is used involving the same body region(s)
- Used largely for the comfort and convenience of the patient
- Used as part of the routine office protocol

Exclusions

- The use of chiropractic manipulation (98940-98943) is not considered a duplication of service or physiological effect when used in conjunction with passive ~~treatment, treatment,~~ except for the following:
 - The National Correct Coding Initiative (NCCI) edits require that the manual therapy techniques be performed in a separate anatomic site than the chiropractic adjustments in order to be reimbursed separately.

BACKGROUND

The preponderance of evidence appears to support either a lack of efficacy or insufficient data to make a judgment on benefit for the modalities evaluated. When a positive outcome was described, the reported treatment effects were modest. Similarly, the duration of treatment effectiveness was typically reported as short (2 weeks to 2 months). Most international guidelines recommend these interventions should only be used reservedly based upon individual circumstances, and not as a principle component of a treatment regime.

The use of passive modalities in the treatment of neuromusculoskeletal conditions presents the inherent risk of promoting passive dependence. It is the responsibility of the treating practitioner to judiciously apply passive modalities and encourage active patient participation in the treatment plan. Passive treatment is generally viewed as appropriate when used for a short period of time and in conjunction with ~~an~~ active care.

Low Back Pain and Passive Interventions

[A review on non-pharmacological therapies for acute and chronic LBP by the American Pain Society and the American College of Physicians concluded that therapies with good evidence of moderate efficacy for chronic or sub-acute LBP are cognitive-behavioral therapy, exercise, spinal manipulation, and inter-disciplinary rehabilitation.](#)¹

[Studies suggest that spinal manipulation may provide modest pain relief and improved function for patients with acute low back pain \(pain that has come on within the last four](#)

weeks) or chronic (longer-term) low back pain and generally appears to be safe.² A meta-analysis of 26 RCTs report statistically significant improvement in both pain and function for patients with LBP who received SMT with only a reporting of minor transient adverse events, including muscle stiffness, pain, and headache.³

Surface electrical muscle stimulators (direct or alternating current, not high-voltage galvanic current) are considered experimental and investigational for the management of idiopathic scoliosis because there is inadequate evidence of its effectiveness and safety in the peer-reviewed published medical literature.⁴

4

For patients with low back pain, the use of transcutaneous electrical nerve stimulation (TENS) is not a recommended intervention.^{5, 6} Green et al found that transcutaneous electrical nerve stimulation (TENS) offers no significant benefit for chronic low back pain particularly concerning multiple disability and quality of life measures, but it does offer a small benefit in pain reduction compared with sham treatment. Their study suggests that TENS may be a useful adjunct in select patients for pain control to reduce the need for medication.⁷ Likewise, a 2017 study of 127 patients with LPB split into five different comparison groups, including TENS, or a control concluded that the “TENS currents and high voltage [electrical stimulation] were helpful, but not as effective. The use of diadynamic currents appears to be useless.”⁸

Guidelines on treatment of low back pain from the National Collaborating Centre for Primary Care found insufficient evidence for the use of interferential stimulation in LBP and recommended against its use for that indication.^{9, 10}

~~A review on non-pharmacological therapies for acute and chronic LBP by the American Pain Society and the American College of Physicians concluded that therapies with good evidence of moderate efficacy for chronic or sub-acute LBP are cognitive behavioral therapy, exercise, spinal manipulation, and inter-disciplinary rehabilitation (Qaseem, 2017).¹~~

~~Studies suggest that spinal manipulation may provide modest pain relief and improved function for patients with acute low back pain (pain that has come on within the last four weeks) or chronic (longer-term) low back pain and generally appears to be safe (Chou, 2018).³~~

A Cochrane review of 32 RCTs involving 2762 participants found that “traction, either alone or in combination with other treatments, has little or no impact on pain intensity, functional status, global improvement and return to work among people with LBP. There is only limited-quality evidence from studies with small sample sizes and moderate to high risk of bias. The effects shown by these studies are small and are not clinically relevant” (Wegner, 2013).¹¹ Similarly, the study by Alrwaily and colleagues found that the use of traction in the treatment of low back pain is inconclusive,¹² and Chou et al (2016) found that traction has little or no impact on pain intensity, functional status, or global improvement.¹³

No high-quality evidence was found to support the use of ultrasound for improving pain or quality of life in patients with non-specific chronic LBP. There is some evidence that therapeutic ultrasound has a small effect on improving low-back function in the short term, but this benefit is unlikely to be clinically important. Evidence from comparisons between other treatments and therapeutic ultrasound for chronic LBP were indeterminate and generally of low quality (Analan 2015).¹⁴ The current evidence does not support the use of therapeutic ultrasound in the management of chronic LBP (Ebadi, 2020).¹⁵

The Cochrane Back and Neck Group reported little confidence that massage is an effective treatment for LBP. Acute, sub-acute, and chronic LBP had improvements in pain outcomes with massage only in the short-term follow-up. Functional improvement was observed in participants with sub-acute and chronic LBP when compared with inactive controls, but only for the short-term follow-up (Furlan, 2015).¹⁶

~~There are insufficient data to draw firm conclusion on the clinical effect of back schools, low-level laser therapy, patient education, massage, traction, superficial heat/cold, and lumbar supports for chronic low back pain (Van Middelkoop, 2011).¹⁷~~

~~A number of nonpharmacological, noninvasive treatments for low back pain are associated with small to moderate, primarily short-term effects on pain versus placebo, sham, wait list, or no treatment. Effects on function are generally smaller than effects on pain. More research is needed to understand optimal selection of treatments, effective combinations, and sequencing of treatments, and effectiveness of treatments for radicular low back pain (Chou, 2017).¹⁸~~

~~Guidelines on treatment of low back pain from the National Collaborating Centre for Primary Care found insufficient evidence for the use of interferential stimulation in LBP and recommended against its use for that indication (Savigny, 2009).^{9, 10}~~

~~Green et al found that transcutaneous electrical nerve stimulation (TENS) offers no significant benefit for chronic low back pain particularly concerning multiple disability and quality of life measures, but it does offer a small benefit in pain reduction compared with sham treatment. Their study suggests that TENS may be a useful adjunct in select patients for pain control to reduce the need for medication (Green, 2016).⁷~~

~~Alrwaily et al found that the use of traction in the treatment of low back pain is inconclusive (Alrwaily, 2018).¹²~~

~~Chou et al (2016)¹² found that traction has little or no impact on pain intensity, functional status, or global improvement.¹⁵~~

A number of nonpharmacological, noninvasive treatments for low back pain are associated with small to moderate, primarily short-term effects on pain versus placebo, sham, wait list, or no treatment. Effects on function are generally smaller than effects on pain. More research is needed to understand optimal selection of treatments, effective combinations, and

sequencing of treatments, and effectiveness of treatments for radicular low back pain.¹⁷

There are insufficient data to draw firm conclusion on the clinical effect of back schools, low-level laser therapy, patient education, massage, traction, superficial heat/cold, and lumbar supports for chronic low back pain.¹⁸

Neck Pain and Passive Interventions

No trials at low risk of bias support the use of traction, stretching, or ultrasound therapy for chronic neck pain. However, Yang et al found that intermittent cervical traction (ICT) may have short-term pain relief effects, but generally the information regarding ICT is inconclusive (Yang, 2017).¹⁹ Likewise, a 2018 systematic review reported “some support to the use of the mechanical and manual traction for CR [cervical radiculopathy] in addition to other physical therapy procedures for pain reduction, but yielding lesser effects on function/disability”, but the reviewers note a lack of homogeneity in diagnostic criteria among the included studies.^{20,}

21

The use of traction for a herniated cervical disc with radiculopathy who completed therapy with a combination treatment of traction and exercise demonstrated good or excellent outcomes with some participants even avoiding surgery.

Low quality evidence suggests clinically important long-term improvements in neck pain, function/disability, and global perceived effect, when manual therapy and exercise are compared to no treatment. High quality evidence suggests greater short-term pain relief than exercise alone,²² but no long-term differences across multiple outcomes for (sub) acute/chronic neck pain with or without cervicogenic headache. Moderate quality evidence supports this treatment combination for pain reduction and improved quality of life, over manual therapy alone for chronic neck pain and suggests greater short-term pain reduction when compared to traditional care for acute whiplash. Evidence regarding radiculopathy was sparse (Miller, 2010).²³

Coulter et al found that there is low to moderate quality evidence that various types of manipulation and/or mobilization will reduce pain and improve function for chronic nonspecific neck pain compared to other interventions, however a multimodal approaches have the greatest potential impact (Coulter, 2019).²⁴ Likewise, Díaz-Pulido et al report that manual therapy resulted in significantly better improvement in patients with chronic neck pain at post-intervention and at 6-month follow-up as compared to patients undergoing TENS therapy.²⁵

Both stretching exercise and manual therapy considerably decreased neck pain and disability in women with non-specific neck pain. The difference in effectiveness between the two treatments was minor. Low-cost stretching exercises can be recommended in the first instance, as an appropriate therapy intervention to relieve pain, at least in the short-term (Ylinen, 2007).²⁶

Combining different forms of manual therapy with exercise is better than manual therapy or exercise alone ~~(Hidalgo, 2017).~~²⁷

Manual trigger point treatment of head and neck muscles may reduce frequency, intensity, and duration of attacks in tension-type headaches and migraine headaches, but the quality of evidence according to GRADE approach was very low for the presence of few studies, high risk of bias, and imprecision of results ~~(Falsiroli, 2018).~~²⁸

There is a linear dose-response relationship between SMT visits and days with cervicogenic headache (CGH). For the highest and most effective dose of 18 SMT visits, CGH days were reduced by half and about 3 more days per month than for the light-massage control ~~(Haas, 2018).~~²⁹

For the treatment of the diagnostic label Non-Specific Neck Pain **(NP)**, strong evidence of efficacy was only found for multimodal care (manipulation/mobilization and supervised exercises) ~~(Tsakitzidis, 2013).~~³⁰ A prospective double-blind randomized controlled trial examining the effects of multimodal care on patients with NP, plus/minus addition of neck-specific aerobic exercise, showed statistically significant reduction in both NP and cervicogenic headache.³¹

Interferential current **(IFC)** therapy is effective in the treatment of chronic neck pain patients. However, the results of clinical trials, to date, have been conflicting regarding whether IFC ~~it~~ has no additional benefit or superiority over neck stabilization exercises ~~(Yesil, 2018).~~ Additional research is required.^{32, 33}

In regards to chronic mechanical neck pain, stabilization exercises with or without connective tissue massage (CTM) might be a useful treatment, however stabilization exercises with CTM are superior in improving pain intensity at night, pressure pain threshold, state anxiety, and mental health compared to stabilization exercises alone ~~(Celenay, 2015).~~³⁴

Thrust manipulation and non-thrust mobilization was less effective when performed alone than when combined with therapeutic exercises for mechanical neck pain with or without headaches.

Cervical traction has not been shown to be effective in the treatment of neck pain.³⁵ A meta-analysis of randomized controlled trials showed that the use of intermittent cervical traction for treating neck pain did not differ significantly from a placebo during the follow-up period after treatment.¹⁹ ~~-(Chou, 2018; UpToDate 2018).~~

TMJ and Passive Interventions

No high-quality evidence was found, indicating that there is great uncertainty about the effectiveness of exercise and manual therapy for treatment of temporomandibular joint dysfunction.

Shoulder Pain and Passive Interventions

For adults with nonspecific shoulder pain of variable duration, cervicothoracic spinal manipulation and mobilization, in addition to usual care, may improve self-perceived recovery compared to usual care alone. For adults with subacromial impingement syndrome of variable duration, neck mobilization in addition to a multimodal shoulder program of care, provides no added benefit.³⁶ ~~(Southerst, 2015)~~

For patients with rotator cuff tendinopathy, based on low to moderate-quality evidence, manual therapy may decrease pain; however, it is unclear whether it can improve function ~~(Desjardins-Charbonneau, 2015).~~³⁷ One meta-analysis notes, “When combined with exercise, manual therapy was superior to exercise alone, but only at the shortest follow-up”.³⁸

There was little evidence that active therapeutic ultrasound is more effective than placebo ultrasound for treating people with pain or a range of musculoskeletal injuries or for promoting soft tissue healing.^{39, 40} A Cochrane Database Review of the use of ultrasound with patients suffering from rotator cuff disease notes no difference between the US and control groups in long-term follow-up (nine months).⁴¹ ~~In regard to patients with rotator cuff disease~~ Additionally, a double-blind study by Analan et al showed that ultrasound does not provide additional benefit to the physiotherapy treatment regimen concerning pain, function, and isokinetic shoulder rotator cuff strength ~~(Analan, 2015).~~¹⁴

Deep tissue friction massage has been shown to be beneficial in improving function and range of motion in supraspinatus tendinitis patients.⁴² ~~(Jindal, 2015).~~

Hip Pain and Passive Interventions

The best available evidence indicates that exercise therapy (whether land-based or water-based) is more effective than minimal control in managing pain associated with hip osteoarthritis (OA) in the short term. Larger high-quality randomized controlled trials (RCT) are needed to establish the effectiveness of exercise and manual therapies in the medium and long term ~~(Beumer, 2016).~~⁴³⁻⁴⁵

Knee Pain and Passive Interventions

Chaves found that ~~in order for~~ deep friction massage (DFM) significantly decreased pain intensity over time in individuals with patellar tendinopathy, regardless of the pressure used. The authors note, “DFM induces an immediate reduction in pain intensity upon palpation,... Notwithstanding, the reader should take into account the small sample size and the caution needed in the results' interpretation.”⁴⁶ ~~to achieve a therapeutic effect in patients with patellar tendinopathy a 1.1 kg/cm² ratio of force needs to be applied (Chaves, 2018).~~⁴⁶

Studies using therapeutic ultrasound for the treatment or management of knee pain have had conflicting results to date. Ultrasound provided no additional benefit in improving pain and function in addition to exercise training in the management of knee osteoarthritis. In a

meta-analysis of the use of ultrasound (US) therapy in treating myofascial pain syndrome (MPS), the researchers conclude, “Owing to the high risk of bias and the across-trial heterogeneity of the studies, the current evidence is not clear enough to support ~~ultrasound~~ US as an effective method to treat ~~myofascial pain syndrome~~ MPS (Xia, 2017).”⁴⁷ Another meta-analysis notes a statistically significant decrease in pain for patients with knee osteoarthritis undergoing therapeutic ultrasound therapy as compared to the individuals in a sham ultrasound control group.⁴⁸

~~The use of ultrasound therapy in patients with knee osteoarthritis had no significant difference when compared to the sham group.~~

For patients with knee osteoarthritis, the use of transcutaneous electrical nerve stimulation (TENS) may provide pain relief; however, the evidence is limited.⁴⁹⁻⁵¹

Ankle Pain and Passive Interventions

For adults with grade I-II ankle sprains of variable duration, lower extremity mobilization, in addition to home exercise and advice, provides greater short-term improvements in activities and function over home exercise and advice alone ~~(Southerst, 2015).~~³⁶

For patients with acute exercise-induced achilles tendinopathy, low level laser therapy may be helpful in reducing inflammation and pain.⁵²

In patients with achilles tendinopathy of < 3 months, the use of iontophoresis was shown to help with pain. However, the control group still demonstrated improvements with the use of a comprehensive rehabilitation program.⁵³

Chronic Pain and Passive Interventions

“Passive therapy (those treatment modalities that do not require energy expenditure on the part of the patient) can provide short-term relief during chronic pain flare-ups and is directed at controlling symptoms such as pain, inflammation, and swelling. Passive therapies may be useful over the short term but have limited benefit for chronic pain conditions overall.”⁵⁴

Electrical Stimulation and Laser Therapy

In a systematic review and meta-analysis, Fuentes analyzed the available information regarding the efficacy of interferential therapy in the management of musculoskeletal pain. Interferential current alone was not significantly better than placebo or other therapy at discharge or follow-up ~~(Fuentes, 2010).~~⁵⁵

There is a paucity of evidence in the peer-reviewed literature regarding the effectiveness of high-voltage, pulsed current treatments in humans as a means of controlling edema and post-traumatic pain; thus, a clear evidence base has not yet been established.

Scientific evidence in the peer review literature is lacking regarding the use, safety, improvement, or effectiveness on health outcomes for light emitting diode (infrared) therapy.

In a systematic review and meta-analysis, Song, et al (2018) found that high intensity laser therapy is able to significantly reduce pain and disability in patients with back and neck pain.⁵⁶

Documentation Requirements:

The treatment plan or plan of care must include the clinical rationale for each service, a description of the service, the area of the body the service will be provided, goals for each service, and a time component, if indicated.

Applicable contraindications for passive modalities (e.g., ultrasound therapy) should be considered.

POLICY HISTORY ~~SUMMARIES:~~

<u>Date</u>	<u>Summary</u>
November <u>December 2021</u>	• <u>Added “General Information” statement</u> • <u>Expansion of the Background section to strengthen the presentation of the evidence-base in support of the indications.</u> • <u>Minor copyediting changes</u>
<u>October 2020</u>	• <u>Modified the definition of “Modality” to include manual/massage therapy</u> • <u>Modified “Appropriate Use of Passive Treatment” by clarifying when passive modalities may be utilized</u> • <u>Modified “Appropriate Use of Passive Treatment” by clarifying passive modalities as not clinically appropriate and/or necessary by adding effectively be performed by the patient or another individual</u> • <u>Modified the “Definitions” section to better distinguish between modalities and passive treatment</u>
<u>January 2020</u>	• <u>No changes to indications</u> • <u>Expansion of the Background section to strengthen the presentation of the evidence-base in support of the indications.</u> • <u>Minor copyediting changes</u>

July 2019

- Older references updated or omitted as appropriate.
- Background information expanded to reflect scope of current evidence base.

July 2019

- ~~Older references updated or omitted as appropriate.~~
- ~~Background information expanded to reflect scope of current evidence base.~~

January 2020

- ~~No changes to indications~~
- ~~Expansion of the Background section to strengthen the presentation of the evidence-base in support of the indications.~~
- ~~Minor copyediting changes~~

October 2020

- ~~Modified the definition of “Modality” to include manual/massage therapy~~
- ~~Modified “Appropriate Use of Passive Treatment” by clarifying when passive modalities may be utilized~~
- ~~Modified “Appropriate Use of Passive Treatment” by clarifying passive modalities as not clinically appropriate and/or necessary by adding effectively be performed by the patient or another individual~~
- ~~Modified the “Definitions” section to better distinguish between modalities and passive treatment~~

REFERENCES

- Alrwaily M, Almutiri M, Schneider M. Assessment of variability in traction interventions for patients with low back pain: a systematic review. *Chiropr Man Therap*. 2018; 26:35.
- Analan PD, Leblebici B, Adam M. Effects of therapeutic ultrasound and exercise on pain, function, and isokinetic shoulder rotator cuff strength of patients with rotator cuff disease. *J Phys Ther Sci*. 2015;27(10):3113-7.
- Armijo-Olivo S, Pitance L, Singh V, et al. Effectiveness of manual therapy and therapeutic exercise for temporomandibular disorders: systematic review and meta-analysis. *Phys Ther*. 2016; 96(1):9-25.
- Babatunde OO, Jordan JL, Van der Windt DA, et al. Effective treatment options for musculoskeletal pain in primary care: A systematic overview of current evidence. *PLoS ONE*. 2017; 12(6):e0178621.
- Baez S, Hoch J, Uhl T. The effectiveness of cervical traction and exercise in decreasing neck and arm pain for patients with cervical radiculopathy: A critically appraised topic. *Int J Athl Ther Train*. 2017; 22(5):4-11.
- Bagheripour B, Kamyab M, Azadinia F. The efficacy of a home mechanical traction unit for patients with mild to moderate cervical osteoarthritis: A pilot study. *Med J Islam Repub Iran*. 2016; 30:386.
- Bryans R, Decina P, Descarreaux M, et al. Evidence-based guidelines for the chiropractic treatment of adults with neck pain. *J Manipulative Physiol Ther*. 2014 Jan;37(1):42-63. doi: 10.1016/j.jmpt.2013.08.010. Epub 2013 Nov 19.
- Beumer L, Wong J, Warden SJ, et al. Effects of exercise and manual therapy on pain associated with hip osteoarthritis: a systematic review and meta-analysis. *Br J Sports Med*. 2016; 50(8):458-463.
- Buchbinder R, Tulder MV, Oberg B, et al. Low Back Pain: A Call for Action. *Lancet*. 2018;391(10137):2384-2388.
- Bukhari SR, Shakil Ur Rehman S, Ahmad S, et al. Comparison between effectiveness of mechanical and manual traction combined with mobilization and exercise therapy in patients with cervical radiculopathy. *Pak J Med Sci*. 2016; 32(1):31-34.
- Cai C, Pua Y, Lim K. A clinical prediction rule for classifying patients with low back pain who demonstrate short-term improvement with mechanical lumbar traction. *Euro Spine J*. 2009; 18(4):554-561.

~~Cakir S, Hepguler S, Ozturk C, et al. Efficacy of therapeutic ultrasound for the management of knee osteoarthritis: a randomized, controlled, and double blind study. *Am J Phys Med Rehabil*. 2014; 93(5):405-412.~~

~~Carey TS, Freburger JK, Holmes GM, et al. A long way to go: Practice patterns and evidence in chronic low back pain care. *Spine*. 2009; 34(7):718-724.~~

~~Celenay ST, Kaya DO, Akbayrak T. Cervical and scapulothoracic stabilization exercises with and without connective tissue massage for chronic mechanical neck pain: a prospective, randomized controlled trial. *Man Ther*. 2016; 21:144-50~~

~~Centers for Medicare & Medicaid Services (CMS). Decision memo for transcutaneous electrical nerve stimulation for chronic low back pain (CAG-00429N). CMS, Baltimore, MD; June 8, 2012. <http://www.cms.gov/medicare-coverage-database/details/nca-decision-memo.aspx?NCAId=256>. Accessed April 21, 2017.~~

~~Chaves, Maria Dias. Is there a role for deep friction massage in the management of patellar tendinopathy? Technique characterization and short term clinical outcomes. Doctorate dissertation in Physiotherapy, presented at the Faculty of Sport, University of Porto, 2018.~~

~~Cherian JJ, Jauregui JJ, Leichter AK, et al. The effects of various physical non-operative modalities on the pain in osteoarthritis of the knee. *Bone Joint J*. 2016; 98(B Suppl 1A): 89-94.~~

~~Chou R. Patient education: Low back pain in adults (Beyond the Basics). UpToDate. Waltham, MA: March 21, 2018.~~

~~Chou R, Côté P, Randhawa K, et al. The Global Spine Care Initiative: applying evidence-based guidelines on the non-invasive management of back and neck pain to low- and middle-income communities. *Eur Spine J*. 2018 Sep;27(Suppl 6):851-860. doi: 10.1007/s00586-017-5433-8. Epub 2018 Feb 19.~~

~~Chou R, Deyo R, Friedly J, et al. Nonpharmacologic therapies for low back pain: A systematic review for an American college of physicians clinical practice guideline. *Ann Intern Med*. 2017; 166(7):493-505.~~

~~Chou R, Deyo R, Friedly J, et al. Noninvasive Treatments for Low Back Pain. Rockville (MD): Agency for Healthcare Research and Quality (US); 2016 Feb. Report No.: 16-EHC004-EF.~~

~~Clar C, Tsertsvadze A, Court R, et al. Clinical effectiveness of manual therapy for the management of musculoskeletal and non-musculoskeletal conditions: Systematic review and update of UK evidence report. *Chiropr Man Therap*. 2014; 22(12). doi: 10.1186/2045-709X-22-12.~~

Coulter ID, Crawford C, Hurwitz EL, et al. Manipulation and mobilization for treating chronic low back pain: a systematic review and meta-analysis. *Spine J*. 2018;18(5):866-879.

Coulter I, Crawford CJ, Vernon H, Hurwitz E, Khorsan R, Booth M, Herman PM. Manipulation and mobilization for treating chronic nonspecific neck pain: A systematic review and meta-analysis for an appropriateness panel. *Pain Physician*. 2019; 22:E55-E70.

Crawford C, Boyd C, MA, Paat CF, et al. The impact of massage therapy on function in pain populations: a systematic review and meta-analysis of randomized controlled trials: part I, patients experiencing pain in the general population. *Pain Med*. July 2016; 17(7): 1353-1375.

Damgaard P, Bartels EM, Ris I, et al. Evidence of physiotherapy interventions for patients with chronic neck pain: A systematic review of randomised controlled trials. *ISRN Pain*. 2013; 567175. doi: 10.1155/2013/567175.

Desjardins-Charbonneau A, Roy JS, Dionne CE, et al. The efficacy of manual therapy for rotator cuff tendinopathy: A systematic review and meta-analysis. *J Orthop Sports Phys Ther*. 2015; 45(5):330-350.

Desmeules F, Boudreault J, Roy JS, et al. The efficacy of therapeutic ultrasound for rotator cuff tendinopathy: A systematic review and meta-analysis. *Phys Ther Sport*. 2015; (3):276-284.

Ebadi S, Henschke N, Forogh B, et al. Ultrasound therapy for chronic low back pain. Cochrane Back and Neck Group. 5 July 2020.

Ebadi S, Henschke N, Nakhostin Ansari N, et al. Therapeutic ultrasound for chronic low back pain. *Cochrane Database Syst Rev*. 2014; (3): CD009169.

Falsiroli Maistrello L, Geri T, Gianola S, et al. Effectiveness of Trigger Point Manual Treatment on the Frequency, Intensity, and Duration of Attacks in Primary Headaches: A Systematic Review and Meta-Analysis of Randomized Controlled Trials. *Front Neurol*. 2018;9:254. doi: 10.3389/fneur.2018.00254

Farooq MN, Mohseni-Bandpei MA, Gilani SA, et al. The effects of neck mobilization in patients with chronic neck pain: a randomized controlled trial. *J Bodyw Mov Ther*. January 2018; 22(1):24-31.

Foster NE, Anema JR, Cherkin D, et al. Prevention and treatment of low back pain: evidence challenges, and promising directions. *Lancet*. 2018;391(10137):2368-2383.

Fritz JM, Thackeray A, Brennan G, et al. Exercise only, exercise with mechanical traction, or exercise with over door traction for patients with cervical radiculopathy, with or without

~~consideration of status on a previously described subgrouping rule: A randomized clinical trial. *J Orthop Sports Phys Ther*. 2014; 44(2):45-57.~~

~~Fuentes JP, Armijo OS, Magee DJ, et al. Effectiveness of interferential current therapy in the management of musculoskeletal pain: a systematic review and meta-analysis. *Phys Ther*. 2010; 90(9):1219-38. doi: 10.2522/ptj.20090335.~~

~~Furlan A, Yazdi F, Tsertsvadze A, et al. A systematic review and meta-analysis of efficacy, cost-effectiveness, and safety of selected complementary and alternative medicine for neck and low back pain. *Evid Based Complement Alternat Med*. 2012; 953139. doi:10.1155/2012/953139.~~

~~Furlan AD, Giraldo M, Baskwill A, et al. Massage for low back pain. *Cochrane Database Syst Rev*. 2015; (9): CD001929. doi: 10.1002/14651858.~~

~~Glazov G, Yelland M, Emery J. Low level laser therapy for chronic non-specific low back pain: a meta-analysis of randomised controlled trials. *Acupunct Med*. 2016; 34:328-341.~~

~~Green MW, Clemente Fuentes RJ. Is transcutaneous electrical nerve stimulation (TENS) more effective than placebo for management of chronic low back pain? *Evidence-Based Practice*. 2016; 19(8): 10-11.~~

~~Haas M, Bronfort G, Evans R, et al. Dose-response and efficacy of spinal manipulation for care of cervicogenic headache: a dual-center randomized controlled trial. *Spine J*. 2018. pii: S1529-9430(18)30077-9.~~

~~Hawk C, Minkalis A, Khorsan R, et al. Systematic review of nondrug, nonsurgical treatment of shoulder conditions. *J Manip Physiol Ther*. 2017; 40(5):293-319.~~

~~Hidalgo B, Hall T, Bossert J, et al. The efficacy of manual therapy and exercise for treating non-specific neck pain: a systematic review. *J Back Musculoskelet Rehabil*. 2017; 30(6):1149-1169.~~

~~Isaac Z, Dec KL. Patient education: Neck pain (Beyond the Basics). UpToDate, 01/25/18. <https://www.uptodate.com/contents/neck-pain-beyond-the-basics>~~

~~Jindal JK, Monika M. Efficacy of Conventional Treatment and Eccentric Exercise with and without Deep Transverse Friction Massage in Supraspinatus Tendinitis Patients a Randomized Clinical Trial. *Indian Journal of Physiotherapy and Occupational Therapy*. 2015; 9(3):249-253~~

~~Johnson MI, Paley CA, Howe TE, et al. Transcutaneous electrical nerve stimulation for acute pain. *Cochrane Database of Syst Rev*. 2015; 6:CD006142.~~

~~Kroeling P, Gross A, Graham N, et al. Electrotherapy for neck pain. *Cochrane Database Syst Rev*. 2013; 8: CD004251.~~

~~Kumar S, Beaton K, Hughes T. The effectiveness of massage therapy for the treatment of nonspecific low back pain: a systematic review of systematic reviews. *Int J Gen Med*. 2013; 6:733-41.~~

~~Littlewood C, May S, Walters S. A review of systematic reviews of the effectiveness of conservative interventions for rotator cuff tendinopathy. *British Elbow & Shoulder Society (BESS)*. February 6, 2017; 5(3):151-167. Available at <https://doi.org/10.1111/sae.12009>.~~

~~Martimbianco AC, Torloni MR, Andriolo BNG, et al. Neuromuscular electrical stimulation (NMES) for patellofemoral pain syndrome. *Cochrane Database of Syst Rev*. 2017; (12):CD011289.~~

~~Miller J, Gross A, D'Silva J, et al. Manual therapy and exercise for neck pain: A systematic review. *Man Ther*. 2010; 15(4):334-354.~~

~~Muhammad A, Mohammed A, Michael S. Assessment of variability in traction interventions for patients with low back pain: a systematic review. *Chiropr Man Therap*. 2018; 26:35.~~

~~Negrini S, Minozzi S, Bettany-Saltikov J, et al. Braces for idiopathic scoliosis in adolescents. *Cochrane Database of Syst Rev*. 2010; (1):CD006850.~~

~~Nelson NL, Churilla JR. Massage therapy for pain and function in patients with arthritis: a systematic review of randomized controlled trials. *Am J Phys Med Rehabil*. 2017; 96(9):665-672.~~

~~Page MJ, Green S, Kramer S, et al. Electrotherapy modalities for adhesive capsulitis (frozen shoulder). *Cochrane Database Syst Rev*. 2014; 10:CD011324.~~

~~Page MJ, Green S, Mrocki MA, et al. Electrotherapy modalities for rotator cuff disease. *Cochrane Database Syst Rev*. 2016; (6):CD012225.~~

~~Palmer E, Redavid L. Therapeutic Ultrasound: Clinical Review. Glendale, CA: *Clinical Information Systems: A division of EBSCO Information Services*. 2016.~~

~~Qaseem A, Wilt TJ, McLean RM, et al. Noninvasive Treatments for Acute, Subacute, and Chronic Low Back Pain: A Clinical Practice Guideline from the American College of Physicians. *Ann Intern Med*. 2017; 166(7):514-530.~~

~~Rankin IA, Sargeant H, Rehman H, et al. Low-level laser therapy for carpal tunnel syndrome. *Cochrane Database of Syst Rev*. 2017; 8:CD012765.~~

~~Remmen R, Tsakitzidis G, Dankaerts W, et al. Non-specific neck pain and evidence-based practice. *European Scientific Journal*. 2013; 9(3):1-19.~~

~~Rutjes AWS, Nüesch E, Sterchi R, et al. Transcutaneous electrostimulation for osteoarthritis of the knee. *Cochrane Database of Syst Rev*. 2009; 4:CD002823.~~

~~Salazar AP, Stein C, Marchese RR, et al. Electric stimulation for pain relief in patients with fibromyalgia: a systematic review and meta-analysis of randomized controlled trials. *Pain Physician*. 2017; 20(2):15-25.~~

~~Sandoval MC, Ramirez C, Camargo DM, et al. Effect of high voltage pulsed current plus conventional treatment on acute ankle sprain. *Rev Bras Fisioter*. 2010; 14(3):193-199.~~

~~Seco J, Kovacs F, Urrutia G. The efficacy, safety, effectiveness, and cost-effectiveness of ultrasound and shock wave therapies for low back pain: A systematic review. *Spine J*. 2011; 11(10):966-977.~~

~~Snyder AR, Perotti AL, Lam KC, et al. Influence of high-voltage electrical stimulation on edema formation after acute injury: A systematic review. *J Sport Rehabil*. 2010; 19(4):436-451.~~

~~Southerst D, Yu H, Randhawa K, et al. The effectiveness of manual therapy for the management of musculoskeletal disorders of the upper and lower extremities: a systematic review by the Ontario Protocol for Traffic Injury Management (OPTIMA) collaboration. *Chiropr Man Therap*. 2015; 23:30.~~

~~Thackeray A, Fritz JM, Childs JD, et al. The Effectiveness of mechanical traction among subgroups of patients with low back pain and leg pain: A randomized trial. *J Orthop Sports Phys Ther*. 2016; 46(3):144-154.~~

~~Thoomes EJ. Effectiveness of manual therapy for cervical radiculopathy: A review. *Chiropr Man Therap*. 2016; 24:45.~~

~~Tsakitzidis G, Remmen R, Dankaerts W, et al. Non-specific neck pain and evidence-based practice. *European Scientific Journal*. 2013; 9(3).~~

~~University of Michigan Health System. Acute low back pain guidelines for clinical care. *University of Michigan Health System*. Ann Arbor, MI. 2003. Revised 2010. Available at www.cme.med.umich.edu.~~

~~Van den Bekerom MP, van der Windt DA, Ter Riet G, et al. Therapeutic ultrasound for acute ankle sprains. *Eur J Phys Rehabil Med*. June 2012; 48(2):325-334.~~

~~Van den Dolder PA, Ferreira PH, Refshauge KM. Effectiveness of soft tissue massage and exercise for the treatment of non-specific shoulder pain: A systematic review with meta-analysis. *Br J Sports Med*. 2014; 48(16):1216.~~

~~Van Middelkoop M, Rubinstein SM, Kuijpers T, et al. A systematic review on the effectiveness of physical and rehabilitation interventions for chronic non-specific low back pain. *Eur Spine J*. 2011; 20(1):19-39.~~

~~Walsh DM, Howe TE, Johnson MI, et al. Transcutaneous electrical nerve stimulation for acute pain. *Cochrane Database of Syst Rev*. 2009; 2:CD006142.~~

~~Wegner I, Widyahening IS, et al. Traction for low back pain with or without sciatica. *Cochrane Database Syst Rev*. August 19, 2013; (8):CD003010. doi: 10.1002/14651858.CD003010.pub5.~~

~~Yang JD, Tam KW, Huang TW, Huang SW, Liou TH, Chen HC. Intermittent Cervical Traction for Treating Neck Pain. *Spine*. 2017; 42(13):959-965~~

~~Yesil H, Hepguler S, Dundar U, et al. Does the Use of Electrotherapies Increase the Effectiveness of Neck Stabilization Exercises for Improving Pain, Disability, Mood, and Quality of life in Chronic Neck Pain? A Randomized, Controlled, Single Blind Study. *Spine*. 2018; 43(20):E1174-E1183.~~

~~Ylinen J, Kautiainen H, Wiren K, et al. Stretching exercises vs manual therapy in treatment of chronic neck pain: A randomized, controlled cross-over trial. *J Rehabil Med*. 2007; 39(2):126-132.~~

~~Xia P, Wang X, Lin Q, et al. Effectiveness of ultrasound therapy for myofascial pain syndrome: a systematic review and meta-analysis. *J Pain Res*. 2017; 10:545-555.~~

~~Zeng C, Li H, Yang T, et al. Electrical stimulation for pain relief in knee osteoarthritis: systematic review and network meta-analysis. *Osteoarthritis Cartilage*. 2015; 23(2):189-202.~~

~~Song HJ, Seo HJ, Lee Y, et al. Effectiveness of high-intensity laser therapy in the treatment of musculoskeletal disorders: A systematic review and meta-analysis of randomized controlled trials. *Medicine (Baltimore)*. 2019; 98(4):e14274.~~

Reviewed/Approved by NIA Clinical Guideline Committee

Reviewed / Approved by
Physical Medicine



Kathleen Norman, D.C., Medical Director,

Disclaimer: Magellan Healthcare service authorization policies do not constitute medical advice and are not intended to govern or otherwise influence the practice of medicine. These policies are not meant to supplant your normal procedures, evaluation, diagnosis, treatment and/or care plans for your patients. Your professional judgement must be exercised and followed in all respects with regard to the treatment and care of your patients. These policies apply to all Magellan Healthcare subsidiaries including, but not limited to, National Imaging Associates (“Magellan”). The policies constitute only the reimbursement and coverage guidelines of Magellan. Coverage for services varies for individual members in accordance with the terms and conditions of applicable Certificates of Coverage, Summary Plan Descriptions, or contracts with governing regulatory agencies. Magellan reserves the right to review and update the guidelines at its sole discretion. Notice of such changes, if necessary, shall be provided in accordance with the terms and conditions of provider agreements and any applicable laws or regulations.

1. Qaseem A, Wilt TJ, McLean RM, et al. Noninvasive Treatments for Acute, Subacute, and Chronic Low Back Pain: A Clinical Practice Guideline From the American College of Physicians. *Ann Intern Med*. Apr 4 2017;166(7):514-530. doi:10.7326/m16-2367
2. Ghasabmahaleh SH, Rezasoltani Z, Dadarkhah A, Hamidipanah S, Mofrad RK, Najafi S. Spinal Manipulation for Subacute and Chronic Lumbar Radiculopathy: A Randomized Controlled Trial. *Am J Med*. Jan 2021;134(1):135-141. doi:10.1016/j.amjmed.2020.08.005
3. Paige NM, Miake-Lye IM, Booth MS, et al. Association of Spinal Manipulative Therapy With Clinical Benefit and Harm for Acute Low Back Pain: Systematic Review and Meta-analysis. *JAMA*. 2017;317(14):1451-1460. doi:10.1001/jama.2017.3086
4. Karimi MT, Rabczuk T. Scoliosis conservative treatment: A review of literature. *J Craniovertebr Junction Spine*. Jan-Mar 2018;9(1):3-8. doi:10.4103/jcvjs.JCVJS_39_17
5. Chiodo A, Bhat S, Harrison R, Shumer G, Wasserman R. Ambulatory Adult Low Back Pain Guideline. Michigan Medicine University of Michigan. Updated December 2020. Accessed October 18, 2021. <https://michmed-public.policystat.com/policy/8093107/latest/>
6. Khadilkar A, Odebiyi DO, Brosseau L, Wells GA. Transcutaneous electrical nerve stimulation (TENS) versus placebo for chronic low-back pain. *Cochrane Database Syst Rev*. 2008;2008(4):CD003008-CD003008. doi:10.1002/14651858.CD003008.pub3
7. Greene MW, Clemente-Fuentes RJW. Is transcutaneous electrical nerve stimulation (TENS) more effective than placebo for management of chronic low back pain? *Evidence-Based Practice*. 2016;19(8):10-11.
8. Rajfur J, Pasternok M, Rajfur K, et al. Efficacy of Selected Electrical Therapies on Chronic Low Back Pain: A Comparative Clinical Pilot Study. *Med Sci Monit*. 2017;23:85-100. doi:10.12659/msm.899461
9. Savigny P, Watson P, Underwood M. Early management of persistent non-specific low back pain: summary of NICE guidance. *Bmj*. Jun 4 2009;338:b1805. doi:10.1136/bmj.b1805

10. Savigny P, Kuntze S, Watson P, et al. Low back pain: early management of persistent non-specific low back pain. *London: National Collaborating Centre for Primary Care and Royal College of General Practitioners*. 2009;14(1):9-13.
11. Wegner I, Widyahening IS, van Tulder MW, et al. Traction for low-back pain with or without sciatica. *Cochrane Database Syst Rev*. 2013;2013(8):CD003010-CD003010. doi:10.1002/14651858.CD003010.pub5
12. Alrwaily M, Almutiri M, Schneider M. Assessment of variability in traction interventions for patients with low back pain: a systematic review. *Chiropr Man Therap*. 2018;26:35-35. doi:10.1186/s12998-018-0205-z
13. Chou R, Deyo R, Friedly J, et al. AHRQ Comparative Effectiveness Reviews. *Noninvasive Treatments for Low Back Pain*. Agency for Healthcare Research and Quality (US); 2016.
14. Analan PD, Leblebici B, Adam M. Effects of therapeutic ultrasound and exercise on pain, function, and isokinetic shoulder rotator strength of patients with rotator cuff disease. *J Phys Ther Sci*. 2015;27(10):3113-3117. doi:10.1589/jpts.27.3113
15. Ebadi S, Henschke N, Forogh B, et al. Therapeutic ultrasound for chronic low back pain. *Cochrane Database Syst Rev*. 2020;7(7):CD009169-CD009169. doi:10.1002/14651858.CD009169.pub3
16. Furlan AD, Giraldo M, Baskwill A, Irvin E, Imamura M. Massage for low-back pain. *Cochrane Database Syst Rev*. Sep 1 2015;(9):Cd001929. doi:10.1002/14651858.CD001929.pub3
17. Chou R, Deyo R, Friedly J, et al. Nonpharmacologic Therapies for Low Back Pain: A Systematic Review for an American College of Physicians Clinical Practice Guideline. *Ann Intern Med*. Apr 4 2017;166(7):493-505. doi:10.7326/m16-2459
18. van Middelkoop M, Rubinstein SM, Kuijpers T, et al. A systematic review on the effectiveness of physical and rehabilitation interventions for chronic non-specific low back pain. *European spine journal : official publication of the European Spine Society, the European Spinal Deformity Society, and the European Section of the Cervical Spine Research Society*. 2011;20(1):19-39. doi:10.1007/s00586-010-1518-3
19. Yang JD, Tam KW, Huang TW, Huang SW, Liou TH, Chen HC. Intermittent Cervical Traction for Treating Neck Pain: A Meta-analysis of Randomized Controlled Trials. *Spine (Phila Pa 1976)*. Jul 1 2017;42(13):959-965. doi:10.1097/brs.0000000000001948
20. Romeo A, Vanti C, Boldrini V, et al. Cervical radiculopathy: effectiveness of adding traction to physical therapy-a systematic review and meta-analysis of randomized controlled trials. *Phys Ther*. 2018;98:231-242. *Phys Ther*. Aug 1 2018;98(8):727. doi:10.1093/ptj/pzy064
21. Romeo A, Vanti C, Boldrini V, et al. Cervical Radiculopathy: Effectiveness of Adding Traction to Physical Therapy-A Systematic Review and Meta-Analysis of Randomized Controlled Trials. *Phys Ther*. Apr 1 2018;98(4):231-242. doi:10.1093/phyth/pzy001
22. Bernal-Utrera C, Gonzalez-Gerez JJ, Anarte-Lazo E, Rodriguez-Blanco C. Manual therapy versus therapeutic exercise in non-specific chronic neck pain: a randomized controlled trial. *Trials*. Jul 28 2020;21(1):682. doi:10.1186/s13063-020-04610-w
23. Miller J, Gross A, D'Sylva J, et al. Manual therapy and exercise for neck pain: a systematic review. *Man Ther*. Aug 2010;15(4):334-54.

24. Coulter ID, Crawford C, Vernon H, et al. Manipulation and Mobilization for Treating Chronic Nonspecific Neck Pain: A Systematic Review and Meta-Analysis for an Appropriateness Panel. *Pain Physician*. 2019;22(2):E55-E70.
25. Díaz-Pulido B, Pérez-Martín Y, Pecos-Martín D, et al. Efficacy of Manual Therapy and Transcutaneous Electrical Nerve Stimulation in Cervical Mobility and Endurance in Subacute and Chronic Neck Pain: A Randomized Clinical Trial. *J Clin Med*. 2021;10(15):3245. doi:10.3390/jcm10153245
26. Ylinen J, Kautiainen H, Wirén K, Häkkinen A. Stretching exercises vs manual therapy in treatment of chronic neck pain: a randomized, controlled cross-over trial. *J Rehabil Med*. Mar 2007;39(2):126-32. doi:10.2340/16501977-0015
27. Hidalgo B, Hall T, Bossert J, Dugeny A, Cagnie B, Pitance L. The efficacy of manual therapy and exercise for treating non-specific neck pain: A systematic review. *J Back Musculoskelet Rehabil*. 2017;30(6):1149-1169. doi:10.3233/BMR-169615
28. Falsiroli Maistrello L, Geri T, Gianola S, Zaninetti M, Testa M. Effectiveness of Trigger Point Manual Treatment on the Frequency, Intensity, and Duration of Attacks in Primary Headaches: A Systematic Review and Meta-Analysis of Randomized Controlled Trials. *Front Neurol*. 2018;9:254-254. doi:10.3389/fneur.2018.00254
29. Haas M, Bronfort G, Evans R, et al. Dose-response and efficacy of spinal manipulation for care of cervicogenic headache: a dual-center randomized controlled trial. *Spine J*. 2018;18(10):1741-1754. doi:10.1016/j.spinee.2018.02.019
30. Tsakitzidis G, Remmen R, Dankaerts W, Van Royen P. Non-specific neck pain and evidence-based practice. *European scientific journal*. 2013;9(3):1-19. doi:<https://doi.org/10.19044/esj.2013.v9n3p%25p>
31. Daher A, Carel RS, Tzipi K, Esther H, Dar G. The effectiveness of an aerobic exercise training on patients with neck pain during a short- and long-term follow-up: a prospective double-blind randomized controlled trial. *Clin Rehabil*. May 2020;34(5):617-629. doi:10.1177/0269215520912000
32. Yesil H, Hepguler S, Dundar U, Taravati S, Isleten B. Does the Use of Electrotherapies Increase the Effectiveness of Neck Stabilization Exercises for Improving Pain, Disability, Mood, and Quality of Life in Chronic Neck Pain?: A Randomized, Controlled, Single-Blind Study. *Spine (Phila Pa 1976)*. Oct 15 2018;43(20):E1174-e1183. doi:10.1097/brs.0000000000002663
33. Albornoz-Cabello M, Pérez-Mármol JM, Barrios Quinta CJ, Matarán-Peñarrocha GA, Castro-Sánchez AM, de la Cruz Olivares B. Effect of adding interferential current stimulation to exercise on outcomes in primary care patients with chronic neck pain: a randomized controlled trial. *Clin Rehabil*. Sep 2019;33(9):1458-1467. doi:10.1177/0269215519844554
34. Celenay ST, Kaya DO, Akbayrak T. Cervical and scapulothoracic stabilization exercises with and without connective tissue massage for chronic mechanical neck pain: A prospective, randomised controlled trial. *Man Ther*. Feb 2016;21:144-50. doi:10.1016/j.math.2015.07.003
35. Chou R, Côté P, Randhawa K, et al. The Global Spine Care Initiative: applying evidence-based guidelines on the non-invasive management of back and neck pain to low- and middle-income communities. *Eur Spine J*. Sep 2018;27(Suppl 6):851-860. doi:10.1007/s00586-017-5433-8

36. Southerst D, Yu H, Randhawa K, et al. The effectiveness of manual therapy for the management of musculoskeletal disorders of the upper and lower extremities: a systematic review by the Ontario Protocol for Traffic Injury Management (OPTIMA) Collaboration. *Chiropr Man Therap*. 2015;23:30-30. doi:10.1186/s12998-015-0075-6
37. Desjardins-Charbonneau A, Roy JS, Dionne CE, Frémont P, MacDermid JC, Desmeules F. The efficacy of manual therapy for rotator cuff tendinopathy: a systematic review and meta-analysis. *J Orthop Sports Phys Ther*. May 2015;45(5):330-50. doi:10.2519/jospt.2015.5455
38. Steuri R, Sattelmayer M, Elsig S, et al. Effectiveness of conservative interventions including exercise, manual therapy and medical management in adults with shoulder impingement: a systematic review and meta-analysis of RCTs. *Br J Sports Med*. 2017;51(18):1340-1347. doi:10.1136/bjsports-2016-096515
39. Pettersson S, Plancher K, Klyve D, Draper D, Ortiz R. Low-Intensity Continuous Ultrasound for the Symptomatic Treatment of Upper Shoulder and Neck Pain: A Randomized, Double-Blind Placebo-Controlled Clinical Trial. *J Pain Res*. 2020;13:1277-1287. doi:10.2147/jpr.S247463
40. Erratum: Low-Intensity Continuous Ultrasound for the Symptomatic Treatment of Upper Shoulder and Neck Pain: A Randomized, Double-Blind Placebo-Controlled Clinical Trial [Corrigendum]. *J Pain Res*. 2020;13:1899-1900. doi:10.2147/jpr.S272270
41. Page MJ, Green S, Mrocki MA, et al. Electrotherapy modalities for rotator cuff disease. *Cochrane Database Syst Rev*. Jun 10 2016;(6):Cd012225. doi:10.1002/14651858.Cd012225
42. Joseph MF, Taft K, Moskwa M, Denegar CR. Deep friction massage to treat tendinopathy: a systematic review of a classic treatment in the face of a new paradigm of understanding. *J Sport Rehabil*. Nov 2012;21(4):343-53. doi:10.1123/jsr.21.4.343
43. Beumer L, Wong J, Warden SJ, Kemp JL, Foster P, Crossley KM. Effects of exercise and manual therapy on pain associated with hip osteoarthritis: a systematic review and meta-analysis. *Br J Sports Med*. 2016;50(8):458-463.
44. Sampath KK, Mani R, Miyamori T, Tumilty S. The effects of manual therapy or exercise therapy or both in people with hip osteoarthritis: a systematic review and meta-analysis. *Clin Rehabil*. Dec 2016;30(12):1141-1155. doi:10.1177/0269215515622670
45. Kemp JL, Mosler AB, Hart H, et al. Improving function in people with hip-related pain: a systematic review and meta-analysis of physiotherapist-led interventions for hip-related pain. *British journal of sports medicine*. 2020;54(23):1382-1394. doi:10.1136/bjsports-2019-101690
46. Chaves P, Simões D, Paço M, et al. Deep Friction Massage in the Management of Patellar Tendinopathy in Athletes: Short-Term Clinical Outcomes. *J Sport Rehabil*. Sep 1 2020;29(7):860-865. doi:10.1123/jsr.2019-0046
47. Xia P, Wang X, Lin Q, Cheng K, Li X. Effectiveness of ultrasound therapy for myofascial pain syndrome: a systematic review and meta-analysis. *J Pain Res*. 2017;10:545-555. doi:10.2147/JPR.S131482
48. Wu Y, Zhu S, Lv Z, et al. Effects of therapeutic ultrasound for knee osteoarthritis: a systematic review and meta-analysis. *Clin Rehabil*. Dec 2019;33(12):1863-1875. doi:10.1177/0269215519866494
49. Rutjes AW, Nüesch E, Sterchi R, et al. Transcutaneous electrostimulation for osteoarthritis of the knee. *Cochrane Database Syst Rev*. 2009;2009(4):CD002823-CD002823. doi:10.1002/14651858.CD002823.pub2

50. Vance CGT, Rakel BA, Blodgett NP, et al. Effects of transcutaneous electrical nerve stimulation on pain, pain sensitivity, and function in people with knee osteoarthritis: a randomized controlled trial. *Phys Ther*. 2012;92(7):898-910. doi:10.2522/ptj.20110183
51. Maeda T, Yoshida H, Sasaki T, Oda A. Does transcutaneous electrical nerve stimulation (TENS) simultaneously combined with local heat and cold applications enhance pain relief compared with TENS alone in patients with knee osteoarthritis? *J Phys Ther Sci*. 2017;29(10):1860-1864. doi:10.1589/jpts.29.1860
52. Tumilty S, Munn J, Abbott JH, McDonough S, Hurley DA, Baxter GD. Laser therapy in the treatment of achilles tendinopathy: a pilot study. *Photomed Laser Surg*. Feb 2008;26(1):25-30. doi:10.1089/pho.2007.2126
53. Neeter C, Thomeé R, Silbernagel KG, Thomeé P, Karlsson J. Iontophoresis with or without dexamethazone in the treatment of acute Achilles tendon pain. *Scand J Med Sci Sports*. Dec 2003;13(6):376-82. doi:10.1046/j.1600-0838.2003.00305.x
54. Zhou J, Salvendy G. *Human Aspects of IT for the Aged Population. Design for Aging: First International Conference, ITAP 2015, Held as Part of HCI International 2015, Los Angeles, CA, USA, August 2-7, 2015. Proceedings, Part I*. vol 9193. Springer; 2015.
55. Fuentes JP, Armijo Olivo S, Magee DJ, Gross DP. Effectiveness of interferential current therapy in the management of musculoskeletal pain: a systematic review and meta-analysis. *Phys Ther*. Sep 2010;90(9):1219-38. doi:10.2522/ptj.20090335
56. Song HJ, Seo H-J, Lee Y, Kim SK. Effectiveness of high-intensity laser therapy in the treatment of musculoskeletal disorders: A systematic review and meta-analysis of randomized controlled trials. *Medicine (Baltimore)*. 2018;97(51):e13126-e13126. doi:10.1097/MD.00000000000013126

ADDITIONAL RESOURCES

1. [Armijo-Olivo S, Pitance L, Singh V, Neto F, Thie N, Michelotti A. Effectiveness of Manual Therapy and Therapeutic Exercise for Temporomandibular Disorders: Systematic Review and Meta-Analysis. *Physical therapy*. 2016;96\(1\):9-25. doi:10.2522/ptj.20140548](#)
2. [Babatunde OO, Jordan JL, Van der Windt DA, Hill JC, Foster NE, Protheroe J. Effective treatment options for musculoskeletal pain in primary care: A systematic overview of current evidence. *PLoS One*. 2017;12\(6\):e0178621-e0178621. doi:10.1371/journal.pone.0178621](#)
3. [Baez S, Hoch JM, Uhl TL. The effectiveness of cervical traction and exercise in decreasing neck and arm pain for patients with cervical radiculopathy: a critically appraised topic. *Int J Athl Ther Train*. 2017;22\(5\):4.](#)
4. [Bagheripour B, Kamyab M, Azadinia F, Amiri A, Akbari M. The efficacy of a home-mechanical traction unit for patients with mild to moderate cervical osteoarthritis: A pilot study. *Med J Islam Repub Iran*. 2016;30:386-386.](#)
5. [Bryans R, Decina P, Descarreaux M, et al. Evidence-based guidelines for the chiropractic treatment of adults with neck pain. *J Manipulative Physiol Ther*. Jan 2014;37\(1\):42-63. doi:10.1016/j.jmpt.2013.08.010](#)
6. [Buchbinder R, van Tulder M, Öberg B, et al. Low back pain: a call for action. *Lancet*. Jun 9 2018;391\(10137\):2384-2388. doi:10.1016/s0140-6736\(18\)30488-4](#)

7. Bukhari SRI, Shakil-Ur-Rehman S, Ahmad S, Naeem A. Comparison between effectiveness of Mechanical and Manual Traction combined with mobilization and exercise therapy in Patients with Cervical Radiculopathy. *Pak J Med Sci.* Jan-Feb 2016;32(1):31-34. doi:10.12669/pjms.321.8923
8. Cai C, Pua YH, Lim KC. A clinical prediction rule for classifying patients with low back pain who demonstrate short-term improvement with mechanical lumbar traction. *Eur Spine J.* 2009;18(4):554-561. doi:10.1007/s00586-009-0909-9
9. Cakir S, Hepguler S, Ozturk C, Korkmaz M, Isleten B, Atamaz FC. Efficacy of therapeutic ultrasound for the management of knee osteoarthritis: a randomized, controlled, and double-blind study. *Am J Phys Med Rehabil.* May 2014;93(5):405-12. doi:10.1097/phm.0000000000000033
10. Carey TS, Freburger JK, Holmes GM, et al. A long way to go: practice patterns and evidence in chronic low back pain care. *Spine.* 2009;34(7):718-724. doi:10.1097/BRS.0b013e31819792b0
11. Centers for Medicare & Medicaid Services. Transcutaneous Electrical Nerve Stimulation for Chronic Low Back Pain: CAG-00429N. Centers for Medicare & Medicaid Services (CMS). Updated June 8, 2012. Accessed September 23, 2012. <https://www.cms.gov/medicare-coverage-database/view/ncacal-decision-memo.aspx?proposed=N&NCAId=256>
12. Cherian JJ, Jauregui JJ, Leichter AK, Elmallah RK, Bhawe A, Mont MA. The effects of various physical non-operative modalities on the pain in osteoarthritis of the knee. *Bone Joint J.* Jan 2016;98-b(1 Suppl A):89-94. doi:10.1302/0301-620x.98b1.36353
13. Chou R. Patient education: Low back pain in adults (Beyond the Basics). Wolters Kluwer. Updated September 20, 2021. Accessed September 23, 2021. <https://www.uptodate.com/contents/low-back-pain-in-adults-beyond-the-basics>
14. Clar C, Tsertsvadze A, Court R, Hundt GL, Clarke A, Sutcliffe P. Clinical effectiveness of manual therapy for the management of musculoskeletal and non-musculoskeletal conditions: systematic review and update of UK evidence report. *Chiropr Man Therap.* 2014;22(1):12-12. doi:10.1186/2045-709X-22-12
15. Coulter ID, Crawford C, Hurwitz EL, et al. Manipulation and mobilization for treating chronic low back pain: a systematic review and meta-analysis. *Spine J.* 2018;18(5):866-879. doi:10.1016/j.spinee.2018.01.013
16. Crawford C, Boyd C, Paat CF, et al. The Impact of Massage Therapy on Function in Pain Populations-A Systematic Review and Meta-Analysis of Randomized Controlled Trials: Part I, Patients Experiencing Pain in the General Population. *Pain Med.* 2016;17(7):1353-1375. doi:10.1093/pm/pnw099
17. Damgaard P, Bartels EM, Ris I, Christensen R, Juul-Kristensen B. Evidence of Physiotherapy Interventions for Patients with Chronic Neck Pain: A Systematic Review of Randomised Controlled Trials. *ISRN Pain.* 2013;2013:567175-567175. doi:10.1155/2013/567175
18. Desmeules F, Boudreault J, Roy JS, Dionne C, Frémont P, MacDermid JC. The efficacy of therapeutic ultrasound for rotator cuff tendinopathy: A systematic review and meta-analysis. *Phys Ther Sport.* Aug 2015;16(3):276-84. doi:10.1016/j.ptsp.2014.09.004

19. Ebadi S, Henschke N, Nakhostin Ansari N, Fallah E, van Tulder MW. Therapeutic ultrasound for chronic low-back pain. *Cochrane Database Syst Rev*. Mar 14 2014;(3):Cd009169. doi:10.1002/14651858.CD009169.pub2
20. Farooq MN, Mohseni-Bandpei MA, Gilani SA, Ashfaq M, Mahmood Q. The effects of neck mobilization in patients with chronic neck pain: A randomized controlled trial. *J Bodyw Mov Ther*. Jan 2018;22(1):24-31. doi:10.1016/j.jbmt.2017.03.007
21. Foster NE, Anema JR, Cherkin D, et al. Prevention and treatment of low back pain: evidence, challenges, and promising directions. *Lancet*. Jun 9 2018;391(10137):2368-2383. doi:10.1016/s0140-6736(18)30489-6
22. Fritz JM, Thackeray A, Brennan GP, Childs JD. Exercise only, exercise with mechanical traction, or exercise with over-door traction for patients with cervical radiculopathy, with or without consideration of status on a previously described subgrouping rule: a randomized clinical trial. *J Orthop Sports Phys Ther*. Feb 2014;44(2):45-57. doi:10.2519/jospt.2014.5065
23. Furlan AD, Yazdi F, Tsertsvadze A, et al. A systematic review and meta-analysis of efficacy, cost-effectiveness, and safety of selected complementary and alternative medicine for neck and low-back pain. *Evid Based Complement Alternat Med*. 2012;2012:953139-953139. doi:10.1155/2012/953139
24. Glazov G, Yelland M, Emery J. Low-level laser therapy for chronic non-specific low back pain: a meta-analysis of randomised controlled trials. *Acupunct Med*. 2016;34(5):328-341. doi:10.1136/acupmed-2015-011036
25. Hawk C, Minkalis AL, Khorsan R, et al. Systematic Review of Nondrug, Nonsurgical Treatment of Shoulder Conditions. *J Manipulative Physiol Ther*. Jun 2017;40(5):293-319. doi:10.1016/j.jmpt.2017.04.001
26. Isaac Z, Dec K. Patient education: Neck pain (Beyond the Basics). Wolters Kluwer. Updated September 3, 2020. Accessed September 23, 2021. <https://www.uptodate.com/contents/neck-pain-beyond-the-basics>
27. Johnson MI, Paley CA, Howe TE, Sluka KA. Transcutaneous electrical nerve stimulation for acute pain. *Cochrane Database Syst Rev*. 2015;2015(6):CD006142-CD006142. doi:10.1002/14651858.CD006142.pub3
28. Kroeling P, Gross A, Graham N, et al. Electrotherapy for neck pain. *Cochrane Database Syst Rev*. Aug 26 2013;(8):Cd004251. doi:10.1002/14651858.CD004251.pub5
29. Walsh DM, Howe TE, Johnson MI, Sluka KA. Transcutaneous electrical nerve stimulation for acute pain. *Cochrane Database Syst Rev*. Apr 15 2009;(2):Cd006142. doi:10.1002/14651858.CD006142.pub2
30. Kumar S, Beaton K, Hughes T. The effectiveness of massage therapy for the treatment of nonspecific low back pain: a systematic review of systematic reviews. *Int J Gen Med*. 2013;6:733-741. doi:10.2147/IJGM.S50243
31. Littlewood C, May S, Walters S. A review of systematic reviews of the effectiveness of conservative interventions for rotator cuff tendinopathy. *Shoulder & Elbow*. 2013;5(3):151-167.
32. Martimbianco ALC, Torloni MR, Andriolo BN, Porfirio GJ, Riera R. Neuromuscular electrical stimulation (NMES) for patellofemoral pain syndrome. *Cochrane Database Syst Rev*. 2017;12(12):CD011289-CD011289. doi:10.1002/14651858.CD011289.pub2

33. Negrini S, Minozzi S, Bettany-Saltikov J, et al. Braces for idiopathic scoliosis in adolescents. *Cochrane Database Syst Rev*. Jan 20 2010;(1):Cd006850. doi:10.1002/14651858.CD006850.pub2
34. Nelson NL, Churilla JR. Massage Therapy for Pain and Function in Patients With Arthritis: A Systematic Review of Randomized Controlled Trials. *Am J Phys Med Rehabil*. Sep 2017;96(9):665-672. doi:10.1097/phm.0000000000000712
35. Page MJ, Green S, Kramer S, Johnston RV, McBain B, Buchbinder R. Electrotherapy modalities for adhesive capsulitis (frozen shoulder). *Cochrane Database Syst Rev*. Oct 1 2014;(10):Cd011324. doi:10.1002/14651858.Cd011324
36. Page MJ, Green S, Mrocki MA, et al. Electrotherapy modalities for rotator cuff disease. *Cochrane Database Syst Rev*. Jun 10 2016;(6):Cd012225. doi:10.1002/14651858.Cd012225
37. Palmer E, Redavid L. Therapeutic Ultrasound: Clinical Review. *Clin Info Sys (EBSCO)*. January 8, 2016 2016:1-14. doi: [https://www.ebscohost.com/assets-sample-content/Therapeutic Ultrasound - CR.pdf](https://www.ebscohost.com/assets-sample-content/Therapeutic%20Ultrasound%20-%20CR.pdf)
38. Rankin IA, Sargeant H, Rehman H, Gurusamy KS. Low-level laser therapy for carpal tunnel syndrome. *Cochrane Database of Systematic Reviews*. 2017;(8)doi:10.1002/14651858.CD012765
39. Salazar AP, Stein C, Marchese RR, Plentz RD, Pagnussat AS. Electric Stimulation for Pain Relief in Patients with Fibromyalgia: A Systematic Review and Meta-analysis of Randomized Controlled Trials. *Pain Physician*. Feb 2017;20(2):15-25.
40. Sandoval MC, Ramirez C, Camargo DM, Salvini TF. Effect of high-voltage pulsed current plus conventional treatment on acute ankle sprain. *Rev Bras Fisioter*. May-Jun 2010;14(3):193-9. doi:10.1590/s1413-35552010000300012
41. Seco J, Kovacs FM, Urrutia G. The efficacy, safety, effectiveness, and cost-effectiveness of ultrasound and shock wave therapies for low back pain: a systematic review. *Spine J*. Oct 2011;11(10):966-77. doi:10.1016/j.spinee.2011.02.002
42. Snyder AR, Perotti AL, Lam KC, Bay RC. The influence of high-voltage electrical stimulation on edema formation after acute injury: a systematic review. *J Sport Rehabil*. Nov 2010;19(4):436-51. doi:10.1123/jsr.19.4.436
43. Thackeray A, Fritz JM, Childs JD, Brennan GP. The Effectiveness of Mechanical Traction Among Subgroups of Patients With Low Back Pain and Leg Pain: A Randomized Trial. *J Orthop Sports Phys Ther*. Mar 2016;46(3):144-54. doi:10.2519/jospt.2016.6238
44. Thoomes EJ. Effectiveness of manual therapy for cervical radiculopathy, a review. *Chiropr Man Therap*. 2016;24:45-45. doi:10.1186/s12998-016-0126-7
45. Chiodo A, Alvarez D, Graziano G, et al. Acute Low Back Pain. Regents of the University of Michigan Updated December 2011. Accessed September 24, 2021. <http://www.med.umich.edu/1info/FHP/practiceguides/back/back.pdf>
46. van den Bekerom MP, van der Windt DA, Ter Riet G, van der Heijden GJ, Bouter LM. Therapeutic ultrasound for acute ankle sprains. *Eur J Phys Rehabil Med*. Jun 2012;48(2):325-34.
47. van den Dolder PA, Ferreira PH, Refshauge KM. Effectiveness of soft tissue massage and exercise for the treatment of non-specific shoulder pain: a systematic review with meta-analysis. *Br J Sports Med*. Aug 2014;48(16):1216-26. doi:10.1136/bjsports-2011-090553

48. [Zeng C, Li H, Yang T, et al. Electrical stimulation for pain relief in knee osteoarthritis: systematic review and network meta-analysis. *Osteoarthritis Cartilage*. Feb 2015;23\(2\):189-202. doi:10.1016/j.joca.2014.11.014](#)

[Reviewed/Approved by NIA Clinical Guideline Committee](#)

GENERAL INFORMATION

It is an expectation that all patients receive care/services from a licensed clinician. All appropriate supporting documentation, including recent pertinent office visit notes, laboratory data, and results of any special testing must be provided. If applicable: All prior relevant imaging results and the reason that alternative imaging cannot be performed must be included in the documentation submitted.

Disclaimer: Magellan Healthcare service authorization policies do not constitute medical advice and are not intended to govern or otherwise influence the practice of medicine. These policies are not meant to supplant your normal procedures, evaluation, diagnosis, treatment and/or care plans for your patients. Your professional judgement must be exercised and followed in all respects with regard to the treatment and care of your patients. These policies apply to all Magellan Healthcare subsidiaries including, but not limited to, National Imaging Associates (“Magellan”). The policies constitute only the reimbursement and coverage guidelines of Magellan. Coverage for services varies for individual members in accordance with the terms and conditions of applicable Certificates of Coverage, Summary Plan Descriptions, or contracts with governing regulatory agencies. Magellan reserves the right to review and update the guidelines at its sole discretion. Notice of such changes, if necessary, shall be provided in accordance with the terms and conditions of provider agreements and any applicable laws or regulations.