



Clinical Policy: Pasireotide (Signifor, Signifor LAR)

Reference Number: LA.PHAR.332

Effective Date: 11.04.23

Last Review Date: 11.19.25~~12.03.24~~

Line of Business: Medicaid

[Coding Implications](#)
[Revision Log](#)

See [Important Reminder](#) at the end of this policy for important regulatory and legal information.

****Please note: This policy is for medical benefit****

Description

Pasireotide (Signifor®, Signifor® LAR) is a somatostatin analog.

FDA Approved Indication(s)

Signifor and Signifor LAR are indicated for the treatment of patients with Cushing's disease for whom pituitary surgery is not an option or has not been curative. Signifor is specifically indicated in adults.

Signifor LAR is also indicated for the treatment of patients with acromegaly who have had an inadequate response to surgery and/or for whom surgery is not an option.

Policy/Criteria

Provider must submit documentation (such as office chart notes, lab results or other clinical information) supporting that member has met all approval criteria.

It is the policy of Louisiana Healthcare Connections that Signifor and Signifor LAR are **medically necessary** when the following criteria are met:

I. Initial Approval Criteria

A. Acromegaly (must meet all):

1. Diagnosis of acromegaly as evidenced by one of the following (a or b):
 - a. Pre-treatment insulin-like growth factor-I (IGF-I) level above the upper limit of normal based on age and gender for the reporting laboratory;
 - b. Serum growth hormone (GH) level $\geq 1 \mu\text{g/L}$ after a 2-hour oral glucose tolerance test;
2. Request is for Signifor LAR;
3. Prescribed by or in consultation with an endocrinologist;
4. Age ≥ 18 years;
5. Inadequate response to surgical resection or pituitary irradiation (*see Appendix D*), or member is not a candidate for such treatment;
6. Dose does not exceed (a and b):
 - a. 60 mg every 4 weeks;
 - b. 1 vial every 4 weeks.

Approval duration: ~~6~~12 months

Formatted: Font: Bold

CLINICAL POLICY

Pasireotide



B. Cushing's Disease (must meet all):

1. Diagnosis of Cushing's disease;
2. Prescribed by or in consultation with an endocrinologist;
3. Age \geq 18 years;
4. Member meets one of the following (a or b):
 - a. Pituitary surgery was not curative;
 - b. Member is not eligible for pituitary surgery;
5. Dose does not exceed one of the following (a or b):
 - a. Signifor (i and ii):
 - i. 1.8 mg per day;
 - ii. 2 ampules per day;
 - b. Signifor LAR (i and ii):
 - i. 40 mg every 4 weeks;
 - ii. 1 vial every 4 weeks.

Approval duration: **12 months**

Formatted: Font: Bold

6 months

Formatted: Font: Not Bold

C. Other diagnoses/indications (must meet 1 or 2):

- 4a.** If this drug has recently (within the last 6 months) undergone a label change (e.g., newly approved indication, age expansion, new dosing regimen) that is not yet reflected in this policy, refer to LA.PMN.255
2. If the requested use (e.g., diagnosis, age, dosing regimen) is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized) AND criterion 1 above does not apply, refer to the off-label use policy **for the relevant line of business**—LA.PMN.53—**for Medicaid**.

Formatted: Indent: Left: 0.75"

II. Continued Therapy

A. Acromegaly (must meet all):

Currently receiving medication via Louisiana Healthcare Connections benefit or member has previously met initial approval criteria;

1. Request is for Signifor LAR;
2. Member is responding positively to therapy (*see Appendix D*);
3. If request is for a dose increase, new dose does not exceed (a and b):
 - a. 60 mg every 4 weeks;
 - b. 1 vial every 4 weeks.

Approval duration: **12 months**

Formatted: Font: Bold

B. Cushing's Disease (must meet all):

- a. Currently receiving medication via Louisiana Healthcare Connections benefit or member has previously met initial approval criteria;
2. Member is responding positively to therapy (*see Appendix D*);
3. If request is for a dose increase, new dose does not exceed one of the following (a or b):
 - a. Signifor (i and ii):
 - i. 1.8 mg per day;
 - ii. 2 ampules per day;

CLINICAL POLICY

Pasireotide



- b. Signifor LAR (i and ii):
 - i. 40 mg every 4 weeks;
 - ii. 1 vial every 4 weeks.

Approval duration: 12 months

Formatted: Font: Bold

C. Other diagnoses/indications (must meet 1 or 2):

- 1. If this drug has recently (within the last 6 months) undergone a label change (e.g., newly approved indication, age expansion, new dosing regimen) that is not yet reflected in this policy, refer to L A one of the following policies (a or b):
 - a.1. For drugs on the formulary (commercial, health insurance marketplace) or PDL (Medicaid), the no coverage criteria policy for the relevant line of business: CP.CPA.190 for commercial, HIM.PA.33 for health insurance marketplace, and CP.PMN.255 for Medicaid; or
 - b. For drugs NOT on the formulary (commercial, health insurance marketplace) or PDL (Medicaid), the non formulary policy for the relevant line of business: CP.CPA.190 for commercial, HIM.PA.103 for health insurance marketplace, and CP.PMN.16 for Medicaid; or
- 2. If the requested use (e.g., diagnosis, age, dosing regimen) is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized) AND criterion 1 above does not apply, refer to the off-label use policy L A for the relevant line of business: CP.CPA.09 for commercial, HIM.PA.154 for health insurance marketplace, and CP.PMN.53 for Medicaid.

III. Diagnoses/Indications for which coverage is NOT authorized:

- A. Non-FDA approved indications, which are not addressed in this policy, unless there is sufficient documentation of efficacy and safety according to the off label use policy yes LACP.CPA.09 for commercial, HIM.PA.154 for health insurance marketplace, and CP.PMN.53 for Medicaid or evidence of coverage documents.

IV-A. Appendices/General Information

Appendix A: Abbreviation/Acronym Key

FDA: Food and Drug Administration

GH: growth hormone

IGF-I: insulin-like growth factor

Appendix B: Therapeutic Alternatives

Not applicable

Appendix C: Contraindications/Boxed Warnings

None reported

Appendix D: General Information

- Treatment response for Cushing's disease may be defined as reduction in 24-hour urinary free cortisol (UFC) levels and/or improvement in signs or symptoms of the disease. Maximum urinary free cortisol reduction is typically seen by two months of treatment.

Formatted: Indent: Left: 0.5"

Formatted: No page break before

Formatted: Indent: Left: 0.25", Numbered + Level: 1 + Numbering Style: A, B, C, ... + Start at: 1 + Alignment: Left + Aligned at: 0.75" + Indent at: 1"

CLINICAL POLICY

Pasireotide



- Examples of treatment response to acromegaly therapy (including somatostatin analogs, surgical resection or pituitary irradiation) include improvement from baseline in or normalization of GH and/or age- and sex-adjusted IGF-I serum concentrations, or tumor mass control.

V.IV. Dosage and Administration

Drug Name	Indication	Dosing Regimen	Maximum Dose
Pasireotide (Signifor)	Cushing's disease	Initial: 0.6 mg or 0.9 mg SC BID Recommended dosing range: 0.3 mg to 0.9 mg SC BID	1.8 mg/day
Pasireotide (Signifor LAR)*	Cushing's disease	10 mg to 40 mg IM every 4 weeks	40 mg/4 weeks
	Acromegaly	40 mg to 60 mg IM every 4 weeks	60 mg/4 weeks

*Signifor LAR must be administered by a healthcare professional

VI.V. Product Availability

Drug Name	Availability
Pasireotide (Signifor)	Single-dose ampules for injection: 0.3 mg/mL, 0.6 mg/mL, 0.9 mg/mL
Pasireotide (Signifor LAR)	Vials for reconstitution and injectable suspension: 10 mg, 20 mg, 30 mg, 40 mg, 60 mg

VII.VI. References

1. Signifor Prescribing Information. East Hanover, NJ: Novartis Pharmaceuticals Corporation; ~~March 2020~~July 2024. Available at: <https://www.signifor.com/pdf/signifor-pi.pdf>. Accessed ~~July 15, 2024~~10, 2025.
2. Signifor LAR Prescribing Information. East Hanover, NJ: Novartis Pharmaceuticals Corporation; July 2024. Available at: <https://signiforlar.com/wp-content/themes/signifor-lar-theme-hcp/dist/pdf/signifor-lar-pi.pdf>. Accessed July ~~25, 2024~~10, 2025.
3. Melmed S, Bronstein MD, Chanson P. A Consensus Statement on acromegaly therapeutic outcomes. Nat Rev Endocrinol. 2018 Sep;14(9):552-561.
4. Katzenelson L, Laws ER, Melmed S, et al. Acromegaly: An Endocrine Society clinical practice guideline. J Clin Endocrinol Metab. 2014; 99(11): 3933-3951.
5. Fleseriu M, Biller BMK, Freda PU, et al. A Pituitary Society update to acromegaly management guidelines. Pituitary. 2021; 24: 1-13.
6. Giustina A, Biermasz N, Casanueva FF, et al; Acromegaly Consensus Group (ACG). Consensus on criteria for acromegaly diagnosis and remission. Pituitary. 2024 Feb;27(1):7-22. doi: 10.1007/s11102-023-01360-1.
7. Giustina A, Barkhoudarian G, Beckers A, et al. Multidisciplinary management of acromegaly: A consensus. Rev Endocr Metab Disord. 2020; 21(4): 667-678.
8. Nieman LK, Biller BMK, Findling JW, et al. Treatment of Cushing's syndrome: An Endocrine Society clinical practice guideline. J Clin Endocrinol Metab. 2015; 100(8): 2807-2831.
9. Fleseriu M, Auchus R, Bancos I, et al. Consensus on diagnosis and management of Cushing's disease: a guideline update. Lancet Diabetes Endocrinol. 2021; 9(12): 847-875.

CLINICAL POLICY

Pasireotide



Coding Implications

Codes referenced in this clinical policy are for informational purposes only. Inclusion or exclusion of any codes does not guarantee coverage. Providers should reference the most up-to-date sources of professional coding guidance prior to the submission of claims for reimbursement of covered services.

HCPCS Codes	Description
J2502	Injection, pasireotide long acting, 1 mg
J3490	Unclassified drugs

Reviews, Revisions, and Approvals	Date	LDH Approval Date
Converted corporate to local policy.	06.19.23	10.05.23
Annual review: no significant changes; added J3490 code for Signifor; references reviewed and updated.	04.09.24	07.10.24
For acromegaly, revised initial criteria from "(GH) level ≥ 1 $\mu\text{g/mL}$ " to "(GH) level $\geq 1 \mu\text{g/L}$ " per PS/ES practice guidelines and ACG; references reviewed and updated.	12.03.24	<u>02.24.25</u>
<u>Annual review: no significant changes; for acromegaly, extended initial approval duration from 6 months to 12 months; for Cushing's disease, extended initial approval duration from 6 months to 12 months; references reviewed and updated</u>	<u>11.19.25</u>	

Important Reminder

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program approval status; evidence-based guidelines and positions of leading national health professional organizations; views of physicians practicing in relevant clinical areas affected by this clinical policy; and other available clinical information. LHCC makes no representations and accepts no liability with respect to the content of any external information used or relied upon in developing this clinical policy. This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved.

The purpose of this clinical policy is to provide a guide to medical necessity, which is a component of the guidelines used to assist in making coverage decisions and administering benefits. It does not constitute a contract or guarantee regarding payment or results. Coverage decisions and the administration of benefits are subject to all terms, conditions, exclusions, and limitations of the coverage documents (e.g., evidence of coverage, certificate of coverage, policy, contract of insurance, etc.), as well as to state and federal requirements and applicable LHCC administrative policies and procedures.

This clinical policy is effective as of the date determined by LHCC. The date of posting may not be the effective date of this clinical policy. This clinical policy may be subject to applicable legal

CLINICAL POLICY

Pasireotide



and regulatory requirements relating to provider notification. If there is a discrepancy between the effective date of this clinical policy and any applicable legal or regulatory requirement, the requirements of law and regulation shall govern. LHCC retains the right to change, amend or withdraw this clinical policy, and additional clinical policies may be developed and adopted as needed, at any time.

This clinical policy does not constitute medical advice, medical treatment, or medical care. It is not intended to dictate to providers how to practice medicine. Providers are expected to exercise professional medical judgment in providing the most appropriate care, and are solely responsible for the medical advice and treatment of members. This clinical policy is not intended to recommend treatment for members. Members should consult with their treating physician in connection with diagnosis and treatment decisions.

Providers referred to in this clinical policy are independent contractors who exercise independent judgment and over whom LHCC has no control or right of control. Providers are not agents or employees of LHCC.

This clinical policy is the property of LHCC. Unauthorized copying, use, and distribution of this clinical policy or any information contained herein are strictly prohibited. Providers, members, and their representatives are bound to the terms and conditions expressed herein through the terms of their contracts. Where no such contract exists, providers, members and their representatives agree to be bound by such terms and conditions by providing services to members and/or submitting claims for payment for such services.

©20254 Louisiana Healthcare Connections. All rights reserved. All materials are exclusively owned by Louisiana Healthcare Connections and are protected by United States copyright law and international copyright law. No part of this publication may be reproduced, copied, modified, distributed, displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise published without the prior written permission of Louisiana Healthcare Connections. You may not alter or remove any trademark, copyright or other notice contained herein. Louisiana Healthcare Connections is a registered trademark exclusively owned by Louisiana Healthcare Connections.