

## Clinical Policy: Mitoxantrone

Reference Number: LA.PHAR.258

Effective Date: 11.04.23

Last Review Date: 04.22.24 ~~06.15.23~~

Line of Business: Medicaid

[Coding Implications](#)

[Revision Log](#)

See [Important Reminder](#) at the end of this policy for important regulatory and legal information.

**\*\*Please note:** This policy is for medical benefit\*\*

### Description

Mitoxantrone is a synthetic antineoplastic anthracenedione.

### FDA Approved Indication(s)

Mitoxantrone is indicated for:

- Reducing neurologic disability and/or the frequency of clinical relapses in patients with secondary (chronic) progressive, progressive relapsing, or worsening relapsing-remitting multiple sclerosis (MS) (i.e., patients whose neurologic status is significantly abnormal between relapses)
- Treatment of patients with pain related to advanced hormone-refractory prostate cancer as initial chemotherapy in combination with corticosteroids
- Initial therapy of acute nonlymphocytic leukemia (ANLL) (including myelogenous, promyelocytic, monocytic, and erythroid acute leukemias) in adults in combination with other approved drug(s)

Limitation(s) of use: Mitoxantrone is not indicated in the treatment of patients with primary progressive MS.

### Policy/Criteria

*Provider must submit documentation (such as office chart notes, lab results or other clinical information) supporting that member has met all approval criteria.*

It is the policy of Louisiana Healthcare Connections that mitoxantrone is **medically necessary** when the following criteria are met:

#### I. Initial Approval Criteria

##### A. Multiple Sclerosis (must meet all):

1. Diagnosis of one of the following (a or b):
  - a. Relapsing-remitting MS, and failure of all of the following at up to maximally indicated doses, unless clinically significant adverse effects are experienced or all are contraindicated (i, ii, iii, and iv):\*
    - i. **Dimethyl fumarate** (generic Tecfidera®);
    - ii. **Teriflunomide** (generic Aubagio®);
    - iii. **Fingolimod** (Gilenya®);

Formatted: Font: Not Bold

Formatted: Font: Not Bold

- iv. An **interferon-beta agent** (Avonex<sup>®</sup>, Betaseron<sup>®</sup>/Extavia<sup>®†</sup>, Rebif<sup>®</sup>, or Plegridy<sup>®</sup>) or **glatiramer** (Copaxone<sup>®</sup>, Glatopa<sup>®</sup>);

*\*Prior authorization ~~is~~may be required for all disease modifying therapies for MS*

*†Extavia is preferred for the Medicaid ~~line of business~~*

- b. Secondary progressive MS;
2. Prescribed by or in consultation with a neurologist;
  3. Age  $\geq 18$  years;
  4. Mitoxantrone is not prescribed concurrently with other disease modifying therapies for MS (*see Appendix D*);
  5. Documentation of both baseline number of relapses per year and expanded disability status scale (EDSS) score;
  6. Dose does not exceed the following (a and b):
    - a. 12 mg/m<sup>2</sup> every 3 months;
    - b. Total cumulative lifetime dose of 140 mg/m<sup>2</sup>.

**Approval duration: 6 months**

**B. Prostate Cancer** (must meet all):

1. Diagnosis of advanced or metastatic prostate cancer;
2. Prescribed by or in consultation with an oncologist;
3. Age  $\geq 18$  years;
4. Disease is hormone-refractory (i.e., castration-resistant);
5. Mitoxantrone is prescribed concurrently with a corticosteroid (e.g., prednisone);
6. Request meets one of the following (a or b):\*
  - a. Dose does not exceed 14 mg/m<sup>2</sup> every 21 days;
  - b. Dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*);
7. Total cumulative lifetime dose does not exceed 140 mg/m<sup>2</sup>.

*\*Prescribed regimen must be FDA-approved or recommended by NCCN.*

**Approval duration: 6 months**

**C. Acute Nonlymphocytic Leukemia** (must meet all):

1. Diagnosis of ANLL (including myelogenous [i.e., acute myelogenous leukemia], promyelocytic, monocytic, and erythroid acute leukemias);
2. Prescribed by or in consultation with an oncologist or hematologist;
3. Age  $\geq 18$  years;
4. Mitoxantrone is prescribed in combination with other therapies for the diagnosis;
5. Request meets one of the following (a or b):\*
  - a. Dose does not exceed 12 mg/m<sup>2</sup> per infusion;
  - b. Dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*);
6. Total cumulative lifetime dose does not exceed 140 mg/m<sup>2</sup>.

*\*Prescribed regimen must be FDA-approved or recommended by NCCN.*

**Approval duration: 6 months**

**D. Lymphoma (off-label)** (must meet all):

1. Diagnosis of one of the following (a, b, or c):
  - a. Classical Hodgkin lymphoma, and both (i and ii):

- i. Refractory to at least 3 prior lines of therapy;
    - ii. Prescribed as a component of MINE (mesna, ifosfamide, mitoxantrone, and etoposide);
  - b. One of the following B-cell lymphomas: follicular lymphoma, diffuse large B-cell lymphoma, high grade B-cell lymphoma, HIV-related B-cell lymphoma, or post-transplant lymphoproliferative disorder; and both (i and ii):
    - i. Prescribed as second line and subsequent therapy;
    - ii. Prescribed as a component of MINE (mesna, ifosfamide, mitoxantrone, and etoposide);
  - c. Symptomatic T-cell prolymphocytic leukemia as a component of FMC (fludarabine, mitoxantrone, and cyclophosphamide);
2. Prescribed by or in consultation with an oncologist or hematologist;
  3. Age  $\geq$  18 years;
  4. Dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*);\*  
*\*Prescribed regimen must be FDA-approved or recommended by NCCN.*
  5. Total cumulative lifetime dose does not exceed 140 mg/m<sup>2</sup>.

**Approval duration: 6 months**

**E. Acute Lymphoblastic Leukemia (off-label) (must meet all):**

1. Diagnosis of acute lymphoblastic leukemia (ALL);
2. Prescribed by or in consultation with an oncologist or hematologist;
3. Member meets one of the following (a or b):
  - a. Age  $\geq$  18 years, and both of the following (i and ii):
    - i. One of the following (1 or 2):
      - 1) Disease is Philadelphia chromosome (Ph)-negative T-ALL or B-ALL, and relapsed or refractory;
      - 2) Disease is Ph-positive B-ALL, and refractory to tyrosine kinase inhibitor therapy (e.g., dasatinib, imatinib, ponatinib, nilotinib, bosutinib);
    - ii. Mitoxantrone is prescribed as a component of an alkylator combination regimen (e.g., etoposide, ifosfamide, and mitoxantrone) or FLAM (fludarabine, cytarabine, and mitoxantrone);
  - b. Age < 18 years, and one of the following (i, ii, or iii):
    - i. Relapsed/refractory Ph-negative B-ALL;
    - ii. Relapsed/refractory Ph-positive B-ALL in combination with dasatinib or imatinib;
    - iii. Relapsed/refractory T-ALL as a component of UKALL R3 Block 1 (dexamethasone, mitoxantrone, pegaspargase, and vincristine);
4. Dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*);\*  
*\*Prescribed regimen must be FDA-approved or recommended by NCCN.*
5. Total cumulative lifetime dose does not exceed 140 mg/m<sup>2</sup>.

**Approval duration: 6 months**

**F. Other diagnoses/indications (must meet 1 or 2):**

**CLINICAL POLICY**  
**Mitoxantrone**



1. If this drug has recently (within the last 6 months) undergone a label change (e.g., newly approved indication, age expansion, new dosing regimen) that is not yet reflected in this policy, refer to LA.PMN.255
2. If the requested use (e.g., diagnosis, age, dosing regimen) is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized) AND criterion 1 above does not apply, refer to the off-label use policy for the relevant line of business: LA.PMN.53 for Medicaid.

**II. Continued Therapy**

**A. Multiple Sclerosis** (must meet all):

1. Currently receiving medication via Louisiana Healthcare Connections benefit or member has previously met initial approval criteria;
2. Member meets one of the following (a or b):
  - a. If member has received < 1 year of total treatment: Member is responding positively to therapy;
  - b. If member has received  $\geq 1$  year of total treatment: Member meets one of the following (i, ii, iii, or iv):
    - i. Member has not had an increase in the number of relapses per year compared to baseline;
    - ii. Member has not had  $\geq 2$  new MRI-detected lesions;
    - iii. Member has not had an increase in EDSS score from baseline;
    - iv. Medical justification supports that member is responding positively to therapy;
3. Mitoxantrone is not prescribed concurrently with other disease modifying therapies for MS (*see Appendix D*);
4. If request is for a dose increase, new dose does not exceed the following (a and b):
  - a. 12 mg/m<sup>2</sup> every 3 months;
  - b. Total cumulative lifetime dose of 140 mg/m<sup>2</sup>.

**Approval duration: 6 months**

**B. All Other Indications in Section I** (must meet all):

1. Currently receiving medication via Louisiana Healthcare Connections benefit or documentation supports that member is currently receiving mitoxantrone for an oncology indication listed in Section I;
2. Member is responding positively to therapy;
3. If request is for a dose increase, request meets one of the following (a, b, or c):\*
  - a. Prostate cancer: New dose does not exceed 14 mg/m<sup>2</sup> every 21 days;
  - b. ANLL: New dose does not exceed 12 mg/m<sup>2</sup> per infusion;
  - c. Any indication: New dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*);

*\*Prescribed regimen must be FDA-approved or recommended by NCCN.*

4. Total cumulative lifetime dose does not exceed 140 mg/m<sup>2</sup>.

**Approval duration: 12 months**

Formatted: Font: Bold

**C. Other diagnoses/indications** (must meet 1 or 2):

1. If this drug has recently (within the last 6 months) undergone a label change (e.g., newly approved indication, age expansion, new dosing regimen) that is not yet reflected in this policy, refer to LA.PMN.255
2. If the requested use (e.g., diagnosis, age, dosing regimen) is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized) AND criterion 1 above does not apply, refer to the off-label use policy for the relevant line of business: LA.PMN.53 for Medicaid.

**III. Diagnoses/Indications for which coverage is NOT authorized:**

- A. Non-FDA approved indications, which are not addressed in this policy, unless there is sufficient documentation of efficacy and safety according to the off label use policy – LA.PMN.53 for Medicaid or evidence of coverage documents;
- B. Primary progressive MS.

**IV. Appendices/General Information**

*Appendix A: Abbreviation/Acronym Key*

ALL: acute lymphoblastic leukemia	MS: multiple sclerosis
ANLL: acute nonlymphocytic leukemia	NCCN: National Comprehensive Cancer Network
EDSS: expanded disability status scale	Ph: Philadelphia chromosome
FDA: Food and Drug Administration	

*Appendix B: Therapeutic Alternatives*

*This table provides a listing of preferred alternative therapy recommended in the approval criteria. The drugs listed here may not be a formulary agent ~~for all relevant lines of business~~ and may require prior authorization.*

Drug Name	Dosing Regimen	Dose Limit/ Maximum Dose
<del>Aubagio® (teriflunomide)</del> (Aubagio®)	7 mg or 14 mg PO QD	14 mg/day
Avonex®, Rebif® (interferon beta-1a)	Avonex: 30 mcg IM Q week Rebif: 22 mcg or 44 mcg SC TIW	Avonex: 30 mcg/week Rebif: 44 mcg TIW
Plegridy® (peginterferon beta-1a)	125 mcg SC Q2 weeks	125 mcg/2 weeks
Betaseron®, Extavia® (interferon beta-1b)	250 mcg SC QOD	250 mg QOD
glatiramer acetate (Copaxone®, Glatopa®)	20 mg SC QD or 40 mg SC TIW	20 mg/day or 40 mg TIW
<del>Gilenya® (fingolimod)</del> (Gilenya®)	0.5 mg PO QD	0.5 mg/day
dimethyl fumarate (Tecfidera®)	120 mg PO BID for 7 days, followed by 240 mg PO BID	480 mg/day

*Therapeutic alternatives are listed as Brand name® (generic) when the drug is available by brand name only and generic (Brand name®) when the drug is available by both brand and generic.*

*Appendix C: Contraindications/Boxed Warnings*

## CLINICAL POLICY

### Mitoxantrone



- Contraindication(s): prior hypersensitivity to mitoxantrone
- Boxed warning(s): cardiotoxicity, secondary leukemia

#### Appendix D: General Information

- Disease-modifying therapies for MS are: glatiramer acetate (Copaxone<sup>®</sup>, Glatopa<sup>®</sup>), interferon beta-1a (Avonex<sup>®</sup>, Rebif<sup>®</sup>), interferon beta-1b (Betaseron<sup>®</sup>, Extavia<sup>®</sup>), peginterferon beta-1a (Plegridy<sup>®</sup>), dimethyl fumarate (Tecfidera<sup>®</sup>), diroximel fumarate (Vumerity<sup>®</sup>), monomethyl fumarate (Bafiertam<sup>™</sup>), fingolimod (Gilenya<sup>®</sup>, Tascenso ODT<sup>™</sup>), teriflunomide (Aubagio<sup>®</sup>), alemtuzumab (Lemtrada<sup>®</sup>), mitoxantrone (Novantrone<sup>®</sup>), natalizumab (Tysabri<sup>®</sup>), ocrelizumab (Ocrevus<sup>®</sup>), cladribine (Mavenclad<sup>®</sup>), siponimod (Mayzent<sup>®</sup>), ozanimod (Zeposia<sup>®</sup>), ponesimod (Ponvory<sup>™</sup>), ublituximab-xiiy (Briumvi<sup>™</sup>), and ofatumumab (Kesimpta<sup>®</sup>).
- Mitoxantrone has Drugdex IIa recommendations for use in anthracycline-resistant breast cancer, liver cancer, and ovarian cancer; however, these indications are not supported by the National Comprehensive Cancer Network (NCCN). Of note, use of mitoxantrone in invasive breast cancer is actually listed as a use no longer recommended by the NCCN.
- Per the NCCN, prostate cancer that stops responding to traditional androgen deprivation therapy (i.e., hormone therapy) is categorized as castration-recurrent (also known as castration-resistant).

## V. Dosage and Administration

Indication	Dosing Regimen	Maximum Dose
Relapsing MS	12 mg/m <sup>2</sup> given as a short (approximately 5 to 15 minutes) intravenous infusion every 3 months	Cumulative lifetime dose of $\geq 140$ mg/m <sup>2</sup>
Hormone-refractory prostate cancer	12 to 14 mg/m <sup>2</sup> given as a short intravenous infusion every 21 days	Cumulative lifetime dose of $\geq 140$ mg/m <sup>2</sup>
ANLL	Induction: 12 mg/m <sup>2</sup> of mitoxantrone injection (concentrate) daily on Days 1 to 3 given as an intravenous infusion. A second induction course (2 days) may be given if there is an incomplete antileukemic response Consolidation: 12 mg/m <sup>2</sup> given by intravenous infusion daily on Days 1 and 2	Cumulative lifetime dose of $\geq 140$ mg/m <sup>2</sup>

## VI. Product Availability

Multidose vial: 20 mg/10 mL, 25 mg/12.5 mL, 30 mg/15 mL

## VII. References

1. Mitoxantrone Prescribing Information. Lake Forest, IL: Hospira Inc.; April 2021. Available at <http://labeling.pfizer.com/ShowLabeling.aspx?id=4536>. Accessed January 31, 2023.
2. Goodin DS, Frohman EM, Garmany GP, et al. Disease modifying therapies in multiple sclerosis: Subcommittee of the American Academy of Neurology and the MS Council for Clinical Practice Guidelines. *Neurology*. 2002; 58(2): 169-178.
3. National Comprehensive Cancer Network Drugs and Biologics Compendium. Available at: [http://www.nccn.org/professionals/drug\\_compendium](http://www.nccn.org/professionals/drug_compendium). Accessed January 31, 2023.

## CLINICAL POLICY

### Mitoxantrone



4. Rae-Grant A, Day GS, Marrie RA, et al. Practice guideline recommendations summary: disease-modifying therapies for adults with multiple sclerosis: report of the Guideline Development, Dissemination, and Implementation Subcommittee of the American Academy of Neurology. *Neurology*. 2018; 90(17): 777-788. Full guideline available at: <https://www.aan.com/Guidelines/home/GetGuidelineContent/904>. Reaffirmed on September 18, 2021.

#### Coding Implications

Codes referenced in this clinical policy are for informational purposes only. Inclusion or exclusion of any codes does not guarantee coverage. Providers should reference the most up-to-date sources of professional coding guidance prior to the submission of claims for reimbursement of covered services.

HCPSC Codes	Description
J9293	Injection, mitoxantrone HCl, per 5 mg

Reviews, Revisions, and Approvals	Date	LDH Approval Date
Converted corporate to local policy.	06.15.23	<u>10.05.23</u>
<u>Added generic references to Aubagio and Gilenya redirections.</u>	<u>04.22.24</u>	

#### Important Reminder

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program approval status; evidence-based guidelines and positions of leading national health professional organizations; views of physicians practicing in relevant clinical areas affected by this clinical policy; and other available clinical information. LHCC makes no representations and accepts no liability with respect to the content of any external information used or relied upon in developing this clinical policy. This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved.

The purpose of this clinical policy is to provide a guide to medical necessity, which is a component of the guidelines used to assist in making coverage decisions and administering benefits. It does not constitute a contract or guarantee regarding payment or results. Coverage decisions and the administration of benefits are subject to all terms, conditions, exclusions and limitations of the coverage documents (e.g., evidence of coverage, certificate of coverage, policy, contract of insurance, etc.), as well as to state and federal requirements and applicable LHCC-level administrative policies and procedures.

This clinical policy is effective as of the date determined by LHCC. The date of posting may not be the effective date of this clinical policy. This clinical policy may be subject to applicable legal and regulatory requirements relating to provider notification. If there is a discrepancy between the effective date of this clinical policy and any applicable legal or regulatory requirement, the requirements of law and regulation shall govern. LHCC retains the right to change, amend or

## CLINICAL POLICY

### Mitoxantrone



withdraw this clinical policy, and additional clinical policies may be developed and adopted as needed, at any time.

This clinical policy does not constitute medical advice, medical treatment or medical care. It is not intended to dictate to providers how to practice medicine. Providers are expected to exercise professional medical judgment in providing the most appropriate care, and are solely responsible for the medical advice and treatment of members. This clinical policy is not intended to recommend treatment for members. Members should consult with their treating physician in connection with diagnosis and treatment decisions.

Providers referred to in this clinical policy are independent contractors who exercise independent judgment and over whom LHCC has no control or right of control. Providers are not agents or employees of LHCC.

This clinical policy is the property of LHCC. Unauthorized copying, use, and distribution of this clinical policy or any information contained herein are strictly prohibited. Providers, members and their representatives are bound to the terms and conditions expressed herein through the terms of their contracts. Where no such contract exists, providers, members and their representatives agree to be bound by such terms and conditions by providing services to members and/or submitting claims for payment for such services.

©2024<sup>13</sup> Louisiana Healthcare Connections. All rights reserved. All materials are exclusively owned by Louisiana Healthcare Connections and are protected by United States copyright law and international copyright law. No part of this publication may be reproduced, copied, modified, distributed, displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise published without the prior written permission of Louisiana Healthcare Connections. You may not alter or remove any trademark, copyright or other notice contained herein. Louisiana Healthcare Connections is a registered trademark exclusively owned by Louisiana Healthcare Connections.