

Revision Log

Clinical Policy: Mogamulizumab-kpkc (Poteligeo)

Reference Number: LA.PHAR.139

Effective Date: 09.15.22

Last Review Date: <u>04.10.24</u> <u>06.02.23</u>

Line of Business: Medicaid

See Important Reminder at the end of this policy for important regulatory and legal information.

Please note: This policy is for medical benefit

Description

Mogamulizumab-kpkc (Poteligeo®) is a CC chemokine receptor type 4 (CCR4)-directed monoclonal antibody.

FDA Approved Indication(s)

Poteligeo is indicated for the treatment of adult patients with relapsed or refractory mycosis fungoides (MF) or Sézary syndrome (SS) after at least one prior systemic therapy.

Policy/Criteria

Provider must submit documentation (such as office chart notes, lab results or other clinical information) supporting that member has met all approval criteria.

It is the policy of Louisiana Healthcare Connections that Poteligeo is **medically necessary** when the following criteria are met:

I. Initial Approval Criteria

- A. Mycosis Fungoides/Sézary Syndrome (must meet all):
 - 1. Diagnosis of MF or SS;
 - 2. Prescribed by or in consultation with an oncologist or hematologist;
 - 3. Age \geq 18 years;
 - 4. Request meets one of the following (a or b):*
 - a. Dose does not exceed 1 mg/kg on days 1, 8, 15, and 22 of the first 28-day cycle and on days 1 and 15 of each subsequent cycle;
 - b. Dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).

*Prescribed regimen must be FDA-approved or recommended by NCCN

Approval duration: 6 months

B. Adult T-Cell Leukemia/Lymphoma (off-label) (must meet all):

- $1. \ \ Diagnosis\ of\ adult\ T-cell\ leukemia/lymphoma\ (ATLL);$
- 2. Prescribed by or in consultation with an oncologist or hematologist;
- 3. Age \geq 18 years;
- Failure of first-line therapy (see Appendix B for examples);*
 *Prior authorization may be required.

Formatted: Font: Bold



 Dose is within FDA maximum limit for any FDA-approved indication or is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (prescriber must submit supporting evidence).*

*Prescribed regimen must be FDA-approved or recommended by NCCN.

Approval duration: 6 months

C. Other diagnoses/indications (must meet 1 or 2):

- If this drug has recently (within the last 6 months) undergone a label change (e.g., newly approved indication, age expansion, new dosing regimen) that is not yet reflected in this policy, refer to LA.PMN.255
- 2. If the requested use (e.g., diagnosis, age, dosing regimen) is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized) AND criterion 1 above does not apply, refer to the off-label use policy LA.PMN.53.

II. Continued Therapy

A. All Indications in Section I (must meet all):

- 1. Currently receiving medication via Louisiana Healthcare Connections benefit, or documentation supports that member is currently receiving Poteligeo for a covered indication and has received this medication for at least one 28-day cycle;
- 2. Member is responding positively to therapy;
- 3. If request is for a dose increase, request meets one of the following (a or b):*
 - a. New dose does not exceed 1 mg/kg on days 1 and 15 of each subsequent 28-day cycle;
 - b. New dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (prescriber must submit supporting evidence).
 *Prescribed regimen must be FDA-approved or recommended by NCCN.

Approval duration:

Medicaid – 12 months

B. Other diagnoses/indications (must meet 1 or 2):

- If this drug has recently (within the last 6 months) undergone a label change (e.g., newly approved indication, age expansion, new dosing regimen) that is not yet reflected in this policy, refer to LA.PMN.255
- 2. If the requested use (e.g., diagnosis, age, dosing regimen) is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized) AND criterion 1 above does not apply, refer to the off-label use policy LA.PMN.53

III. Diagnoses/Indications for which coverage is NOT authorized:

A. Non-FDA approved indications, which are not addressed in this policy, unless there is sufficient documentation of efficacy and safety according to the off label use policy LA.PMN.53

IV. Appendices/General Information

Appendix A: Abbreviation/Acronym Key ATLL: adult T-cell leukemia/lymphoma CCR4: CC chemokine receptor type 4 CTCL: cutaneous T-cell lymphoma

FDA: Food and Drug Administration

MF: mycosis fungoides

Formatted: Font: Bold

Formatted: Indent: Left: 0"

Formatted: Font: Bold
Formatted: Indent: Left: 0.5"

Formatted: Indent: Left: 0", Hanging: 0.13"



NCCN: National Comprehensive Cancer

SS: Sézary syndrome

Network

Appendix B: Therapeutic Alternatives

This table provides a listing of preferred alternative therapy recommended in the approval

Drug Name	Dosing Regimen	Dose Limit/ Maximum Dose
ATLL: examples of first-line therapy:	Varies	Varies
• Brentuximab vedotin + CHP (cyclophosphamide,		
doxorubicin, and prednisone)		
 CHOP (cyclophosphamide, doxorubicin, vincristine, prednisone) 		
 CHOEP (cyclophosphamide, doxorubicin, vincristine, etoposide, prednisone) 		
• Dose-adjusted EPOCH (etoposide, prednisone, vincristine, cyclophosphamide, doxorubicin)		
• HyperCVAD (cyclophosphamide, vincristine, doxorubicin,		
dexamethasone) alternating with high-dose methotrexate and cytarabine		

Therapeutic alternatives are listed as Brand name[®] (generic) when the drug is available by brand name only and generic (Brand name®) when the drug is available by both brand and generic.

Appendix C: Contraindications/Boxed Warnings None reported

V. Dosage and Administration

Indication	Dosing Regimen	Maximum Dose
MF, SS	1 mg/kg IV over at least 60 minutes on days 1, 8, 15, and 22 of the first 28-day cycle and on days 1 and 15 of each subsequent cycle until disease progression or unacceptable toxicity	1 mg/kg/dose

VI. Product Availability

Single-dose vial: 20 mg/5 mL (4 mg/mL)

VII. References

- 1. Poteligeo Prescribing Information. Bedminster, NJ: Kyowa Kirin, Inc.; March 2022/2023. Available at: https://www.poteligeohcp.com.https://www.poteligeohcp.com. Accessed August 08, 202210, 2023.
- 2. National Comprehensive Cancer Network Drugs and Biologics Compendium. Available at: http://www.nccn.org/professionals/drug_compendium. Accessed August 08, 202210, 2023.
- 3. National Comprehensive Cancer Network. Primary Cutaneous Lymphomas Version 2.2022<u>1.2023</u>. Available at:

https://www.ncen.org/professionals/physician_gls/pdf/primary_cutaneous.pdf.https://www.n

Formatted: Indent: Left: 0" Formatted: Indent: Left: 0", Hanging: 0.13"

Formatted Table

Formatted Table



<u>ccn.org/professionals/physician_gls/pdf/primary_cutaneous.pdf.</u> Accessed August 8, 202210, 2023.

National Comprehensive Cancer Network. T-Cell Lymphomas Version 2.20221.2023.
 Available at: https://www.nccn.org/professionals/physician_gls/pdf/t-cell.pdf.
 https://www.nccn.org/professionals/physician_gls/pdf/t-cell.pdf.
 Accessed August 8, 202210, 2023.

Coding Implications

Codes referenced in this clinical policy are for informational purposes only. Inclusion or exclusion of any codes does not guarantee coverage. Providers should reference the most up-to-date sources of professional coding guidance prior to the submission of claims for reimbursement of covered services.

HCPCS Codes	Description
J9204	Injection, mogamulizumab-kpkc, 1 mg

Reviews, Revisions, and Approvals	Date	LDH Approval Date
Converted corporate to local policy.	09.22	09.15.22
Nno significant changes; references reviewed and updated.	06.02.23	10.05.23
Template changes applied to other diagnoses/indications.		
Added verbiage this policy is for medical benefit only.		
Annual review: no significant changes; references reviewed and	04.10.24	
<u>updated.</u>		

Important Reminder

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program approval status; evidence-based guidelines and positions of leading national health professional organizations; views of physicians practicing in relevant clinical areas affected by this clinical policy; and other available clinical information. LHCC makes no representations and accepts no liability with respect to the content of any external information used or relied upon in developing this clinical policy. This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved.

The purpose of this clinical policy is to provide a guide to medical necessity, which is a component of the guidelines used to assist in making coverage decisions and administering benefits. It does not constitute a contract or guarantee regarding payment or results. Coverage decisions and the administration of benefits are subject to all terms, conditions, exclusions and limitations of the coverage documents (e.g., evidence of coverage, certificate of coverage, policy, contract of insurance, etc.), as well as to state and federal requirements and applicable LHCC administrative policies and procedures.



This clinical policy is effective as of the date determined by LHCC. The date of posting may not be the effective date of this clinical policy. This clinical policy may be subject to applicable legal and regulatory requirements relating to provider notification. If there is a discrepancy between the effective date of this clinical policy and any applicable legal or regulatory requirement, the requirements of law and regulation shall govern. LHCC retains the right to change, amend or withdraw this clinical policy, and additional clinical policies may be developed and adopted as needed, at any time.

This clinical policy does not constitute medical advice, medical treatment or medical care. It is not intended to dictate to providers how to practice medicine. Providers are expected to exercise professional medical judgment in providing the most appropriate care, and are solely responsible for the medical advice and treatment of members. This clinical policy is not intended to recommend treatment for members. Members should consult with their treating physician in connection with diagnosis and treatment decisions.

Providers referred to in this clinical policy are independent contractors who exercise independent judgment and over whom LHCC has no control or right of control. Providers are not agents or employees of LHCC.

This clinical policy is the property of LHCC. Unauthorized copying, use, and distribution of this clinical policy or any information contained herein are strictly prohibited. Providers, members and their representatives are bound to the terms and conditions expressed herein through the terms of their contracts. Where no such contract exists, providers, members and their representatives agree to be bound by such terms and conditions by providing services to members and/or submitting claims for payment for such services.

©20243 Louisiana Healthcare Connections. All rights reserved. All materials are exclusively owned by Louisiana Healthcare Connections and are protected by United States copyright law and international copyright law. No part of this publication may be reproduced, copied, modified, distributed, displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise published without the prior written permission of Louisiana Healthcare Connections. You may not alter or remove any trademark, copyright or other notice contained herein. Louisiana Healthcare Connections is a registered trademark exclusively owned by Louisiana Healthcare Connections.