

Clinical Policy: Dostarlimab-gxly (Jemperli)

Reference Number: LA.PHAR.540

Effective Date: 09.29.23

Last Review Date: ~~06.20.25~~09.17.24

Line of Business: Medicaid

[Revision Log](#)

See [Important Reminder](#) at the end of this policy for important regulatory and legal information.

****Please note: This policy is for medical benefit****

Formatted: Font: Bold

Description

Dostarlimab-gxly (Jemperli™) is a programmed death receptor-1 (PD-1)–blocking antibody.

FDA Approved Indication(s)

Jemperli is indicated for:

- **Endometrial cancer (EC)**
 - In combination with carboplatin and paclitaxel, followed by Jemperli as a single agent for the treatment of adult patients with primary advanced or recurrent EC ~~that is mismatch repair deficient (dMMR), as determined by an FDA-approved test, or microsatellite instability-high (MSI-H)~~
 - As a single agent for the treatment of adult patients with mismatch repair deficient (dMMR) recurrent or advanced EC, as determined by an FDA-approved test, that has progressed on or following prior treatment with a platinum-containing regimen in any setting and are not candidates for curative surgery or radiation
- **dMMR recurrent or advanced solid tumors**
 - As a single agent for the treatment of adult patients with dMMR recurrent or advanced solid tumors, as determined by an FDA-approved test, that have progress on or following prior treatment and who have no satisfactory alternative treatment options*

**This indication is approved under accelerated approval based on tumor response rate and durability of response. Continued approval for this indication may be contingent upon verification and description of clinical benefit in a confirmatory trial(s).*

Policy/Criteria

Provider must submit documentation (such as office chart notes, lab results or other clinical information) supporting that member has met all approval criteria.

It is the policy of Louisiana Healthcare Connections that Jemperli is **medically necessary** when the following criteria are met:

I. Initial Approval Criteria**A. Endometrial Carcinoma** (must meet all):

1. Diagnosis of EC;
2. Prescribed by or in consultation with an oncologist;
3. Age ≥ 18 years;
4. One of the following (a or b):

CLINICAL POLICY

Dostarlimab-gxly

- a. Prescribed in combination with carboplatin and paclitaxel for advanced (i.e., stage III-IV) or recurrent disease;
- b. All of the following (i, ii, iii, and iv):
 - i. Disease is recurrent or advanced;
 - ii. Disease is dMMR (i.e., disease is indicative of MMR gene mutation or loss of expression) or microsatellite instability-high (MSI-H);
 - iii. Disease has progressed following prior treatment with a platinum-containing regimen (e.g., carboplatin/cisplatin);
 - iv. Member is not a candidate for curative surgery or radiation;
5. Request meets one of the following (a, b, or c):*
 - a. Combination therapy: Dose does not exceed 500 mg every 3 weeks for 6 dosecycles, in combination with carboplatin and paclitaxel, followed by 1,000 mg monotherapy every 6 weeks starting 3 weeks after dosecycle 6;
 - b. Single-agent use: Dose does not exceed 500 mg every 3 weeks for 4 dosecycles, followed by 1,000 mg every 6 weeks starting 3 weeks after dosecycle 4;
 - c. Dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).

**Prescribed regimen must be FDA-approved or recommended by NCCN*

Approval duration: 6 months

B. Solid Tumor (must meet all):

1. Diagnosis of solid tumor (e.g., ampullary adenocarcinoma, breast cancer, colon cancer [including appendiceal adenocarcinoma], esophageal and esophagogastric junction cancers, gallbladder cancer, gastric cancer, hepatocellular carcinoma, extra/intrahepatic cholangiocarcinoma, occult primary cancer, ovarian/fallopian tube/primary peritoneal cancer, pancreatic adenocarcinoma, rectal cancer, small bowel adenocarcinoma);
2. Prescribed by or in consultation with an oncologist;
3. Age ≥ 18 years;
4. One of the following (a, b, c, or d):
 - a. Disease is metastatic, recurrent, or advanced;
 - b. Gastric cancer only: Disease is early stage or surgically unresectable, or member is not a surgical candidate;
 - c. Colon (including appendiceal adenocarcinoma) or rectal cancer only: Disease is locally unresectable or medically inoperable;
 - d. Esophageal or esophagogastric junction cancer only: Member is not a surgical candidate;
5. One of the following (a, b, or c):
 - a. Disease is dMMR (i.e., disease is indicative of MMR gene mutation or loss of expression);
 - b. Disease is MSI-H;
 - c. Colon (including appendiceal adenocarcinoma) and rectal cancer only: Disease is positive for polymerase epsilon/delta [POLE/POLD1] mutation;

CLINICAL POLICY

Dostarlimab-gxly

6. One of the following (a, b, or c):
 - a. Disease has progressed on or following prior treatment and who have no satisfactory alternative options;
 - b. Request is for palliative therapy for gastric, esophageal, or esophagogastric junction cancer;
 - c. Request is for colon cancer (including appendiceal adenocarcinoma), esophageal or esophagogastric junction cancer with a planned esophagectomy, gastric cancer that is either early stage or surgically unresectable, pancreatic adenocarcinoma, rectal cancer, or small bowel adenocarcinoma;
 7. Prescribed as a single agent;
 8. Request meets one of the following (a or b):*
 - a. Dose does not exceed 500 mg every 3 weeks for 4 ~~doses~~dose cycles, followed by 1,000 mg every 6 weeks starting 3 weeks after ~~dose~~dose cycle 4;
 - b. Dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).
- *Prescribed regimen must be FDA-approved or recommended by NCCN*

Approval duration: 6 months

C. Other diagnoses/indications (must meet 1 or 2):

1. If this drug has recently (within the last 6 months) undergone a label change (e.g., newly approved indication, age expansion, new dosing regimen) that is not yet reflected in this policy, refer to LA.PMN.255
2. If the requested use (e.g., diagnosis, age, dosing regimen) is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized) AND criterion 1 above does not apply, refer to the off-label use policy ~~for the relevant line of business: LA.PMN.53 for Medicaid.~~

II. Continued Therapy

A. All Indications in Section I (must meet all):

1. Currently receiving medication via Louisiana Healthcare Connections benefit, or documentation supports that member is currently receiving Jemperli and has received this medication for at least 30 days;
2. Member is responding positively to therapy;
3. If request is for a dose increase, request meets one of the following (a or b):*
 - a. New dose does not exceed 1,000 mg every 6 weeks;
 - b. New dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).

**Prescribed regimen must be FDA-approved or recommended by NCCN*

Approval duration: 12 months

CLINICAL POLICY

Dostarlimab-gxly

B. Other diagnoses/indications (must meet 1 or 2):

1. If this drug has recently (within the last 6 months) undergone a label change (e.g., newly approved indication, age expansion, new dosing regimen) that is not yet reflected in this policy, refer to LA.PMN.255
2. If the requested use (e.g., diagnosis, age, dosing regimen) is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized) AND criterion 1 above does not apply, refer to the off-label use policy ~~for the relevant line of business: LA.PMN.53 for Medicaid.~~

III. Diagnoses/Indications for which coverage is NOT authorized:

- A. Non-FDA approved indications, which are not addressed in this policy, unless there is sufficient documentation of efficacy and safety according to the off-label use policy ~~yes – LA.PMN.53 for Medicaid or evidence of coverage documents.~~

IV. Appendices/General Information

Appendix A: Abbreviation/Acronym Key

dMMR: mismatch repair deficient

EC: endometrial carcinoma

FDA: Food and Drug Administration

MSI-H: microsatellite instability-high

NCCN: National Comprehensive Cancer Network

POLE/POLD1: polymerase epsilon/delta

Appendix B: Therapeutic Alternatives

This table provides a listing of preferred alternative therapy recommended in the approval criteria. The drugs listed here may not be a formulary agent and may require prior authorization.

Drug Name	Dosing Regimen	Dose Limit/Maximum Dose
EC systemic therapies: carboplatin, cisplatin, carboplatin/paclitaxel, cisplatin/docetaxel, cisplatin/doxorubicin, cisplatin/doxorubicin/paclitaxel, carboplatin/paclitaxel/bevacizumab, carboplatin/paclitaxel/trastuzumab, cisplatin/ifosfamide	Varies	Varies

Therapeutic alternatives are listed as Brand name® (generic) when the drug is available by brand name only and generic (Brand name®) when the drug is available by both brand and generic.

Appendix C: Contraindications/Boxed Warnings

None reported

Formatted: Keep with next

V. Dosage and Administration

CLINICAL POLICY

Dostarlimab-gxly

Indication	Dosing Regimen	Maximum Dose
EC as combination therapy	Dose 1 through 6: 500 mg <u>IV</u> every 3 weeks <u>for 6 cycles</u> in combination with carboplatin and paclitaxel Subsequent dosing beginning 3 weeks after Dose 6 (Dose 7 onwards); followed by 1,000 mg <u>IV</u> as monotherapy every 6 weeks <u>for all cycles thereafter until disease progression, unacceptable toxicity, or up to 3 years</u>	See do ing dosing regimen
EC as single agent therapy; solid tumors	Dose 1 through 4: 500 mg <u>IV</u> every 3 weeks Subsequent dosing beginning 3 weeks after Dose 4 (Dose 5 onwards); for 4 cycles followed by 1,000 mg <u>IV</u> every 6 weeks <u>for all cycles thereafter until disease progression or unacceptable toxicity</u>	See dosing regimen

VI. Product Availability

Single-dose vial: 500 mg/10 mL

VII. References

1. Jemperli Prescribing Information. Philadelphia, PA: GlaxoSmithKline LLC; ~~March~~ August 2024. Available at: <https://jemperli.com/>. Accessed ~~May~~ August 8, 2024.
2. Dostarlimab-hxly In: National Comprehensive Cancer Network Drugs and Biologics Compendium. Available at: http://www.nccn.org/professionals/drug_compendium. Accessed ~~May 15~~ August 8, 2024.
3. Mirza MR, Chase DM, Slomovitz BM, et al. Dostarlimab for primary advanced or recurrent endometrial cancer. N Engl J Med. 2023 Jun 8;388(23):2145-2158. doi: 10.1056/NEJMoa2216334. Epub 2023 Mar 27. PMID: 36972026.

Reviews, Revisions, and Approvals	Date	LDH Approval Date
Policy created.	05.01.23	08.28.23
For EC, added indication and pathway for first-line use when prescribed in combination with carboplatin and paclitaxel for stage III-IV or recurrent disease; for solid tumors, added gallbladder cancer and pancreatic cancer, specified types of hepatobiliary cancers, and added bypass of prior therapies for small bowel adenocarcinoma or pancreatic adenocarcinoma per NCCN; references reviewed and updated.	02.01.24	05.10.24

CLINICAL POLICY

Dostarlimab-gxly

Reviews, Revisions, and Approvals	Date	LDH Approval Date
Annual review: revised solid tumors criteria per NCCN – added additional disease qualifiers of early stage or unresectable for gastric cancer and locally unresectable or medically inoperable for colon and rectal cancers, added pathway to allow members who are not surgical candidates for gastric and esophageal/esophagogastric junction cancers, added POLE/POLD1 mutation for colon and rectal cancers, and added bypass of prior therapies for colon cancer, esophageal/esophagogastric junction cancer with planned esophagectomy or if request is for palliative therapy, gastric cancer that is early stage or surgically unresectable or if request is for palliative therapy, and rectal cancer; references reviewed and updated.	09.17.24	<u>01.02.25</u>
<u>Updated FDA approved indication to remove requirement for disease to be dMMR/MSI-H when prescribed in combination with carboplatin and paclitaxel per expanded label, and clarified in criteria that stage III-IV is advanced</u>	<u>06.20.25</u>	

Important Reminder

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program approval status; evidence-based guidelines and positions of leading national health professional organizations; views of physicians practicing in relevant clinical areas affected by this clinical policy; and other available clinical information. LHCC makes no representations and accepts no liability with respect to the content of any external information used or relied upon in developing this clinical policy. This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved.

The purpose of this clinical policy is to provide a guide to medical necessity, which is a component of the guidelines used to assist in making coverage decisions and administering benefits. It does not constitute a contract or guarantee regarding payment or results. Coverage decisions and the administration of benefits are subject to all terms, conditions, exclusions, and limitations of the coverage documents (e.g., evidence of coverage, certificate of coverage, policy, contract of insurance, etc.), as well as to state and federal requirements and applicable LHCC administrative policies and procedures.

This clinical policy is effective as of the date determined by LHCC. The date of posting may not be the effective date of this clinical policy. This clinical policy may be subject to applicable legal and regulatory requirements relating to provider notification. If there is a discrepancy between the effective date of this clinical policy and any applicable legal or regulatory requirement, the

CLINICAL POLICY

Dostarlimab-gxly

requirements of law and regulation shall govern. LHCC retains the right to change, amend or withdraw this clinical policy, and additional clinical policies may be developed and adopted as needed, at any time.

This clinical policy does not constitute medical advice, medical treatment, or medical care. It is not intended to dictate to providers how to practice medicine. Providers are expected to exercise professional medical judgment in providing the most appropriate care, and are solely responsible for the medical advice and treatment of members. This clinical policy is not intended to recommend treatment for members. Members should consult with their treating physician in connection with diagnosis and treatment decisions.

Providers referred to in this clinical policy are independent contractors who exercise independent judgment and over whom LHCC has no control or right of control. Providers are not agents or employees of LHCC.

This clinical policy is the property of LHCC. Unauthorized copying, use, and distribution of this clinical policy or any information contained herein are strictly prohibited. Providers, members, and their representatives are bound to the terms and conditions expressed herein through the terms of their contracts. Where no such contract exists, providers, members and their representatives agree to be bound by such terms and conditions by providing services to members and/or submitting claims for payment for such services.

©2025⁵⁴ Louisiana Healthcare Connections. All rights reserved. All materials are exclusively owned by Louisiana Healthcare Connections and are protected by United States copyright law and international copyright law. No part of this publication may be reproduced, copied, modified, distributed, displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise published without the prior written permission of Louisiana Healthcare Connections. You may not alter or remove any trademark, copyright or other notice contained herein. Louisiana Healthcare Connections is a registered trademark exclusively owned by Louisiana Healthcare Connections.