

Louisiana Morbidity Report



JOHN BEL EDWARDS
GOVERNOR

Office of Public Health - Infectious Disease Epidemiology Section
P.O. Box 60630, New Orleans, LA 70160 - Phone: (504) 568-8313
www.ldh.louisiana.gov/LMR



Infectious Disease Epidemiology Main Webpage
www.infectiousdisease.dhh.louisiana.gov

REBEKAH E. GEE MD MPH
SECRETARY

September - October, 2016

Volume 27, Number 5

An Unusual Case of *Neisseria* Louisiana, 2016

A 51-year old male presents at an emergency department with fever (104°F, 40°C) and tachycardia (136). His job required him to be outside delivering materials to multiple addresses and to wear a uniform not well adapted to summer temperatures. The presumptive diagnoses were heat exhaustion or early sepsis. Multiple workup tests done did not show any major abnormalities.

After being admitted, a blood culture was done and the patient was treated empirically with ceftriaxone. During the night, his temperature decreased to 101°F and his white blood cell count increased from 9,000 to 16,200. His other lab tests were unremarkable. The patient was well hydrated and was discharged the next day with diagnoses of heat exhaustion, hypertension and hypokalemia.

At 24 hours the blood culture showed no growth, but the next day after discharge a presumptive identification of *Neisseria* species was made and later confirmed by polymerase chain reaction (PCR) as *Neisseria cinerea*.

N. cinerea is part of a group of non-pathogenic *Neisseria* (*N. lactamica*, *N. polysaccharea*, *N. sicca*, *N. subflava*, *N. flavescens*, *N. mucosa*, *N. elongate*) that are colonizers of the human upper respiratory tract.

(continued on page 6)

Sudden Unexpected Infant Death: Louisiana, 2013-2014

Cara Bergo, MPH

The Louisiana infant mortality rate, death before the age of one year, is the third highest in the United States at 8.0 deaths per 1,000 live births compared to the United States at 5.8 deaths. The third leading cause of infant mortality in Louisiana is Sudden Unexpected Infant Death (SUID). SUID is defined by the Centers for Disease Control and Prevention (CDC) as the death of an infant younger than one year of age that cannot be explained after a thorough investigation is conducted, including a full autopsy, examination of the death scene, and a review of the infant's clinical history.

In 2013, there were 100 SUID cases in Louisiana; in 2014, there were 104. With help from the CDC SUID Case Registry Grant, the nine Louisiana Regional Maternal and Child Health (MCH) Coordinators with the Bureau of Family Health collect records regarding these deaths to abstract and review each case. Of the 204 SUID deaths in Louisiana from 2013 and 2014, 199 were abstracted and reviewed within regional case review teams. The abstraction and reviews provide information on demographics, scene of event, risk factors involved, and possible prevention strategies to implement in the future.

There are four types of SUID: Sudden Infant Death Syndrome (92 cases, 46.2%), unknown cause of death (7 cases, 3.5%), accidental suffocation and strangulation in bed (84 cases, 42.2%) and asphyxiation and suffocation (ASSB), (16 cases, 8.0%), (Table 1, Figure).

Table 1: Characteristics of Sudden Unexpected Infant Deaths Louisiana, 2013-2014

Demographics	Count (%)
Age at Death	N=199
<28 days	17 (8.5)
28-112 days	119 (59.8)
113-224 days	52 (26.1)
225-365 days	11 (5.5)
Sex	
Male	91 (45.7)
Female	108 (54.3)
Race/ Ethnicity	
Black, Non-Hispanic	106 (53.3)
White, Non-Hispanic	83 (41.7)
Hispanic	10 (5.0)

(continued on page 4)

Inside

Louisiana Service Worker Wellness Report: Results from the BRFSS, 2013-2014	2
Announcements: NSHN and Emerging Disease Workshops; Updates, IDEpi Webpages	3
National Lead Poisoning Prevention Week: October 23-29, 2016	5
Louisiana Fact: At the Forefront of Modern Dentistry	5
Zika and Animals	6
Zika - Confirmed and Probable Cases; United States, 2016	6

Louisiana Service Worker Wellness Report: Results from the Behavioral Risk Factor Surveillance System, 2013-2014

Jocelyn Lewis, PhD; Vanessa Paul, MPH; Laurie Freyder, MPH; Michelle Lackovic, MPH

The Behavioral Risk Factor Surveillance System (BRFSS) is a cross-sectional annual health-related telephone survey of civilian, non-institutionalized state residents aged 18 years and older. It is coordinated by the Centers for Disease Control and Prevention and conducted by all US states, Washington D.C. and several U.S. territories. The BRFSS uses a multistage sampling design to select a representative sample of the adult population in each state, allowing for state-to-state and national comparisons.

Beginning in 2013, Louisiana added an optional industry and occupation module to collect information from employed respondents about their “type of business/industry” and the “type of work/occupation.” Verbatim responses were collected, coded by the National Institute of Occupational Safety and Health certified coders into 3-digit census occupation codes and then aggregated into broader occupational categories. Using a combined 2013/2014 data set, population-based estimates on the health and well-being of service workers in Louisiana were calculated. For Louisiana and many other states, the BRFSS is the only available data source of timely and accurate data on health-related behaviors.

Service workers represent a broadly defined group that includes: healthcare support, protective service, food service, cleaning and maintenance and personal care and service occupations. Together, these workers make up about 17 percent of Louisiana’s workforce, with food service workers making up almost one-third of all service workers, followed by personal care and service workers at 21 percent. Although there is variation among service occupations, many of the jobs are held by women and minorities and involve shift work, low wages and minimum job security. These jobs, especially service jobs that pay below \$20 per hour and require only a high school education or less, are predicted to grow far more quickly than higher-wage jobs over the next decade. Between 2010 and 2014, the number of food service workers increased approximately 12 percent. The other service occupations also experienced growth during this time period, excluding protective service workers.

Methods

Prevalence estimates for 19 survey responses from the BRFSS were calculated to compare Louisiana service workers with all other workers. These 19 were key questions addressing issues of chronic health and risk behaviors grouped into six main categories: health care access; health status; chronic health conditions; housing and food insecurity; risk factors and behaviors and immunization. The BRFSS criteria for publicly reporting data results are that: 1) each cell size must have a count of at least 50 and 2) the coefficient of variation (CV) of the prevalent estimate must be less than 0.30 to indicate stability. Unreliable estimates are not included in the report. SAS 9.3 was used for all data analyses, and Microsoft Excel was used to create all figures and tables. Rao-Scott chi-squared tests were calculated for all

prevalence estimates to determine differences between occupation groups. Statistical significance for all tests was set at $p < 0.05$.

Results

Table 1 shows the annual average distribution of 2013 and 2014 LA BRFSS Service worker respondents grouped by 2010 US Census Bureau Occupation Codes compared to the Bureau of Labor Statistics’ Current Population Survey (CPS). The CPS is a monthly survey of approximately 60,000 randomly sampled US households representative of the civilian non-institutional population. It includes demographic and labor force and employment data for each household member at least 15 years-old. The five service workers’ occupational categories were combined to create a service group, and all other occupation groups were combined to form the all other workers group for comparison.

Table 1: Average Workforce Distribution of BRFSS Service Worker Respondents by Major Occupation Group - Louisiana, 2013 and 2014

Occupation Group/Service	Example Occupations	Workforce Distribution (percent)	
		BRFSS	Current Population Survey
Healthcare Support	massage therapist, dental assistant, phlebotomist, pharmacy aides	3.1	2.5
Protective Service	detective, fish and game warden, firefighter, police officer, correctional officer, security guard	3.6	2.5
Food Prep and Serving Related	chef, bartender, dishwasher, counter attendant, host and hostess, waiter and waitress	6.7	5.5
Building and Grounds Cleaning and Maintenance	janitor, housekeeper, pest control worker, groundskeeper, lawn service worker	4.3	3.1
Personal Care and Service	barber, animal trainer, hairdresser, usher, mortician, baggage porter, tour guide, childcare attendant	3.8	3.7

(continued on page 3)

Louisiana Morbidity Report	
Volume 27, Number 5	September - October, 2016
<p>The Louisiana Morbidity Report is published bimonthly by the LHH OPH Infectious Disease Epidemiology Section to inform physicians, nurses, and public health professionals about disease trends and patterns in Louisiana. Address correspondence to Louisiana Morbidity Report, Infectious Disease Epidemiology Section, Louisiana Department of Health and Hospitals, P.O. Box 60630, New Orleans, LA 70160.</p>	
<i>Assistant Secretary OPH</i>	<i>Beth Scalco</i>
<i>State Epidemiologist</i>	<i>Raoult Ratard, MD, MPH</i>
<i>Editors</i>	<i>Theresa Sokol, MPH</i> <i>Julie Hand, MSPH</i> <i>Rosemarie Robertson, BS, MT(C), CNMT</i>

(Louisiana Service Worker ... continued from page 2)

Table 2 shows the distribution of socio-demographic characteristics among Louisiana service workers and all other workers combined for Louisiana’s BRFSS 2013 and 2014 data. Service workers in Louisiana were more likely to be female, younger than 45 years-old, African-American, not have a high school education, and have an annual income less than \$50,000 in comparison with other workers.

Table 2: Distribution of Demographic Data Among Service Workers and All Other Workers - BRFSS - Louisiana, 2013 and 2014

Characteristic	Service Occupation Group (N=913)	All Other Workers (N=4,270)
	%	%
Sex		
Male	41.1	56.5
Female	58.9	43.5
Age Group (Years)		
18-44	68.4	54.9
45-54	17.9	23.8
55-64	10.6	15.7
Older than or equal to 65	3.2	5.6
Race/ethnicity		
White non-Hispanic	47.0	66.5
Black non-Hispanic	44.1	26.3
Hispanic	5.7	4.0
Other	3.2	3.3
Education		
Some high school	21.4	11.6
High school graduate	41.9	29.8
Some college	28.4	29.8
College graduate	8.2	28.8
Household Income		
Less than \$50,000	77.3	44.1
Greater or equal to \$50,000	22.7	55.9

Table 3 shows that service workers reported significantly greater prevalences of poor health, chronic health conditions, food and housing insecurity, and risk behaviors than other workers.

Table 3: Significant Prevalence Estimates - BRFSS - Louisiana, 2013 and 2014

BRFSS Variable	Significant Prevalence Estimates (%)		
	Service Workers	All Other Workers	p-value
Health Care Access			
No health care coverage	38.5	18.7	<0.0001
Unable to see doctor due to medical costs	29.5	14.9	<0.0001
Health Status			
Poor total health (physical and/or mental health poor 11-30 days)	20.9	14.9	0.003
Chronic Health Conditions			
Current asthma	7.4	5.0	0.04
Chronic obstructive pulmonary disease	6.0	3.5	0.008
Depressive disorder	17.7	13.2	0.01
Diabetes	8.4	6.3	0.01
Insecurities			
Food insecurity	34.2	19.2	0.0001
Housing/shelter insecurity	17.0	14.3	0.007
Risk Factors and Behaviors			
Current smoking	32.1	23.8	0.0004

Conclusion

This report provides critical information on the health and well-being of service workers in Louisiana that can be used by policymakers, community leaders, and business leaders to better

understand the economic hardships, chronic health conditions, and other quality of life issues faced by this growing occupational sector. This information, in turn, can better inform policy and legislation and health intervention and prevention programs.

Data in this report pre-date Medicaid expansion which, as pointed out in the recently released Louisiana State Health Assessment and Improvement plan, is the single greatest action our state can take to ensure that every working Louisianan has access to health care.

Lastly, for the purposes of presenting data that meet minimum reporting standards, data for the five service occupations were aggregated, which may have resulted in masking occupation-specific issues. As additional years of data are received, a similar analysis can be done for each of the five service sectors. This analysis sets the important foundation for that work.

The full Louisiana Service Worker BRFSS Report can be accessed at http://new.dhh.louisiana.gov/assets/oph/Center-EH/envepi/occ_health/Documents/LA_Service_Worker_Wellness_Report_BRFSS_2016_FINAL.pdf. For more information, please contact Dr. Jocelyn Lewis at Jocelyn.lewis@la.gov.

Announcements

National Healthcare Safety Network/Emerging Infectious Diseases Workshops - 2016

Metairie - November 14 Bossier City - November 16
Alexandria - November 17

This is a one-day workshop sponsored by the Department of Health’s, Office of Public Health, Infectious Disease Epidemiology Section. It is targeted towards infectious disease preventionists that are seeking to reduce the number of hospital acquired infections and prepare for emerging infectious disease threats that may present in their facility.

This workshop is free to attend, but must be registered for because of seating limitations and to provide the adequate number of handouts. Nurse and laboratory education credits have been applied for.

Please go to <http://new.dhh.louisiana.gov/index.cfm/page/2566> for a registration form and more information.

Updates: Infectious Disease Epidemiology (IDEpi) Webpages
www.infectiousdisease.dhh.louisiana.gov

Annual: Brucellosis; Creutzfeldt Jakob Disease; Cryptosporidiosis; Cyclosporiasis; Eastern Equine and La Crosse (California) Encephalitis; Gullain Barre; Hepatitis C; St. Louis Encephalitis

Arboviral: WNV Weekly Report


Epidemiology Manual: Brain-eating Ameba (Naegleria) Public Information; Pertussis; U.S. Laboratories Testing for Zika Virus Infection

Hepatitis: Epidemiologic Profile of Hepatitis C Virus Infection in Louisiana - 2015; What is the Prevalence of Hepatitis C in Louisiana?

Influenza: ACIP Recommendations 2016-17; Monthly Report

(Sudden Unexpected Infant ... continued from page 1)

Figure: Type of SUID by Year: Louisiana, 2004-2014



The majority of deaths (75.6%) occurred before the infant was 4-months old and the deaths were more likely to be female, (54.3%). East Baton Rouge Parish (Region 2*) had 26 deaths in 2013 and 2014, the most of any parish in Louisiana. Out of parishes with at least 10 SUID deaths, East Baton Rouge Parish had the highest incidence rate of SUID at 2.12 deaths per 1,000 live births during 2013-2014 followed by Calcasieu (Region 7) at 2.02 per 1,000 live births, Orleans (Region 1) at 1.72 per 1,000 live births, and Lafayette (Region 4) at 1.58 per 1,000 live births.

Following a SUID case, a death scene investigation is done by the parish coroner’s office or local law enforcement. The scene of the event is investigated, caregiver is interviewed and a doll reenactment is done to better understand the situation surrounding the death. SUID cases were most likely to occur in the child’s home (83.4%) and secondly in a relative’s home (12.1%). The majority of cases had supervision at the time of death (91.5%) and the caregiver at the time of death was usually the female biological parent (97.0%). More than half (59.8%) were receiving social services. (Table 2).

Table 2: Scene of Event of SUID Death: Louisiana, 2013-2014

Factor	Count (%)
Caregiver	
Biological Parent	192 (96.5)
< 20 Years Old	28 (14.1)
Female	193 (97.0)
Unemployed	59 (29.6)
Low Income	105 (52.8)
Receiving Social Services	119 (59.8)
Any Substance Abuse	43 (21.6)
Supervision at Time of Incident Leading to death	182 (91.5)
Incident	
Child's Home	166 (83.4)
Relative's Home	24 (12.1)

The ASSB cause of death has increased over the past decade

partly due to better investigation and reporting surrounding the scene and situation of the death.

There are known sleep environment risk factors associated with SUID including not sleeping on the back, soft bedding, and sleeping on an adult mattress. In Louisiana, about half of cases were sleeping in an adult bed (50.8%) and 48.3% were found sleeping on their stomach or side. Known risk factors of SUID were present in the sleep environment including pillows (44.7%), comforters (40.7%) and other people (65.8%). Many of these deaths (79.4%) were tragically found to be preventable after review due to the ability to mitigate many of these risk factors (Table 3).

Table 3: Sleep Environment Risk Factors: Louisiana, 2013-2014

Factor	Count (%)
Child's activity was sleeping at time of incident	195 (98.0)
Death related to sleeping or the sleep environment	180 (90.5)
Incident Sleep Place	
Adult bed	101 (50.8)
Crib/ bassinette	42 (21.1)
Couch	18 (9.0)
Other	19 (9.5)
Found Position	
Stomach	75 (37.7)
Back	45 (22.6)
Side	21 (10.6)
Placed Position	
Stomach	50 (25.1)
Back	66 (33.2)
Side	26 (13.1)
Objects in Sleeping Area	
Mattress	126 (63.3)
Adults	98 (49.2)
Pillow	89 (44.7)
Comforter, quilt or other	81 (40.7)
Children	33 (16.6)
Toys	15 (7.5)
Bumper pads	10 (5.0)
Was This Death Found to be Preventable?	
Yes	158 (79.4)
No	41 (20.6)

The Louisiana Department of Health, Bureau of Family Health implements a multifaceted prevention effort, including community awareness campaigns for safe sleep, providing safe sleep resources to community partners, such as child care centers and all birthing hospitals. Training coroners and death scene investigators on proper protocol and procedures including doll reenactment and maternal interview has helped better correctly classify and understand SUID. At the local level, the Regional MCH Coordinators convene community action teams that work to build community capacity to prevent these deaths. SUID is a risk for all young babies and the Bureau of Family Health is working to minimize the risk and reduce infant deaths in Louisiana.

For more references or more information, email jane.herwehe@la.gov or go to http://dcfs.louisiana.gov/index.cfm?md=p_agebuilder&tmp=home&pid=357 or <http://www.cdc.gov/sids/aboutsuidandsids.htm>.

National Lead Poisoning Prevention Week

October 23-29, 2016

Trina Evans Williams, MPH

Lead poisoning is the most preventable environmental disease among young children, yet half a million US children have blood lead levels above 5 mcg/dl, the reference level at which Centers for Disease Control and Prevention (CDC), recommends public health actions be initiated. A simple blood test can prevent permanent damage that will last a lifetime.

Homes painted before 1978, when household lead paint was outlawed, are more likely to have lead, as are older water pipes and service lines.

Other actions include:

- getting the home tested if there are children with elevated blood lead levels, or whose parents suspect lead contamination;
- warn workers not to spread lead from job sites to the family

vehicle or home if there has been contact within church or community... especially with disaster volunteers;
 - share dangers of remodeling or refinishing old homes and furnishings with young women who are pregnant or have toddlers.

Resources on lead safety can be found at websites: [National Lead Poisoning Prevention Week](#) (CDC); [You Can Prevent Lead Poisoning](#) (LSU Ag Center); and [Louisiana Healthy Homes and Childhood Lead Poisoning Prevention Program's "Toolkit"](#) (LDH OPH).

For more information, please contact Trina Evans Williams at (504) 568-8254 or email Trina.Evans@la.gov.

Louisiana Fact

At the Forefront of Modern Dentistry

In the 1820's sales of teeth instruments to physicians and laymen were quite common. Although dentistry was beginning to emerge as a distinct profession, in the 1860's and 1870's most dental work was still performed by general practitioners or charlatans. A rural doctor could make and sell his own medicines as well as perform veterinary or dental services. Some medical account books in these years show that the extraction of one tooth would have a standard price of one dollar, the same price as administering medicine to a horse.

Dental education has a long history in Louisiana, dating as far back as 1861, with the establishment of the New Orleans Dental College. Classes in the dental college were suspended due to the Civil War but started again in 1867 and continued until 1877. This dental college accepted women on an equal basis, unlike the medical school which refused to accept them.



Photo: C. Edmund Kells, *The Dentist's Own Book*, 1925

C. Edmund Kells of New Orleans dentist, researcher, and inventor, who did pioneering work with x-rays, was the first in the U.S. to use dental xrays on a live patient in April, 1896.

He also was the first dentist to use compressed air in the dental office for drying the teeth and invented the oral suction pump. Surgeons around the world quickly adopted his suction pump for use in operating rooms, the idea of which is still in use today. Besides his

many innovations for the dental profession, Dr. Kells had among his 30 patents, the fire extinguisher, automotive jack, electric thermostat, automatic window closer, and elevator starter and brake.

He was a vocal proponent of keeping natural teeth, pulpless or not, which were regularly pulled by most of the dental community. Dr. Kells believed that the diagnosis of dental disease should be done by dentists, not physicians and followed the latest treatments of infection available.

In 1885, the first female dental assistant was employed by Dr. Kells to offer chair-side assistance, instrument cleaning, inventory, appointments, bookkeeping, and reception. He believed that the 'lady assistant' was absolutely essential to the modern dental office and would survive as long as dentistry lived.

- Thanks to the American Dental Association and Andrea Matlak, Archivist

- Ruldoph Matas, Edited by John Duffy, *The History of Medicine in Louisiana Vol II*, 29,352, 549-550

Submitted by Abhaykumar Unjiya, BDS, MSPH Candidate




Photo courtesy of Yahoo.com

Table: Communicable Disease Surveillance, Incidence by Region and Time Period, July-August, 2016

DISEASE	HEALTH REGION									TIME PERIOD					
	1	2	3	4	5	6	7	8	9	Jul-Aug 2016	Jul-Aug 2015	Jan-Dec Cum 2016	Jan-Dec Cum 2015	Jan-Dec % Chg*	
<u>Vaccine-preventable</u>															
Hepatitis B	Cases	0	3	0	0	2	0	1	0	3	9	12	35	55	-36.4
	Rate ¹	0	0.5	0	0	0.7	0	0.2	0	0.8	0.2	0.3	0.8	1.3	NA*
Measles		0	0	0	0	0	0	0	0	0	0	0	0	0	NA*
Mumps		0	0	0	0	0	0	0	0	0	0	0	0	0	NA*
Rubella		0	0	0	0	0	0	0	0	0	0	0	0	0	NA*
Pertussis		1	3	0	0	0	0	6	0	1	11	4	35	31	NA*
<u>Sexually-transmitted</u>															
HIV/AIDS	Cases ²	72	60	13	13	6	5	27	16	6	218	200	863	770	12.1
	Rate ¹	8.6	9.0	3.2	2.2	2.1	1.6	5.0	4.5	1.1	4.8	4.4	19.0	17.0	NA*
Chlamydia	Cases ^{1,3}	1,382	636	394	572	241	377	657	544	458	5,261	5,730	20,936	20,763	0.8
	Rate ¹	154.4	93.2	97.1	94.1	80.6	123.1	120.4	153.2	79.9	112.6	122.7	448.2	444.5	NA*
Gonorrhea	Cases ^{1,3}	553	270	130	222	81	112	228	205	133	1,934	1,935	7,401	6,608	12.0
	Rate ¹	61.8	39.6	32.0	36.5	27.1	36.6	41.8	57.7	23.2	41.4	41.4	158.5	141.5	NA*
Syphilis (P&S)	Cases ^{1,3}	33	5	16	13	8	7	26	2	5	115	177	428	457	-6.3
	Rate ¹	3.7	0.7	3.9	2.1	2.7	2.3	4.8	0.6	0.9	2.5	3.8	9.2	9.8	NA*
<u>Enteric</u>															
Campylobacter	Cases	1	12	5	1	2	2	5	9	9	46	50	168	159	5.7
Hepatitis A	Cases	0	0	0	0	0	0	0	0	0	0	1	9	3	200.0
	Rate ¹	0	0	0	0	0	0	0	0	0	0	0	0.2	0.1	NA*
Salmonella	Cases	26	35	28	60	19	27	18	27	46	286	342	808	820	-1.5
	Rate ¹	2.5	6.2	7.4	11.6	7.1	8.9	3.6	7.7	11.9	6.6	7.9	18.7	19.0	NA*
Shigella	Cases	8	5	9	6	1	5	6	12	4	56	42	247	123	100.8
	Rate ¹	0.8	0.9	2.4	1.2	0.4	1.6	1.2	3.4	1.0	1.3	1.0	5.7	2.9	NA*
Vibrio, cholera	Cases	0	0	0	0	0	0	0	0	0	0	0	0	0	NA*
Vibrio, other	Cases	5	1	3	2	0	0	0	0	3	14	8	35	40	-12.5
<u>Other</u>															
<i>H. influenzae (other)</i>		0	3	0	0	0	0	0	2	1	6	6	37	49	-24.5
<i>N. Meningitidis</i>		0	0	0	0	0	0	0	0	0	0	0	2	4	NA*

¹ = Cases Per 100 000 Population.

² = These totals reflect people with HIV infection whose status was first detected during the specified time period. This includes people who were diagnosed with AIDS at the time HIV first was detected. Because of delays in reporting HIV/AIDS cases, the number of persons reported is a minimal estimate. Data should be considered provisional.

³ = Preliminary data.

* = Percent change not calculated for rates or count differences less than 5.


Table 2. Diseases of Low Frequency, January-December, 2016

Disease	Total to Date
Legionellosis	20
Lyme Disease	4
Malaria	10
Rabies, animal	2
Varicella	46

Table 3. Animal Rabies, Jul-Aug, 2016

Parish	No. Cases	Species
Caddo	1	Bat

Figure: Department of Health Regional Map



Sanitary Code - State of Louisiana Part II - The Control of Disease

LAC 51:II.105: The following diseases/conditions are hereby declared reportable with reporting requirements by Class:

Class A Diseases/Conditions - Reporting Required Within 24 Hours

Diseases of major public health concern because of the severity of disease and potential for epidemic spread-report by telephone immediately upon recognition that a case, a suspected case, or a positive laboratory result is known; [in addition, all cases of rare or exotic communicable diseases, unexplained death, unusual cluster of disease and all outbreaks shall be reported.

Acute Flaccid Paralysis	Fish/Shellfish Poisoning (domoic acid, neurotoxic shellfish poisoning, ciguatera, paralytic shellfish poisoning, scombroid)	Plague (<i>Yersinia pestis</i>)	Smallpox
Anthrax	Foodborne Infection	Poliomyelitis (paralytic & non-paralytic)	<i>Staphylococcus aureus</i> , Vancomycin Intermediate or Resistant (VISA/VRSA)
Avian or Novel Strain Influenza A (initial detection)	<i>Haemophilus influenzae</i> (invasive infection)	Q Fever (<i>Coxiella burnetii</i>)	Staphylococcal Enterotoxin B (SEB) Pulmonary Poisoning
Botulism	Influenza-associated Mortality	Rabies (animal and human)	Tularemia (<i>Francisella tularensis</i>)
Brucellosis	Measles (Rubeola imported or indigenous)	Ricin Poisoning	Viral Hemorrhagic Fever (Ebola, Lassa, Marburg, Crimean Congo, etc.)
Cholera	Neisseria meningitidis (invasive infection)	Rubella (congenital syndrome)	Yellow Fever
<i>Clostridium perfringens</i> (foodborne infection)	Outbreaks of Any Infectious Disease	Rubella (German Measles)	
Diphtheria	Pertussis	Severe Acute Respiratory Syndrome-associated Coronavirus (SARS-CoV)	

Class B Diseases/Conditions - Reporting Required Within 1 Business Day

Diseases of public health concern needing timely response because of potential of epidemic spread-report by the end of the next business day after the existence of a case, a suspected case, or a positive laboratory result is known.

Amoeba (free living infection: <i>Acanthamoeba</i> , <i>Naegleria</i> , <i>Balamuthia</i> , others)	Chagas Disease	Hepatitis B (perinatal infection)	Mumps
Anaplasmosis	Chancroid	Hepatitis E	Salmonellosis
Arthropod-Borne Viral Infections (West Nile, Dengue, St. Louis, California, Eastern Equine, Western Equine, Chikungunya, Usutu, and others)	<i>Escherichia coli</i> , Shiga-toxin producing (STEC), including <i>E. coli</i> O157:H7	Herpes (neonatal)	Shigellosis
Aseptic Meningitis	Granuloma Inguinale	Human Immunodeficiency Virus ² [(HIV), infection in pregnancy]	Syphilis ¹
Babesiosis	Hantavirus (infection or Pulmonary Syndrome)	Human Immunodeficiency Virus ² [(HIV), perinatal exposure]	Tetanus
	Hemolytic-Uremic Syndrome	Legionellosis	Tuberculosis ³ (due to <i>M. tuberculosis</i> , <i>M. bovis</i> , or <i>M. africanum</i>)
	Hepatitis A (acute illness)	Malaria	Typhoid Fever
	Hepatitis B (acute illness and carriage in pregnancy)		

Class C Diseases/Conditions - Reporting Required Within 5 Business Days

Diseases of significant public health concern-report by the end of the workweek after the existence of a case, suspected case, or a positive laboratory result is known.

Acquired Immune Deficiency Syndrome ³ (AIDS)	Giardiasis	Listeriosis	Staphylococcal Toxic Shock Syndrome
<i>Anaplasma Phagocytophilum</i>	Glanders (<i>Burkholderia mallei</i>)	Lyme Disease	Streptococcal Disease, Group A (invasive disease)
Blastomycosis	Gonorrhea ¹ (genital, oral, ophthalmic, pelvic inflammatory disease, rectal)	Lymphogranuloma Venereum ¹	Streptococcal Disease, Group B (invasive disease)
Campylobacteriosis	Hansen's Disease (leprosy)	Melioidosis (<i>Burkholderia pseudomallei</i>)	Streptococcal Toxic Shock Syndrome
Chlamydial infection ¹	Hepatitis C (acute illness)	Meningitis, Eosinophilic (including those due to <i>Angiostrongylus</i> infection)	<i>Streptococcus pneumoniae</i> , invasive disease
Coccidioidomycosis	Histoplasmosis	Nipah Virus Infection	Transmissible Spongiform Encephalopathies (Creutzfeldt-Jacob Disease & variants)
Cryptococcosis (<i>C. neoformans</i> and <i>C. gattii</i>)	Human Immunodeficiency Virus ² (HIV) (infection other than as in Class B)	Non-gonococcal Urethritis	Trichinosis
Cryptosporidiosis	Human T Lymphocyte Virus (HTLV I and II infection)	Ophthalmia neonatorum	Varicella (chickenpox)
Cyclosporiasis	Leptospirosis	Psittacosis	<i>Vibrio</i> Infections (other than cholera)
Ehrlichiosis (human granulocytic, human monocytic, <i>E. chaffeensis</i> and <i>E. ewingii</i>)		Spotted Fevers [<i>Rickettsia</i> species including Rocky Mountain Spotted Fever (RMSF)]	Yersiniosis
<i>Enterococcus</i> , Vancomycin Resistant [(VRE), invasive disease]		<i>Staphylococcus aureus</i> (MRSA), invasive infection	

Class D Diseases/Conditions - Reporting Required Within 5 Business Days

Cancer	Heavy Metal (arsenic, cadmium, mercury) Exposure and/or Poisoning (all ages) ⁵	Phenylketonuria ⁴	Severe Traumatic Head Injury
Carbon Monoxide Exposure and/or Poisoning ⁵	Hemophilia ⁴	Pneumoconiosis (asbestosis, berylliosis, silicosis, byssinosis, etc.)	Severe Undernutrition (severe anemia, failure to thrive)
Complications of Abortion	Lead Exposure and/or Poisoning (all ages) ^{4,5}	Radiation Exposure, Over Normal Limits	Sickle Cell Disease ⁴ (newborns)
Congenital Hypothyroidism ⁴	Pesticide-Related Illness or Injury (all ages) ⁵	Reye's Syndrome	Spinal Cord Injury
Galactosemia ⁴			Sudden Infant Death Syndrome (SIDS)

Case reports not requiring special reporting instructions (see below) can be reported by mail or facsimile on Confidential Disease Report forms (2430), facsimile (504) 568-8290, telephone (504) 568-8313, or (800) 256-2748 for forms and instructions.

¹Report on STD-43 form. Report cases of syphilis with active lesions by telephone, within one business day, to (504) 568-8374.

²Report to the Louisiana HIV/AIDS Program: Visit www.hiv.dhh.louisiana.gov or call 504-568-7474 for regional contact information.

³Report on form TB 2431 (8/94). Mail form to TB Control Program, DHH-OPH, P.O. Box 60630, New Orleans, LA. 70160-0630 or fax both sides of the form to (504) 568-5016

⁴Report to the Louisiana Genetic Diseases Program and Louisiana Childhood Lead Poisoning Prevention Programs: www.genetics.dhh.louisiana.gov or facsimile (504) 568-8253, telephone (504) 568-8254, or (800) 242-3112

⁵Report to the Section of Environmental Epidemiology and Toxicology: www.seet.dhh.louisiana.gov or call (225) 342-7136 or (888) 293-7020

All **laboratory facilities** shall, in addition to reporting tests indicative of conditions found in §105, report positive or suggestive results for additional conditions of public health interest. The following findings shall be reported as detected by laboratory facilities: 1. adenoviruses; 2. coronaviruses; 3. enteroviruses; 4. hepatitis B (carriage other than in pregnancy); 5. hepatitis C (past or present infection); 6. human metapneumovirus; 7. parainfluenza viruses; 8. respiratory syncytial virus; and 9. rhinoviruses.