

SUPPLEMENTAL THIRD DOSE VS. BOOSTER SHOT

WHAT'S THE DIFFERENCE?

SUPPLEMENTAL THIRD DOSE

(28 DAYS AFTER YOUR SECOND DOSE OF A MRNA COVID-19 VACCINE)

Based on CDC recommendations, third vaccine doses are available now for people who are considered moderately or severely immunosuppressed. These would include those who:

- Receive active cancer treatment for tumors or cancers of the blood.
- Received an organ transplant and are taking medicine to suppress the immune system.
- Have either had a stem cell transplant within the last two years or are taking medicine to suppress the immune system.
- Were diagnosed with DiGeorge syndrome or Wiskott-Aldrich syndrome.
- Are diagnosed with HIV and have a high viral load or low CD4 count, or are not currently taking medication to treat HIV.
- Are taking drugs such as high-dose steroids or other medications that may cause severe suppression of the immune system.

BOOSTER

(6 MONTHS AFTER LAST PFIZER VACCINE)

Effective immediately, booster shots of the Pfizer-BioNTech COVID-19 vaccine are available to adults with high risk medical conditions for COVID, or are at high risk for getting COVID due to their occupation, per new guidance from the Centers for Disease Control and Prevention (CDC).

If you are 6 months out from your two-dose Pfizer series, you are now eligible to receive a booster shot if you are:

- Age 65 or older
- Age 18 or older and have one of many underlying medical conditions
- Age 18 or older and live or work in a setting that places you at risk of being exposed to COVID

QUESTIONS?

Talk to your healthcare provider or call our vaccine hotline: 1-855-453-0774

BRING BACK
LOUISIANA

LOUISIANA
DEPARTMENT OF
HEALTH