

First Quarter 2016

Vol. 14, No. 1



**Louisiana HIV, AIDS and Early Syphilis Surveillance
Quarterly Report
March 31, 2016**

**Louisiana Department of Health and Hospitals
Office of Public Health
STD/HIV Program (SHP)**

**1450 Poydras Street, Suite 2136
New Orleans, Louisiana 70112**

504.568.7474

haprequest@la.gov

<http://dhh.louisiana.gov/hiv>

www.hiv411.org

Louisiana Department of Health and Hospitals
Office of Public Health
STD/HIV Program

DeAnn Gruber, PhD, LCSW
Administrative Director

Chaquetta Johnson, DNP, MPH
Deputy Program Director

Debbie Wendell, PhD, MPH
Data Management/Analysis Manager

Jessica Fridge, MSPH
Surveillance Program Manager

Javone Davis, MPH
Regional Operations Manager

Stephanie Taylor, MD
STD Medical Director

Joy Ewell
CDC Lead PHA-Louisiana

Editor/Production:

Demerial Banks, MPH
Catherine Desmarais, MPH

Data Management/Analysis and Surveillance:

Demerial Banks, MPH, *HIV Surveillance Epidemiologist*

Tania Bechtel, *Incidence & Resistance Coordinator*

Antoine Brantley, MPH, *Data Analyst*

Lakeesha Brumfield, *HIV Data Entry Specialist*

Catherine Desmarais, DrPH, *STD Surveillance Epidemiologist and Data Manager*

Ashley Hoover, *STD/HIV Perinatal Surveillance Supervisor*

Mohammad Rahman, MBBS, MPH, PhD, *STD Epidemiologist Supervisor*

Denise Robertson, *ICCR Coordinator*

William T. Robinson, PhD., *Biostatistician*

Josette Russell, *HIV Data Entry Specialist*

Molly Votaw, MPH, *HIV Surveillance Data Management Supervisor*

Carmen Washington, *HIV Data Manager*

HIV Field Epidemiologists:

Betsy Shepard, PAC, MPH, *Epidemiologist Supervisor*

James Hubbard

Region 1

Cheryl Edney, BS MT(ASCP)

Regions 4, 5, 6

Monica Pendelton

Regions 1, 3

Gayle Norris, MA

Regions 7, 8

Gale Terry, LPN

Regions 2, 9

Regional STD Supervisors:

Michael Carter

Region 1

Patrick Williams

Region 5 & 6

Megan King

Region 2

Charles Jones

Region 7

Javone Davis

Region 8

John Lyons

Region 3 & 9

Ken May

Region 4

TABLE OF CONTENTS

| Contents | Page |
|--|-------------|
| Surveillance Summary | 1 |
| Louisiana: Persons Living with HIV Infection | 3 |
| Persons newly diagnosed with HIV and AIDS | 4 |
| Persons diagnosed with Early Syphilis | 5 |
| Parish: Persons Living with HIV Infection | 6 |
| Persons diagnosed with Early Syphilis | 7 |
| Metro New Orleans: Persons Living with HIV Infection | 8 |
| Persons newly diagnosed with HIV and AIDS | 9 |
| Persons diagnosed with Early Syphilis | 10 |
| Metro Baton Rouge: Persons Living with HIV Infection | 11 |
| Persons newly diagnosed with HIV and AIDS | 12 |
| Persons diagnosed with Early Syphilis | 13 |
| Region 1: Persons Living with HIV Infection | 14 |
| Persons newly diagnosed with HIV and AIDS | 15 |
| Persons diagnosed with Early Syphilis | 16 |
| Region 2: Persons Living with HIV Infection | 17 |
| Persons newly diagnosed with HIV and AIDS | 18 |
| Persons diagnosed with Early Syphilis | 19 |
| Region 3: Persons Living with HIV Infection | 20 |
| Persons newly diagnosed with HIV and AIDS | 21 |
| Persons diagnosed with Early Syphilis | 22 |
| Region 4: Persons Living with HIV Infection | 23 |
| Persons newly diagnosed with HIV and AIDS | 24 |
| Persons diagnosed with Early Syphilis | 25 |
| Region 5: Persons Living with HIV Infection | 26 |
| Persons newly diagnosed with HIV and AIDS | 27 |
| Persons diagnosed with Early Syphilis | 28 |
| Region 6: Persons Living with HIV Infection | 29 |
| Persons newly diagnosed with HIV and AIDS | 30 |
| Persons diagnosed with Early Syphilis | 31 |
| Region 7: Persons Living with HIV Infection | 32 |
| Persons newly diagnosed with HIV and AIDS | 33 |
| Persons diagnosed with Early Syphilis | 34 |
| Region 8: Persons Living with HIV Infection | 35 |
| Persons newly diagnosed with HIV and AIDS | 36 |
| Persons diagnosed with Early Syphilis | 37 |
| Region 9: Persons Living with HIV Infection | 38 |
| Persons newly diagnosed with HIV and AIDS | 39 |
| Persons diagnosed with Early Syphilis | 40 |
| List of Figures | |
| Louisiana: 10-Year Trends New HIV and New AIDS Diagnoses (2006-2015) | 41 |
| New HIV Diagnoses | |
| New AIDS Diagnoses | |
| Louisiana: 10-Year Persons Living with HIV Infection (2006-2015) | 42 |
| Louisiana: 7-Year Trends P&S Syphilis Diagnoses (2009-2015) | 42 |
| Louisiana: 7-Year Trends Early Syphilis Diagnoses (2009-2015) | 43 |
| Louisiana 2015 STD Update Fact Sheet | 44 |

Surveillance Summary

Data in this report includes cases diagnosed with HIV, AIDS or Early Syphilis

- In 2014, Louisiana ranked **2nd** in the nation for estimated HIV case rates (30.4 per 100,000 population) and **9th** in the estimated number of HIV cases. The Baton Rouge MSA ranked **1st** and the New Orleans MSA ranked **3rd** for estimated HIV case rates (44.7 and 36.9 per 100,000, respectively), among the large metropolitan areas in the nation. The most recent *CDC HIV Surveillance Report* (vol. 26) was released in November 2015 and provided the 2014 rankings.
- According to the same report, Louisiana ranked **2nd** highest in estimated state AIDS case rates (13.7 per 100,000) and **11th** in the number of estimated AIDS cases in 2014. The Baton Rouge metropolitan area ranked **1st** in estimated AIDS case rates (21.6 per 100,000) and the New Orleans metropolitan area ranked **4th** in estimated AIDS case rates (17.0 per 100,000) in 2014 among the large metropolitan areas in the nation.
- In 2014, Louisiana ranked **2nd** in the nation for primary and secondary (P&S) syphilis case rates (12.4 per 100,000 population) according to the 2014 *CDC STD Surveillance Report*. A total of 575 P&S syphilis cases were reported.
- As of March 31, 2016, a total of **20,912** persons are living with HIV infection (PLWH) in Louisiana; of these individuals, **10,962** persons (52%) have an AIDS diagnosis.
- In 2015, **1,207** new HIV cases were diagnosed in Louisiana. Of these cases, **33%** (n=394) occurred in Region 1 (New Orleans) and **22%** (n=268) occurred in Region 2 (Baton Rouge).
- In the first quarter of 2016, a total of **217** persons were diagnosed with early syphilis* in Louisiana; **132** persons were diagnosed with P&S syphilis, and **85** persons were diagnosed with early latent syphilis. Of the total early syphilis cases, **27%** (n=59) occurred in Region 1 (New Orleans) and **22%** (n=47) occurred in Region 7 (Shreveport).
- The primary risk factor for new HIV diagnoses in Louisiana is men who have sex with men (MSM). In 2015, of the persons who reported a risk factor, **68%** reported being MSM, and an additional **3%** reported being MSM and an injection drug user (IDU). As of March 31, 2016, **54%** of all people living with HIV infection in Louisiana were MSM.
- The primary risk factor for early syphilis diagnoses in Louisiana is heterosexual contact. In the first quarter of 2016, of the persons who reported a risk factor, **60%** reported engaging in heterosexual activity, and **44%** reported being MSM. *Persons could report multiple risk factors for syphilis diagnoses.*
- In Louisiana, **76%** of new HIV cases, **71%** of new AIDS cases, and **70%** of the early syphilis diagnoses were among men.
- African Americans in Louisiana are experiencing a severe health disparity in regards to HIV and STDs. In 2015, **71%** of newly diagnosed HIV cases and **73%** of newly diagnosed AIDS cases were among African Americans. In the first quarter of 2016, **75%** of early syphilis cases were among African Americans. African Americans make up only 32% of Louisiana's overall population.
- A large percentage of new HIV and syphilis diagnoses occurred among adolescents and young adults. In 2015, **25%** of all new HIV diagnoses occurred among persons 13-24 years old and an additional **33%** occurred among persons 25-34 years old. In the first quarter of 2016, **36%** of all early syphilis diagnoses occurred among persons 13-24 years old and an additional **37%** occurred among persons 25-34 years old.

* A syphilis diagnosis with less than one year duration is considered to be "early syphilis." This designation covers the primary, secondary, and early latent stages of syphilis.

New Quarterly Report Format:

SHP's new quarterly report format includes P&S syphilis and early latent syphilis diagnoses for Louisiana, the New Orleans and Baton Rouge MSAs, and all nine of the public health regions. It is important to note that the time periods for the HIV and syphilis data are not equivalent. Please pay special attention to the titles above each table. There are different reporting delays and methods for HIV and syphilis which impact the timeliness of those data in this report. The 'Persons Living with AIDS' excerpts have been removed.

AIDS Case Definition Change – Effective for all 2014 and future AIDS Diagnoses:

Prior to 2014, the surveillance case definition for an AIDS diagnosis was: a CD4 cell count <200, a CD4 percentage <14%, or the diagnosis of an opportunistic infection (OI) such as Kaposi Sarcoma. The CDC has changed the case surveillance definition for AIDS. For all AIDS diagnoses, starting January 1, 2015, the case definition is: a CD4 cell count <200 or the diagnosis of an OI. A CD4 percent <14% is no longer part of the AIDS case definition, unless the CD4 count is not available. Previously, a person with a CD4 count of 300 but a percent of 10%, would have been defined as an AIDS case. Because that CD4 count is not below 200, that case is no longer considered a new AIDS diagnosis. If the CD4 count of 300 was not available and only the percent of 10% was available, then the case would be newly diagnosed as AIDS. This change in surveillance case definition will reduce the number of AIDS diagnoses that occur each year, starting with 2014 data.

Time Periods for this Quarterly Report (First Quarter 2016 Report):

Persons Living with HIV Infection – As of March 31, 2016

New HIV Diagnoses – Calendar Year 2015

New AIDS Diagnoses – Calendar Year 2015

New P&S Syphilis Diagnoses – January to March 31, 2016

New Early Latent Syphilis Diagnoses – January to March 31, 2016

Future Quarterly Report:

Second Quarter 2016 Report: Release in August 2016

Persons Living with HIV Infection, As of June 30, 2016

HIV and AIDS Diagnoses, Calendar Year 2015 (Updated)

P&S Syphilis and Early Latent Syphilis Diagnoses, First and Second Quarter 2016 (January-June 2016)

Third Quarter 2016 Report: Release in November 2016

Persons Living with HIV Infection, As of September 30, 2016

HIV and AIDS Diagnoses, Calendar Year 2015 (Updated)

P&S Syphilis and Early Latent Syphilis Diagnoses, First-Third Quarter 2016 (January-September 2016)

Fourth Quarter 2016 Report: Release in February 2017

Persons Living with HIV Infection, As of December 31, 2016

HIV and AIDS Diagnoses, Calendar Year 2015 (Updated)

P&S Syphilis and Early Latent Syphilis Diagnoses, Calendar Year 2016

Persons Living with HIV Infection
Louisiana
as of March 31, 2016

- **36,829** cumulative cases have been detected in Louisiana, including **351** pediatric cases.
- **13,470** cumulative deaths among persons with HIV infection have occurred in Louisiana, including **91** pediatric deaths.
- **20,912** persons are currently living with HIV infection in Louisiana.
 Persons living with HIV: **9,986**
 Persons living with AIDS: **10,926**

Gender

| | Cases | Percent |
|-------------------|---------------|----------------|
| Women | 6,057 | 29 % |
| Men | 14,686 | 70 % |
| Transgender Women | 167 | 1 % |
| Transgender Men | 2 | <1 % |
| Total | 20,912 | 100 % |

Age Category

| Current Age | Cases | Percent |
|--------------------|---------------|----------------|
| 0 - 12 | 52 | <1 % |
| 13 - 19 | 115 | 1 % |
| 20 - 24 | 812 | 4 % |
| 25 - 29 | 1,847 | 9 % |
| 30 - 34 | 2,139 | 10 % |
| 35 - 39 | 2,499 | 12 % |
| 40 - 44 | 2,396 | 11 % |
| 45 - 49 | 2,664 | 13 % |
| 50 - 54 | 3,112 | 15 % |
| 55 - 59 | 2,520 | 12 % |
| 60+ | 2,756 | 13 % |
| Total | 20,912 | 100 % |

Race/Ethnicity by Birth Sex

| | Men | Women | Total | Percent |
|------------------------|---------------|--------------|---------------|----------------|
| Black, Non-Hispanic | 9,323 | 4,978 | 14,301 | 68 % |
| White, Non-Hispanic | 4,606 | 853 | 5,459 | 26 % |
| Hispanic | 709 | 166 | 875 | 4 % |
| Asian/Pacific Islander | 72 | 14 | 86 | <1 % |
| American Indian | 30 | 11 | 41 | <1 % |
| Unknown | 11 | 1 | 12 | <1 % |
| Multi-race | 102 | 36 | 138 | 1 % |
| Total | 14,853 | 6,059 | 20,912 | 100 % |

Exposure Category by Birth Sex

| Cases with a reported risk: | Men | Women | Total | Percent |
|--|---------------|--------------|---------------|----------------|
| Men who have sex with men | 7,644 | 0 | 7,644 | 54 % |
| Injection drug use | 1,034 | 641 | 1,675 | 12 % |
| Men who have sex with men and inject drugs | 858 | 0 | 858 | 6 % |
| Heterosexual contact | 1,147 | 2,452 | 3,599 | 26 % |
| Transfusion/Hemophilia | 40 | 30 | 70 | <1 % |
| Confirmed Other | 0 | 1 | 1 | <1 % |
| Perinatal | 96 | 112 | 208 | 1 % |
| Total | 10,819 | 3,236 | 14,055 | 100 % |
| <i>No Identified Risk</i> | <i>4,034</i> | <i>2,823</i> | <i>6,857</i> | |

**Persons Newly Diagnosed with HIV and AIDS
Louisiana**

HIV CASES DIAGNOSED January - December 2015

• **1,207** persons were newly diagnosed with HIV in Louisiana.

| Gender | Cases | Percent | Race/Ethnicity | Cases | Percent |
|-------------------|--------------|----------------|------------------------|--------------|----------------|
| Men | 914 | 76 % | Black, Non-Hispanic | 853 | 70 % |
| Women | 287 | 24 % | White, Non-Hispanic | 256 | 22 % |
| Transgender Women | 6 | <1 % | Hispanic | 74 | 6 % |
| Total | 1,207 | 100 % | Asian/Pacific Islander | 4 | 1 % |
| | | | American Indian | 10 | <1 % |
| | | | Unknown | 5 | <1 % |
| | | | Multi-race | 5 | 1 % |
| | | | Total | 1,207 | 100 % |

| Age at Diagnosis | Cases | Percent | Cases with a reported risk (by birth sex): | Cases | Percent |
|-------------------------|--------------|----------------|---|--------------|----------------|
| 0 - 12 | 2 | <1 % | Men who have sex with men | 422 | 68 % |
| 13 - 24 | 298 | 25 % | Injection drug use | 42 | 7 % |
| 25 - 34 | 395 | 33 % | Men who have sex with men and inject drugs | 21 | 3 % |
| 35 - 44 | 230 | 19 % | Heterosexual contact | 129 | 21 % |
| 45+ | 282 | 23 % | Perinatal | 3 | <1 % |
| Total | 1,207 | 100 % | Total | 617 | 100 % |
| | | | <i>No Identified Risk</i> | <i>590</i> | |

AIDS CASES DIAGNOSED January - December 2015

• **538** persons were newly diagnosed with AIDS in Louisiana.

| Gender | Cases | Percent | Race/Ethnicity | Cases | Percent |
|-------------------|--------------|----------------|------------------------|--------------|----------------|
| Men | 381 | 71 % | Black, Non-Hispanic | 393 | 74 % |
| Women | 154 | 29 % | White, Non-Hispanic | 110 | 19 % |
| Transgender Women | 3 | <1 % | Hispanic | 23 | 5 % |
| Total | 538 | 100 % | Asian/Pacific Islander | 2 | <1 % |
| | | | American Indian | 6 | <1 % |
| | | | Multi-race | 4 | 1 % |
| | | | Total | 538 | 100 % |

| Age at Diagnosis | Cases | Percent | Cases with a reported risk (by birth sex): | Cases | Percent |
|-------------------------|--------------|----------------|---|--------------|----------------|
| 0 - 12 | 1 | <1 % | Men who have sex with men | 160 | 55 % |
| 13 - 24 | 47 | 9 % | Injection drug use | 27 | 9 % |
| 25 - 34 | 152 | 28 % | Men who have sex with men and inject drugs | 10 | 3 % |
| 35 - 44 | 135 | 25 % | Heterosexual contact | 92 | 32 % |
| 45+ | 203 | 38 % | Perinatal | 2 | 1 % |
| Total | 538 | 100 % | Total | 291 | 100 % |
| | | | <i>No Identified Risk</i> | <i>247</i> | |

**Persons Diagnosed with Early Syphilis
Louisiana**

PERSONS DIAGNOSED WITH P&S SYPHILIS January – March 2016

- 132 persons were diagnosed with P&S Syphilis in Louisiana between January 1 and March 31, 2016.

| <u>Gender</u> | <u>Cases</u> | <u>Percent</u> | <u>Race/Ethnicity</u> | <u>Cases</u> | <u>Percent</u> |
|---------------|--------------|----------------|------------------------|--------------|----------------|
| Men | 96 | 73 % | American Indian | 0 | 0 % |
| Women | 36 | 27 % | Asian/Pacific Islander | 0 | 0 % |
| Total | 132 | 100 % | Black, Non-Hispanic | 100 | 76 % |
| | | | Hispanic | 4 | 3 % |
| | | | White, Non-Hispanic | 28 | 21 % |
| | | | Other/Multi-race | 0 | 0 % |
| | | | Unknown | 0 | 0 % |
| | | | Total | 132 | 100 % |

| <u>Age at Diagnosis</u> | <u>Cases</u> | <u>Percent</u> | <u>Cases with a reported risk (Can have >1 risk)</u> | <u>Cases</u> | <u>Percent</u> |
|-------------------------|--------------|----------------|---|--------------|----------------|
| 0 - 12 | 0 | 0 % | Men who have sex with men and inject drugs | 0 | 0 % |
| 13 - 19 | 13 | 10 % | Men who have sex with men | 51 | 43 % |
| 20 – 29 | 64 | 48 % | Injection drug use | 3 | 3 % |
| 30 - 39 | 35 | 27 % | Transactional sex | 6 | 5 % |
| 40+ | 20 | 15 % | High risk heterosexual contact | 22 | 18 % |
| Total | 132 | 100 % | Heterosexual contact | 53 | 45 % |
| | | | Incarceration | 12 | 10 % |
| | | | Illegal drug use (non-injection) | 10 | 8 % |
| | | | <i>No Reported Risk</i> | <i>13</i> | |

PERSONS DIAGNOSED WITH EARLY LATENT SYPHILIS January - March 2016

- 85 persons were diagnosed with Early Latent Syphilis in Louisiana between January 1 and March 31, 2016.

| <u>Gender</u> | <u>Cases</u> | <u>Percent</u> | <u>Race/Ethnicity</u> | <u>Cases</u> | <u>Percent</u> |
|---------------|--------------|----------------|------------------------|--------------|----------------|
| Men | 56 | 66 % | American Indian | 0 | 0 % |
| Women | 29 | 34 % | Asian/Pacific Islander | 1 | 1 % |
| Total | 85 | 100 % | Black, Non-Hispanic | 63 | 74 % |
| | | | Hispanic | 1 | 1 % |
| | | | White, Non-Hispanic | 20 | 24 % |
| | | | Other/Multi-race | 0 | 0 % |
| | | | Unknown | 0 | 0 % |
| | | | Total | 85 | 100 % |

| <u>Age at Diagnosis</u> | <u>Cases</u> | <u>Percent</u> | <u>Cases with a reported risk (Can have >1 risk)</u> | <u>Cases</u> | <u>Percent</u> |
|-------------------------|--------------|----------------|---|--------------|----------------|
| 0 - 12 | 0 | 0 % | Men who have sex with men and inject drugs | 1 | 1 % |
| 13 - 19 | 9 | 11 % | Men who have sex with men | 37 | 45 % |
| 20 – 29 | 37 | 44 % | Injection drug use | 1 | 1 % |
| 30 - 39 | 26 | 31 % | Transactional sex | 3 | 4 % |
| 40+ | 13 | 15 % | High risk heterosexual contact | 10 | 12 % |
| Total | 85 | 100 % | Heterosexual contact | 36 | 43 % |
| | | | Incarceration | 16 | 19 % |
| | | | Illegal drug use (non-injection) | 9 | 11 % |
| | | | <i>No Reported Risk</i> | <i>2</i> | |

* A syphilis diagnosis with less than one year duration is considered to be “early syphilis.” This designation covers the primary, secondary, and early latent stages of syphilis.

Persons Living with HIV Infection by Parish
as of March 31, 2016

First Quarter 2016

| <u>Parish</u> | <u>Living Cases</u> | <u>Parish</u> | <u>Living Cases</u> |
|----------------------|----------------------------|----------------------|----------------------------|
| Acadia | 119 | Sabine | 18 |
| Allen | 262 | St. Bernard | 177 |
| Ascension | 241 | St. Charles | 118 |
| Assumption | 33 | St. Helena | 20 |
| Avoyelles | 162 | St. James | 73 |
| Beauregard | 48 | St. John the Baptist | 169 |
| Bienville | 30 | St. Landry | 297 |
| Bossier | 223 | St. Martin | 112 |
| Caddo | 1,170 | St. Mary | 95 |
| Calcasieu | 708 | St. Tammany | 473 |
| Caldwell | 24 | Tangipahoa | 395 |
| Cameron | 3 | Tensas | 37 |
| Catahoula | 26 | Terrebonne | 211 |
| Claiborne | 73 | Union | 31 |
| Concordia | 49 | Vermilion | 93 |
| De Soto | 54 | Vernon | 73 |
| East Baton Rouge | 3,991 | Washington | 187 |
| East Carroll | 25 | Webster | 89 |
| East Feliciana | 163 | West Baton Rouge | 102 |
| Evangeline | 70 | West Carroll | 13 |
| Franklin | 52 | West Feliciana | 150 |
| Grant | 53 | Winn | 70 |
| Iberia | 124 | | |
| Iberville | 313 | | |
| Jackson | 20 | | |
| Jefferson | 1,921 | | |
| Jefferson Davis | 63 | | |
| La Salle | 32 | | |
| Lafayette | 764 | | |
| Lafourche | 159 | | |
| Lincoln | 102 | | |
| Livingston | 227 | | |
| Madison | 37 | | |
| Morehouse | 64 | | |
| Natchitoches | 121 | | |
| Orleans | 5,098 | | |
| Ouachita | 635 | | |
| Plaquemines | 42 | | |
| Pointe Coupee | 51 | | |
| Rapides | 493 | | |
| Red River | 15 | | |
| Richland | 49 | | |
| | | Total | 20,912 |

**Persons Diagnosed with Early Syphilis* by Parish
January - March 2016**

| <u>Parish</u> | <u>Diagnoses</u> | <u>Parish</u> | <u>Diagnoses</u> |
|------------------|------------------|----------------------|------------------|
| Acadia | 1 | Sabine | 0 |
| Allen | 0 | St. Bernard | 0 |
| Ascension | 1 | St. Charles | 1 |
| Assumption | 0 | St. Helena | 0 |
| Avoyelles | 0 | St. James | 2 |
| Beauregard | 0 | St. John the Baptist | 3 |
| Bienville | 1 | St. Landry | 6 |
| Bossier | 5 | St. Martin | 3 |
| Caddo | 37 | St. Mary | 1 |
| Calcasieu | 7 | St. Tammany | 3 |
| Caldwell | 0 | Tangipahoa | 1 |
| Cameron | 0 | Tensas | 0 |
| Catahoula | 0 | Terrebonne | 3 |
| Claiborne | 0 | Union | 0 |
| Concordia | 1 | Vermilion | 1 |
| De Soto | 0 | Vernon | 0 |
| East Baton Rouge | 26 | Washington | 1 |
| East Carroll | 0 | Webster | 0 |
| East Feliciana | 2 | West Baton Rouge | 2 |
| Evangeline | 1 | West Carroll | 0 |
| Franklin | 0 | West Feliciana | 0 |
| Grant | 2 | Winn | 0 |
| Iberia | 2 | | |
| Iberville | 0 | | |
| Jackson | 1 | | |
| Jefferson | 16 | | |
| Jefferson Davis | 0 | | |
| La Salle | 0 | | |
| Lafayette | 13 | | |
| Lafourche | 3 | | |
| Lincoln | 1 | | |
| Livingston | 1 | | |
| Madison | 1 | | |
| Morehouse | 0 | | |
| Natchitoches | 3 | | |
| Orleans | 43 | | |
| Ouachita | 9 | | |
| Plaquemines | 0 | | |
| Pointe Coupee | 0 | | |
| Rapides | 12 | | |
| Red River | 1 | | |
| Richland | 0 | | |
| | | Total | 217 |

* A syphilis diagnosis with less than one year duration is considered to be "early syphilis." This designation covers the primary, secondary, and early latent stages of syphilis.

**Persons Living with HIV Infection
Metro New Orleans
as of March 31, 2016**

Jefferson, Orleans, St. Tammany, St. John the Baptist, Plaquemines, St. Bernard, St. James, St. Charles Parishes

- **16,715** cumulative cases have been detected in the New Orleans MSA, including **168** pediatric cases.
- **5,780** cumulative deaths among persons with HIV infection have occurred in the New Orleans MSA, including **45** pediatric deaths
- **8,071** persons are currently living with HIV infection in the New Orleans MSA.
Persons living with HIV: **3,837**
Persons living with AIDS: **4,234**

Gender

| | Cases | Percent |
|-------------------|--------------|----------------|
| Women | 2,032 | 25 % |
| Men | 5,943 | 74 % |
| Transgender Women | 94 | 1% |
| Transgender Men | 2 | <1% |
| Total | 8,071 | 100 % |

Age Category

| Current Age | Cases | Percent |
|--------------------|--------------|----------------|
| 0 - 12 | 21 | <1 % |
| 13 - 19 | 46 | 1 % |
| 20 - 24 | 276 | 3 % |
| 25 - 29 | 682 | 8 % |
| 30 - 34 | 810 | 10 % |
| 35 - 39 | 909 | 11 % |
| 40 - 44 | 881 | 11 % |
| 45 - 49 | 1,001 | 12 % |
| 50 - 54 | 1,238 | 15 % |
| 55 - 59 | 1,028 | 13 % |
| 60+ | 1,179 | 15 % |
| Total | 8,071 | 100 % |

Race/Ethnicity by Birth Sex

| | Men | Women | Total | Percent |
|------------------------|--------------|--------------|--------------|----------------|
| Black, Non-Hispanic | 3381 | 1645 | 5026 | 62 % |
| White, Non-Hispanic | 2176 | 259 | 2435 | 30 % |
| Hispanic | 388 | 114 | 502 | 6 % |
| Asian/Pacific Islander | 42 | 6 | 48 | 1 % |
| American Indian | 8 | 0 | 8 | <1 % |
| Unknown | 8 | 1 | 9 | <1 % |
| Multi-race | 34 | 9 | 43 | 1 % |
| Total | 6,037 | 2,034 | 8,071 | 100 % |

Exposure Category by Birth Sex

| Cases with a reported risk: | Men | Women | Total | Percent |
|--|--------------|--------------|--------------|----------------|
| Men who have sex with men | 3,457 | 0 | 3,457 | 61 % |
| Injection drug use | 321 | 216 | 537 | 10 % |
| Men who have sex with men and inject drugs | 334 | 0 | 334 | 6 % |
| Heterosexual contact | 376 | 826 | 1,202 | 21 % |
| Transfusion/Hemophilia | 18 | 13 | 31 | 1 % |
| Perinatal | 35 | 45 | 80 | 1 % |
| Total | 4,541 | 1,100 | 5,641 | 100 % |
| <i>No Identified Risk</i> | <i>1,496</i> | <i>934</i> | <i>2,430</i> | |

**Persons Newly Diagnosed with HIV and AIDS
Metro New Orleans**

Jefferson, Orleans, St. Tammany, St. John the Baptist, Plaquemines, St. Bernard, St. James, St. Charles Parishes

HIV CASES DIAGNOSED January - December 2015

- 433 persons were newly diagnosed with HIV in the New Orleans MSA.

| <u>Gender</u> | <u>Cases</u> | <u>Percent</u> | <u>Race/Ethnicity</u> | <u>Cases</u> | <u>Percent</u> |
|-------------------|--------------|----------------|------------------------|--------------|----------------|
| Men | 349 | 81 % | Black, Non-Hispanic | 289 | 67 % |
| Women | 79 | 18 % | White, Non-Hispanic | 97 | 22 % |
| Transgender Women | 5 | 1 % | Hispanic | 38 | 9 % |
| Total | 433 | 100 % | Asian/Pacific Islander | 2 | <1 % |
| | | | American Indian | 1 | <1 % |
| | | | Unknown | 4 | 1 % |
| | | | Multi-race | 2 | <1 % |
| | | | Total | 433 | 100 % |

| <u>Age at Diagnosis</u> | <u>Cases</u> | <u>Percent</u> | <u>Cases with a reported risk (by birth sex):</u> | <u>Cases</u> | <u>Percent</u> |
|-------------------------|--------------|----------------|---|--------------|----------------|
| 0 - 12 | 2 | <1 % | Men who have sex with men | 175 | 67 % |
| 13 - 24 | 101 | 23 % | Injection drug use | 24 | 9 % |
| 25 - 34 | 152 | 35 % | Men who have sex with men and inject drugs | 14 | 5 % |
| 35 - 44 | 88 | 20 % | Heterosexual contact | 46 | 18 % |
| 45+ | 90 | 21 % | Perinatal | 2 | 1 % |
| Total | 433 | 100 % | Total | 261 | 100 % |
| | | | <i>No Identified Risk</i> | <i>172</i> | |

AIDS CASES DIAGNOSED January - December 2015

- 191 persons were newly diagnosed with AIDS in the New Orleans MSA.

| <u>Gender</u> | <u>Cases</u> | <u>Percent</u> | <u>Race/Ethnicity</u> | <u>Cases</u> | <u>Percent</u> |
|-------------------|--------------|----------------|------------------------|--------------|----------------|
| Men | 139 | 73 % | Black, Non-Hispanic | 134 | 70 % |
| Women | 51 | 27 % | White, Non-Hispanic | 42 | 22 % |
| Transgender Women | 1 | <1 % | Hispanic | 12 | 6 % |
| Total | 191 | 100 % | Asian/Pacific Islander | 1 | 1 % |
| | | | American Indian | 1 | 1 % |
| | | | Multi-race | 1 | 1 % |
| | | | Total | 191 | 100 % |

| <u>Age at Diagnosis</u> | <u>Cases</u> | <u>Percent</u> | <u>Cases with a reported risk (by birth sex):</u> | <u>Cases</u> | <u>Percent</u> |
|-------------------------|--------------|----------------|---|--------------|----------------|
| 0 - 12 | 1 | 1 % | Men who have sex with men | 62 | 51 % |
| 13 - 24 | 16 | 8 % | Injection drug use | 15 | 12 % |
| 25 - 34 | 50 | 26 % | Men who have sex with men and inject drugs | 5 | 4 % |
| 35 - 44 | 50 | 26 % | Heterosexual contact | 38 | 31 % |
| 45+ | 74 | 39 % | Perinatal | 2 | 2 % |
| Total | 191 | 100 % | Total | 122 | 100 % |
| | | | <i>No Identified Risk</i> | <i>69</i> | |

**Persons Diagnosed with Early Syphilis
Metro New Orleans**

Jefferson, Orleans, St. Tammany, St. John the Baptist, Plaquemines, St. Bernard, St. James, St. Charles Parishes

PERSONS DIAGNOSED WITH P&S SYPHILIS January - March 2016

- 29 persons were diagnosed with P&S Syphilis in the New Orleans MSA from January 1 through March 31, 2016.

| <u>Gender</u> | <u>Cases</u> | <u>Percent</u> | <u>Race/Ethnicity</u> | <u>Cases</u> | <u>Percent</u> |
|---------------|--------------|----------------|------------------------|--------------|----------------|
| Men | 25 | 86 % | American Indian | 0 | 0 % |
| Women | 4 | 14 % | Asian/Pacific Islander | 0 | 0 % |
| Total | 29 | 100 % | Black, Non-Hispanic | 19 | 66 % |
| | | | Hispanic | 0 | 0 % |
| | | | White, Non-Hispanic | 10 | 34 % |
| | | | Other/Multi-race | 0 | 0 % |
| | | | Unknown | 0 | 0 % |
| | | | Total | 29 | 100 % |

| <u>Age at Diagnosis</u> | <u>Cases</u> | <u>Percent</u> | <u>Cases with a reported risk (Can have >1 risk)</u> | <u>Cases</u> | <u>Percent</u> |
|-------------------------|--------------|----------------|---|--------------|----------------|
| 0 - 12 | 0 | 0 % | Men who have sex with men and inject drugs | 0 | 0 % |
| 13 - 19 | 2 | 7 % | Men who have sex with men | 16 | 70 % |
| 20 - 29 | 15 | 52 % | Injection drug use | 0 | 0 % |
| 30 - 39 | 8 | 28 % | Transactional sex | 0 | 0 % |
| 40+ | 4 | 14 % | High risk heterosexual contact | 1 | 4 % |
| Total | 29 | 100 % | Heterosexual contact | 8 | 35 % |
| | | | Incarceration | 3 | 13 % |
| | | | Illegal drug use (non-injection) | 1 | 4 % |
| | | | <i>No Reported Risk</i> | 6 | |

PERSONS DIAGNOSED WITH EARLY LATENT SYPHILIS January - March 2016

- 39 persons were diagnosed with Early Latent Syphilis in the New Orleans MSA from January 1 through March 31, 2016.

| <u>Gender</u> | <u>Cases</u> | <u>Percent</u> | <u>Race/Ethnicity</u> | <u>Cases</u> | <u>Percent</u> |
|---------------|--------------|----------------|------------------------|--------------|----------------|
| Men | 31 | 79 % | American Indian | 0 | 0 % |
| Women | 8 | 21 % | Asian/Pacific Islander | 1 | 3 % |
| Total | 39 | 100 % | Black, Non-Hispanic | 30 | 77 % |
| | | | Hispanic | 0 | 0 % |
| | | | White, Non-Hispanic | 8 | 21 % |
| | | | Other/Multi-race | 0 | 0 % |
| | | | Unknown | 0 | 0 % |
| | | | Total | 39 | 100 % |

| <u>Age at Diagnosis</u> | <u>Cases</u> | <u>Percent</u> | <u>Cases with a reported risk (Can have >1 risk)</u> | <u>Cases</u> | <u>Percent</u> |
|-------------------------|--------------|----------------|---|--------------|----------------|
| 0 - 12 | 0 | 0 % | Men who have sex with men and inject drugs | 0 | 0 % |
| 13 - 19 | 0 | 0 % | Men who have sex with men | 24 | 65 % |
| 20 - 29 | 18 | 46 % | Injection drug use | 0 | 0 % |
| 30 - 39 | 12 | 31 % | Transactional sex | 0 | 0 % |
| 40+ | 9 | 23 % | High risk heterosexual contact | 4 | 11 % |
| Total | 39 | 100 % | Heterosexual contact | 9 | 24 % |
| | | | Incarceration | 7 | 19 % |
| | | | Illegal drug use (non-injection) | 1 | 3 % |
| | | | <i>No Reported Risk</i> | 2 | |

**Persons Living with HIV Infection
Metro Baton Rouge
as of March 31, 2016**

Ascension, East Baton Rouge, East Feliciana, Iberville, Livingston, Pointe Coupee, St. Helena, West Baton Rouge, West Feliciana Parishes

- **8,515** cumulative cases have been detected in the Baton Rouge MSA, including **80** pediatric cases.
- **3,432** cumulative deaths among persons with HIV infection have occurred in the Baton Rouge MSA, including **17** pediatric deaths.
- **5,258** persons are currently living with HIV infection in the Baton Rouge MSA.
Persons living with HIV: **2,484**
Persons living with AIDS: **2,774**

Gender

| | Cases | Percent |
|-------------------|--------------|----------------|
| Women | 1,834 | 35 % |
| Men | 3,381 | 64 % |
| Transgender Women | 43 | 1% |
| Total | 5,258 | 100 % |

Age Category

| Current Age | Cases | Percent |
|--------------------|--------------|----------------|
| 0 - 12 | 19 | <1 % |
| 13 - 19 | 24 | <1 % |
| 20 - 24 | 205 | 4 % |
| 25 - 29 | 471 | 9 % |
| 30 - 34 | 572 | 11 % |
| 35 - 39 | 685 | 13 % |
| 40 - 44 | 661 | 13 % |
| 45 - 49 | 665 | 13 % |
| 50 - 54 | 726 | 14 % |
| 55 - 59 | 580 | 11 % |
| 60+ | 650 | 12 % |
| Total | 5,258 | 100 % |

Race/Ethnicity by Birth Sex

| | Men | Women | Total | Percent |
|------------------------|--------------|--------------|--------------|----------------|
| Black, Non-Hispanic | 2630 | 1676 | 4306 | 82 % |
| White, Non-Hispanic | 654 | 168 | 822 | 16 % |
| Hispanic | 64 | 19 | 83 | 2 % |
| Asian/Pacific Islander | 11 | 5 | 16 | <1 % |
| American Indian | 2 | 0 | 2 | <1 % |
| Multi-race | 20 | 9 | 29 | 1 % |
| Total | 3,381 | 1,877 | 5,258 | 100 % |

Exposure Category by Birth Sex

| Cases with a reported risk: | Men | Women | Total | Percent |
|--|--------------|--------------|--------------|----------------|
| Men who have sex with men | 1,432 | 0 | 1,432 | 44 % |
| Injection drug use | 355 | 211 | 566 | 17 % |
| Men who have sex with men and inject drugs | 189 | 0 | 189 | 6 % |
| Heterosexual contact | 320 | 684 | 1,004 | 31 % |
| Transfusion/Hemophilia | 6 | 8 | 14 | <1 % |
| Perinatal | 31 | 27 | 58 | 2 % |
| Total | 2,333 | 930 | 3,263 | 100 % |
| <i>No Identified Risk</i> | <i>1,048</i> | <i>947</i> | <i>1,995</i> | |

Persons Newly Diagnosed with HIV and AIDS**Metro Baton Rouge***Ascension, East Baton Rouge, East Feliciana, Iberville, Livingston, Pointe Coupee, St. Helena, West Baton Rouge, West Feliciana Parishes***HIV CASES DIAGNOSED January - December 2015**

- 285 persons were newly diagnosed with HIV in the Baton Rouge MSA.

| <u>Gender</u> | <u>Cases</u> | <u>Percent</u> | <u>Race/Ethnicity</u> | <u>Cases</u> | <u>Percent</u> |
|-------------------|--------------|----------------|------------------------|--------------|----------------|
| Men | 198 | 69 % | Black, Non-Hispanic | 241 | 85 % |
| Women | 86 | 30 % | White, Non-Hispanic | 33 | 12 % |
| Transgender Women | 1 | <1 % | Hispanic | 8 | 3 % |
| Total | 285 | 100 % | Asian/Pacific Islander | 1 | <1 % |
| | | | Multi-race | 2 | 1 % |
| | | | Total | 285 | 100 % |

| <u>Age at Diagnosis</u> | <u>Cases</u> | <u>Percent</u> | <u>Cases with a reported risk (by birth sex):</u> | <u>Cases</u> | <u>Percent</u> |
|-------------------------|--------------|----------------|---|--------------|----------------|
| 13 - 24 | 75 | 26 % | Men who have sex with men | 99 | 72 % |
| 25 - 34 | 91 | 32 % | Injection drug use | 5 | 4 % |
| 35 - 44 | 59 | 21 % | Men who have sex with men and inject drugs | 4 | 3 % |
| 45+ | 60 | 21 % | Heterosexual contact | 28 | 20 % |
| Total | 285 | 100 % | Perinatal | 1 | 1 % |
| | | | Total | 137 | 100 % |

*No Identified Risk**148***AIDS CASES DIAGNOSED January - December 2015**

- 138 persons were newly diagnosed with AIDS in the Baton Rouge MSA.

| <u>Gender</u> | <u>Cases</u> | <u>Percent</u> | <u>Race/Ethnicity</u> | <u>Cases</u> | <u>Percent</u> |
|-------------------|--------------|----------------|------------------------|--------------|----------------|
| Men | 96 | 70 % | Black, Non-Hispanic | 112 | 89 % |
| Women | 41 | 30 % | White, Non-Hispanic | 23 | 10 % |
| Transgender Women | 1 | <1 % | Hispanic | 1 | 1 % |
| Total | 138 | 100 % | Asian/Pacific Islander | 1 | 1 % |
| | | | Multi-race | 1 | 1 % |
| | | | Total | 138 | 100 % |

| <u>Age at Diagnosis</u> | <u>Cases</u> | <u>Percent</u> | <u>Cases with a reported risk (by birth sex):</u> | <u>Cases</u> | <u>Percent</u> |
|-------------------------|--------------|----------------|---|--------------|----------------|
| 13 - 24 | 10 | 7 % | Men who have sex with men | 39 | 59 % |
| 25 - 34 | 39 | 28 % | Injection drug use | 7 | 11 % |
| 35 - 44 | 38 | 28 % | Men who have sex with men and inject drugs | 4 | 6 % |
| 45+ | 51 | 37 % | Heterosexual contact | 16 | 24 % |
| Total | 138 | 100 % | Total | 66 | 100 % |

*No Identified Risk**72*

**Persons Diagnosed with Early Syphilis
Metro Baton Rouge**

Ascension, East Baton Rouge, East Feliciana, Iberville, Livingston, Pointe Coupee, St. Helena, West Baton Rouge, West Feliciana Parishes

PERSONS DIAGNOSED WITH P&S SYPHILIS January – March 2015

- 26 persons were diagnosed with P&S Syphilis in the Baton Rouge MSA from January 1 through March 31, 2016.

| <u>Gender</u> | <u>Cases</u> | <u>Percent</u> | <u>Race/Ethnicity</u> | <u>Cases</u> | <u>Percent</u> |
|---------------|--------------|----------------|------------------------|--------------|----------------|
| Men | 20 | 77 % | American Indian | 0 | 0 % |
| Women | 6 | 23 % | Asian/Pacific Islander | 0 | 0 % |
| Total | 26 | 100 % | Black, Non-Hispanic | 21 | 81 % |
| | | | Hispanic | 0 | 0 % |
| | | | White, Non-Hispanic | 5 | 19 % |
| | | | Other/Multi-race | 0 | 0 % |
| | | | Unknown | 0 | 0 % |
| | | | Total | 26 | 100 % |

| <u>Age at Diagnosis</u> | <u>Cases</u> | <u>Percent</u> | <u>Cases with a reported risk (Can have >1 risk)</u> | <u>Cases</u> | <u>Percent</u> |
|-------------------------|--------------|----------------|---|--------------|----------------|
| 0 - 12 | 0 | 0 % | Men who have sex with men and inject drugs | 0 | 0 % |
| 13 - 19 | 3 | 12 % | Men who have sex with men | 15 | 60 % |
| 20 – 29 | 11 | 42 % | Injection drug use | 0 | 0 % |
| 30 - 39 | 5 | 19 % | Transactional sex | 1 | 4 % |
| 40+ | 7 | 27 % | High risk heterosexual contact | 2 | 8 % |
| Total | 26 | 100 % | Heterosexual contact | 8 | 32 % |
| | | | Incarceration | 0 | 0 % |
| | | | Illegal drug use (non-injection) | 0 | 0 % |
| | | | <i>No Reported Risk</i> | <i>1</i> | |

PERSONS DIAGNOSED WITH EARLY LATENT SYPHILIS January – March 2015

- 6 persons were diagnosed with Early Latent Syphilis in the Baton Rouge MSA from January 1 through March 31, 2016.

| <u>Gender</u> | <u>Cases</u> | <u>Percent</u> | <u>Race/Ethnicity</u> | <u>Cases</u> | <u>Percent</u> |
|---------------|--------------|----------------|------------------------|--------------|----------------|
| Men | 6 | 100 % | American Indian | 0 | 0 % |
| Women | 0 | 0 % | Asian/Pacific Islander | 0 | 0 % |
| Total | 6 | 100 % | Black, Non-Hispanic | 4 | 67 % |
| | | | Hispanic | 1 | 17 % |
| | | | White, Non-Hispanic | 1 | 17 % |
| | | | Other/Multi-race | 0 | 0 % |
| | | | Unknown | 0 | 0 % |
| | | | Total | 6 | 100 % |

| <u>Age at Diagnosis</u> | <u>Cases</u> | <u>Percent</u> | <u>Cases with a reported risk (Can have >1 risk)</u> | <u>Cases</u> | <u>Percent</u> |
|-------------------------|--------------|----------------|---|--------------|----------------|
| 0 - 12 | 0 | 0 % | Men who have sex with men and inject drugs | 0 | 0 % |
| 13 - 19 | 2 | 33 % | Men who have sex with men | 3 | 50 % |
| 20 – 29 | 3 | 50 % | Injection drug use | 0 | 0 % |
| 30 - 39 | 1 | 17 % | Transactional sex | 0 | 0 % |
| 40+ | 0 | 0 % | High risk heterosexual contact | 0 | 0 % |
| Total | 6 | 100 % | Heterosexual contact | 2 | 33 % |
| | | | Incarceration | 1 | 17 % |
| | | | Illegal drug use (non-injection) | 2 | 33 % |
| | | | <i>No Reported Risk</i> | <i>0</i> | |

Persons Living with HIV Infection

Region 1: New Orleans Region

as of March 31, 2016

Jefferson, Orleans, Plaquemines, St. Bernard Parishes

- **15,610** cumulative cases have been detected in Region 1, including **149** pediatric cases.
- **5,630** cumulative deaths among persons in Region 1 with HIV infection have occurred, including **44** pediatric deaths.
- **7,238** persons are currently living with HIV infection in Region 1.
Persons living with HIV: **3,443**
Persons living with AIDS: **3,795**

Gender

| | Cases | Percent |
|-------------------|--------------|--------------|
| Women | 1,817 | 25 % |
| Men | 5,333 | 74 % |
| Transgender Women | 86 | 1 % |
| Transgender Men | 2 | <1 % |
| Total | 7,238 | 100 % |

Age Category

| <u>Current Age</u> | Cases | Percent |
|--------------------|--------------|--------------|
| 0 - 12 | 15 | <1 % |
| 13 - 19 | 41 | 1 % |
| 20 - 24 | 243 | 3 % |
| 25 - 29 | 607 | 8 % |
| 30 - 34 | 733 | 10 % |
| 35 - 39 | 819 | 11 % |
| 40 - 44 | 792 | 11 % |
| 45 - 49 | 889 | 12 % |
| 50 - 54 | 1,105 | 15 % |
| 55 - 59 | 918 | 13 % |
| 60+ | 1,076 | 15 % |
| Total | 7,238 | 100 % |

Race/Ethnicity by Sex at Birth

| | Men | Women | Total | Percent |
|------------------------|--------------|--------------|--------------|--------------|
| Black, Non-Hispanic | 3,106 | 1,502 | 4,608 | 64 % |
| White, Non-Hispanic | 1,868 | 198 | 2,066 | 29 % |
| Hispanic | 363 | 105 | 468 | 6 % |
| Asian/Pacific Islander | 39 | 4 | 43 | 1 % |
| American Indian | 7 | 0 | 7 | <1 % |
| Unknown | 8 | 1 | 9 | <1 % |
| Multi-race | 28 | 9 | 37 | 1 % |
| Total | 5,419 | 1,819 | 7,238 | 100 % |

Exposure Category by Sex at Birth

| Cases with a reported risk: | Men | Women | Total | Percent |
|--|--------------|------------|--------------|--------------|
| Men who have sex with men | 3,088 | 0 | 3,088 | 61 % |
| Injection drug use | 294 | 194 | 488 | 10 % |
| Men who have sex with men and inject drugs | 306 | 0 | 306 | 6 % |
| Heterosexual contact | 342 | 745 | 1,087 | 21 % |
| Transfusion/Hemophilia | 16 | 13 | 29 | 1 % |
| Perinatal | 27 | 36 | 63 | 1 % |
| Total | 4,073 | 988 | 5,061 | 100 % |
| <i>No Identified Risk</i> | <i>1,346</i> | <i>831</i> | <i>2,177</i> | |

Persons Newly Diagnosed with HIV and AIDS

Region 1: New Orleans Region

Jefferson, Orleans, Plaquemines, St. Bernard Parishes

HIV CASES DIAGNOSED January - December 2015

- 394 persons were newly diagnosed with HIV in Region 1.

| <u>Gender</u> | <u>Cases</u> | <u>Percent</u> | <u>Race/Ethnicity</u> | <u>Cases</u> | <u>Percent</u> |
|-------------------|--------------|----------------|------------------------|--------------|----------------|
| Men | 319 | 81 % | Black, Non-Hispanic | 264 | 65 % |
| Women | 71 | 18 % | White, Non-Hispanic | 85 | 24 % |
| Transgender Women | 4 | 1 % | Hispanic | 37 | 9 % |
| Total | 394 | 100 % | Asian/Pacific Islander | 1 | <1 % |
| | | | American Indian | 1 | <1 % |
| | | | Unknown | 4 | 1 % |
| | | | Multi-race | 2 | 1 % |
| | | | Total | 394 | 100 % |

| <u>Age at Diagnosis</u> | <u>Cases</u> | <u>Percent</u> | <u>Cases with a reported risk (by birth sex):</u> | <u>Cases</u> | <u>Percent</u> |
|-------------------------|--------------|----------------|---|--------------|----------------|
| 0 -12 | 1 | <1 % | Men who have sex with men | 163 | 68 % |
| 13 - 24 | 95 | 24 % | Injection drug use | 23 | 10 % |
| 25 - 34 | 135 | 34 % | Men who have sex with men and inject drugs | 13 | 5 % |
| 35 - 44 | 81 | 21 % | Heterosexual contact | 41 | 17 % |
| 45+ | 82 | 21 % | Perinatal | 1 | <1 % |
| Total | 394 | 100 % | Total | 241 | 100 % |

No Identified Risk 153

AIDS CASES DIAGNOSED January - December 2015

- 175 persons were newly diagnosed with AIDS in Region 1.

| <u>Gender</u> | <u>Cases</u> | <u>Percent</u> | <u>Race/Ethnicity</u> | <u>Cases</u> | <u>Percent</u> |
|-------------------|--------------|----------------|-----------------------|--------------|----------------|
| Men | 124 | 71 % | Black, Non-Hispanic | 125 | 71 % |
| Women | 50 | 29 % | White, Non-Hispanic | 36 | 21 % |
| Transgender Women | 1 | <1 % | Hispanic | 12 | 7 % |
| Total | 175 | 100 % | American Indian | 1 | 1 % |
| | | | Multi-race | 1 | 1 % |
| | | | Total | 175 | 100 % |

| <u>Age at Diagnosis</u> | <u>Cases</u> | <u>Percent</u> | <u>Cases with a reported risk (by birth sex):</u> | <u>Cases</u> | <u>Percent</u> |
|-------------------------|--------------|----------------|---|--------------|----------------|
| 0 - 12 | 1 | 1 % | Men who have sex with men | 56 | 50 % |
| 13 - 24 | 16 | 9 % | Injection drug use | 13 | 12 % |
| 25 - 34 | 44 | 25 % | Men who have sex with men and inject drugs | 5 | 4 % |
| 35 - 44 | 47 | 27 % | Heterosexual contact | 37 | 33 % |
| 45+ | 67 | 38 % | Perinatal | 2 | 2 % |
| Total | 175 | 100 % | Total | 113 | 100 % |

No Identified Risk 62

Persons Diagnosed with Early Syphilis**Region 1: New Orleans Region***Jefferson, Orleans, Plaquemines, St. Bernard Parishes***PERSONS DIAGNOSED WITH P&S SYPHILIS January – March 2016**

- 24 persons were diagnosed with P&S Syphilis in Region 1 from January 1 through March 31, 2016.

| <u>Gender</u> | <u>Cases</u> | <u>Percent</u> | <u>Race/Ethnicity</u> | <u>Cases</u> | <u>Percent</u> |
|---------------|--------------|----------------|------------------------|--------------|----------------|
| Men | 21 | 88 % | American Indian | 0 | 0 % |
| Women | 3 | 13 % | Asian/Pacific Islander | 0 | 0 % |
| Total | 24 | 100 % | Black, Non-Hispanic | 16 | 67 % |
| | | | Hispanic | 0 | 0 % |
| | | | White, Non-Hispanic | 8 | 33 % |
| | | | Other/Multi-race | 0 | 0 % |
| | | | Unknown | 0 | 0 % |
| | | | Total | 24 | 100 % |

| <u>Age at Diagnosis</u> | <u>Cases</u> | <u>Percent</u> | <u>Cases with a reported risk (Can have >1 risk)</u> | <u>Cases</u> | <u>Percent</u> |
|-------------------------|--------------|----------------|---|--------------|----------------|
| 0 - 12 | 0 | 0 % | Men who have sex with men and inject drugs | 0 | 0 % |
| 13 - 19 | 1 | 4 % | Men who have sex with men | 14 | 70 % |
| 20 – 29 | 12 | 50 % | Injection drug use | 0 | 0 % |
| 30 - 39 | 7 | 29 % | Transactional sex | 0 | 0 % |
| 40+ | 4 | 17 % | High risk heterosexual contact | 0 | 0 % |
| Total | 24 | 100 % | Heterosexual contact | 7 | 35 % |
| | | | Incarceration | 2 | 10 % |
| | | | Illegal drug use (non-injection) | 1 | 5 % |
| | | | <i>No Reported Risk</i> | 4 | |

PERSONS DIAGNOSED WITH EARLY LATENT SYPHILIS January – March 2016

- 35 persons were diagnosed with Early Latent Syphilis in Region 1 from January 1 through March 31, 2016.

| <u>Gender</u> | <u>Cases</u> | <u>Percent</u> | <u>Race/Ethnicity</u> | <u>Cases</u> | <u>Percent</u> |
|---------------|--------------|----------------|------------------------|--------------|----------------|
| Men | 28 | 80 % | American Indian | 0 | 0 % |
| Women | 7 | 20 % | Asian/Pacific Islander | 1 | 3 % |
| Total | 35 | 100 % | Black, Non-Hispanic | 26 | 74 % |
| | | | Hispanic | 0 | 0 % |
| | | | White, Non-Hispanic | 8 | 23 % |
| | | | Other/Multi-race | 0 | 0 % |
| | | | Unknown | 0 | 0 % |
| | | | Total | 35 | 100 % |

| <u>Age at Diagnosis</u> | <u>Cases</u> | <u>Percent</u> | <u>Cases with a reported risk (Can have >1 risk)</u> | <u>Cases</u> | <u>Percent</u> |
|-------------------------|--------------|----------------|---|--------------|----------------|
| 0 - 12 | 0 | 0 % | Men who have sex with men and inject drugs | 0 | 0 % |
| 13 - 19 | 0 | 0 % | Men who have sex with men | 21 | 64 % |
| 20 – 29 | 15 | 43 % | Injection drug use | 0 | 0 % |
| 30 - 39 | 11 | 31 % | Transactional sex | 0 | 0 % |
| 40+ | 9 | 26 % | High risk heterosexual contact | 4 | 12 % |
| Total | 35 | 100 % | Heterosexual contact | 8 | 24 % |
| | | | Incarceration | 6 | 18 % |
| | | | Illegal drug use (non-injection) | 1 | 3 % |
| | | | <i>No Reported Risk</i> | 2 | |

**Persons Living with HIV Infection
Region 2: Baton Rouge Region
as of March 31, 2016**

Ascension, East Baton Rouge, East Feliciana, Iberville, Pointe Coupee, West Baton Rouge, West Feliciana Parishes

- **8,205** cumulative cases have been detected in Region 2, including **75** pediatric cases.
- **3,316** cumulative deaths among persons in Region 2 with HIV infection have occurred, including **15** pediatric deaths.
- **5,011** persons are currently living with HIV infection in Region 2.
Persons living with HIV: **2,374**
Persons living with AIDS: **2,637**

Gender

| | Cases | Percent |
|-------------------|--------------|----------------|
| Women | 1,803 | 36 % |
| Men | 3,168 | 63 % |
| Transgender Women | 40 | 1 % |
| Total | 5,011 | 100 % |

Age Category

| Current Age | Cases | Percent |
|--------------------|--------------|----------------|
| 0 - 12 | 18 | <1 % |
| 13 - 19 | 24 | <1 % |
| 20 - 24 | 199 | 4 % |
| 25 - 29 | 454 | 9 % |
| 30 - 34 | 553 | 11 % |
| 35 - 39 | 657 | 13 % |
| 40 - 44 | 624 | 12 % |
| 45 - 49 | 621 | 12 % |
| 50 - 54 | 691 | 14 % |
| 55 - 59 | 552 | 11 % |
| 60+ | 618 | 12 % |
| Total | 5,011 | 100 % |

Race/Ethnicity by Sex at Birth

| | Men | Women | Total | Percent |
|------------------------|--------------|--------------|--------------|----------------|
| Black, Non-Hispanic | 2,563 | 1,643 | 4,206 | 84 % |
| White, Non-Hispanic | 551 | 130 | 681 | 14 % |
| Hispanic | 61 | 17 | 78 | 2 % |
| Asian/Pacific Islander | 11 | 4 | 15 | <1 % |
| American Indian | 2 | 0 | 2 | <1 % |
| Multi-race | 20 | 9 | 29 | 1 % |
| Total | 3,208 | 1,803 | 5,011 | 100 % |

Exposure Category by Sex at Birth

| Cases with a reported risk: | Men | Women | Total | Percent |
|--|--------------|--------------|--------------|----------------|
| Men who have sex with men | 1339 | 0 | 1339 | 43 % |
| Injection drug use | 342 | 202 | 544 | 18 % |
| Men who have sex with men and inject drugs | 179 | 0 | 179 | 6 % |
| Heterosexual contact | 305 | 656 | 961 | 31 % |
| Transfusion/Hemophilia | 4 | 8 | 12 | <1 % |
| Perinatal | 31 | 25 | 56 | 2 % |
| Total | 2,200 | 891 | 3,091 | 100 % |
| <i>No Identified Risk</i> | <i>1,008</i> | <i>912</i> | <i>1,920</i> | |

Persons Newly Diagnosed with HIV and AIDS**Region 2: Baton Rouge Region***Ascension, East Baton Rouge, East Feliciana, Iberville, Pointe Coupee, West Baton Rouge, West Feliciana Parishes***HIV CASES DIAGNOSED January - December 2015**

- 268 persons were newly diagnosed with HIV in Region 2.

| <u>Gender</u> | Cases | Percent | <u>Race/Ethnicity</u> | Cases | Percent |
|-------------------|------------|--------------|------------------------|------------|--------------|
| Men | 189 | 71 % | Black, Non-Hispanic | 230 | 86 % |
| Women | 78 | 29 % | White, Non-Hispanic | 27 | 10 % |
| Transgender Women | 1 | <1 % | Hispanic | 8 | 3 % |
| Total | 268 | 100 % | Asian/Pacific Islander | 1 | <1 % |
| | | | Multi-race | 2 | 1 % |
| | | | Total | 268 | 100 % |

| <u>Age at Diagnosis</u> | Cases | Percent | <u>Cases with a reported risk (by birth sex):</u> | Cases | Percent |
|-------------------------|------------|--------------|---|------------|--------------|
| 13 - 24 | 74 | 28 % | Men who have sex with men | 93 | 73 % |
| 25 - 34 | 87 | 32 % | Injection drug use | 5 | 4 % |
| 35 - 44 | 52 | 19 % | Men who have sex with men and inject drugs | 4 | 3 % |
| 45+ | 55 | 21 % | Heterosexual contact | 25 | 20 % |
| Total | 268 | 100 % | Perinatal | 1 | 1 % |
| | | | Total | 128 | 100 % |

No Identified Risk 140

AIDS CASES DIAGNOSED January - December 2015

- 124 persons were newly diagnosed with AIDS in Region 2.

| <u>Gender</u> | Cases | Percent | <u>Race/Ethnicity</u> | Cases | Percent |
|-------------------|------------|--------------|------------------------|------------|--------------|
| Men | 86 | 69 % | Black, Non-Hispanic | 104 | 84 % |
| Women | 37 | 30 % | White, Non-Hispanic | 17 | 14 % |
| Transgender Women | 1 | 1 % | Hispanic | 1 | 1 % |
| Total | 124 | 100 % | Asian/Pacific Islander | 1 | 1 % |
| | | | Multi-race | 1 | 1 % |
| | | | Total | 124 | 100 % |

| <u>Age at Diagnosis</u> | Cases | Percent | <u>Cases with a reported risk (by birth sex):</u> | Cases | Percent |
|-------------------------|------------|--------------|---|-----------|--------------|
| 13 - 24 | 10 | 8 % | Men who have sex with men | 35 | 59 % |
| 25 - 34 | 37 | 30 % | Injection drug use | 6 | 10 % |
| 35 - 44 | 34 | 27 % | Men who have sex with men and inject drugs | 4 | 7 % |
| 45+ | 43 | 35 % | Heterosexual contact | 14 | 24 % |
| Total | 124 | 100 % | Total | 59 | 100 % |

No Identified Risk 65

Persons Diagnosed with Early Syphilis**Region 2: Baton Rouge Region***Ascension, East Baton Rouge, East Feliciana, Iberville, Pointe Coupee, West Baton Rouge, West Feliciana Parishes***PERSONS DIAGNOSED WITH P&S SYPHILIS January – March 2016**

- 26 persons were diagnosed with P&S Syphilis in Region 2 from January 1 through March 31, 2016.

| <u>Gender</u> | <u>Cases</u> | <u>Percent</u> | <u>Race/Ethnicity</u> | <u>Cases</u> | <u>Percent</u> |
|---------------|--------------|----------------|------------------------|--------------|----------------|
| Men | 20 | 77 % | American Indian | 0 | 0 % |
| Women | 6 | 23 % | Asian/Pacific Islander | 0 | 0 % |
| Total | 26 | 100 % | Black, Non-Hispanic | 21 | 81 % |
| | | | Hispanic | 0 | 0 % |
| | | | White, Non-Hispanic | 5 | 19 % |
| | | | Other/Multi-race | 0 | 0 % |
| | | | Unknown | 0 | 0 % |
| | | | Total | 26 | 100 % |

| <u>Age at Diagnosis</u> | <u>Cases</u> | <u>Percent</u> | <u>Cases with a reported risk (Can have >1 risk)</u> | <u>Cases</u> | <u>Percent</u> |
|-------------------------|--------------|----------------|---|--------------|----------------|
| 0 - 12 | 0 | 0 % | Men who have sex with men and inject drugs | 0 | 0 % |
| 13 - 19 | 3 | 12 % | Men who have sex with men | 15 | 60 % |
| 20 – 29 | 11 | 42 % | Injection drug use | 0 | 0 % |
| 30 - 39 | 5 | 19 % | Transactional sex | 1 | 4 % |
| 40+ | 7 | 27 % | High risk heterosexual contact | 2 | 8 % |
| Total | 26 | 100 % | Heterosexual contact | 8 | 32 % |
| | | | Incarceration | 0 | 0 % |
| | | | Illegal drug use (non-injection) | 0 | 0 % |
| | | | <i>No Reported Risk</i> | <i>1</i> | |

PERSONS DIAGNOSED WITH EARLY LATENT SYPHILIS January – March 2016

- 5 persons were diagnosed with Early Latent Syphilis in Region 2 from January 1 through March 31, 2016.

| <u>Gender</u> | <u>Cases</u> | <u>Percent</u> | <u>Race/Ethnicity</u> | <u>Cases</u> | <u>Percent</u> |
|---------------|--------------|----------------|------------------------|--------------|----------------|
| Men | 5 | 100 % | American Indian | 0 | 0 % |
| Women | 0 | 0 % | Asian/Pacific Islander | 0 | 0 % |
| Total | 5 | 100 % | Black, Non-Hispanic | 4 | 80 % |
| | | | Hispanic | 0 | 0 % |
| | | | White, Non-Hispanic | 1 | 20 % |
| | | | Other/Multi-race | 0 | 0 % |
| | | | Unknown | 0 | 0 % |
| | | | Total | 5 | 100 % |

| <u>Age at Diagnosis</u> | <u>Cases</u> | <u>Percent</u> | <u>Cases with a reported risk (Can have >1 risk)</u> | <u>Cases</u> | <u>Percent</u> |
|-------------------------|--------------|----------------|---|--------------|----------------|
| 0 - 12 | 0 | 0 % | Men who have sex with men and inject drugs | 0 | 0 % |
| 13 - 19 | 0 | 0 % | Men who have sex with men | 3 | 60 % |
| 20 – 29 | 1 | 20 % | Injection drug use | 0 | 0 % |
| 30 - 39 | 3 | 60 % | Transactional sex | 0 | 0 % |
| 40+ | 1 | 20 % | High risk heterosexual contact | 0 | 0 % |
| Total | 5 | 100 % | Heterosexual contact | 2 | 40 % |
| | | | Incarceration | 1 | 20 % |
| | | | Illegal drug use (non-injection) | 1 | 20 % |
| | | | <i>No Reported Risk</i> | <i>0</i> | |

**Persons Living with HIV Infection
Region 3: Houma Region
as of March 31, 2016**

Assumption, Lafourche, St. Charles, St. James, St. John the Baptist, St. Mary, Terrebonne Parishes

- **1,294** cumulative cases have been detected in Region 3, including **20** pediatric cases.
- **371** cumulative deaths among persons in Region 3 with HIV infection have occurred, including **2** pediatric deaths.
- **858** persons are currently living with HIV infection in Region 3.
Persons living with HIV: **388**
Persons living with AIDS: **470**

| <u>Gender</u> | <u>Cases</u> | <u>Percent</u> |
|-------------------|--------------|----------------|
| Women | 282 | 33 % |
| Men | 568 | 66 % |
| Transgender Women | 8 | 1 % |
| Total | 858 | 100 % |

Age Category

| <u>Current Age</u> | <u>Cases</u> | <u>Percent</u> |
|--------------------|--------------|----------------|
| 0 - 12 | 4 | <1 % |
| 13 - 19 | 8 | 1 % |
| 20 - 24 | 46 | 5 % |
| 25 - 29 | 90 | 10 % |
| 30 - 34 | 98 | 11 % |
| 35 - 39 | 90 | 10 % |
| 40 - 44 | 101 | 12 % |
| 45 - 49 | 115 | 13 % |
| 50 - 54 | 102 | 12 % |
| 55 - 59 | 98 | 11 % |
| 60+ | 106 | 12 % |
| Total | 858 | 100 % |

Race/Ethnicity by Sex at Birth

| | <u>Men</u> | <u>Women</u> | <u>Total</u> | <u>Percent</u> |
|------------------------|------------|--------------|--------------|----------------|
| Black, Non-Hispanic | 305 | 214 | 519 | 60 % |
| White, Non-Hispanic | 239 | 58 | 297 | 35 % |
| Hispanic | 26 | 6 | 32 | 4 % |
| Asian/Pacific Islander | 2 | 1 | 3 | <1 % |
| American Indian | 1 | 1 | 2 | <1 % |
| Multi-race | 3 | 2 | 5 | 1 % |
| Total | 576 | 282 | 858 | 100 % |

Exposure Category by Sex at Birth

| <u>Cases with a reported risk:</u> | <u>Men</u> | <u>Women</u> | <u>Total</u> | <u>Percent</u> |
|--|----------------|----------------|----------------|----------------|
| Men who have sex with men | 342 | 0 | 342 | 57 % |
| Injection drug use | 34 | 17 | 51 | 9 % |
| Men who have sex with men and inject drugs | 18 | 0 | 18 | 3 % |
| Heterosexual contact | 36 | 132 | 168 | 28 % |
| Perinatal | 6 | 11 | 17 | 3 % |
| Total | 436 | 160 | 596 | 100 % |
| <i>No Identified Risk</i> | <i>140</i> | <i>122</i> | <i>262</i> | |

Persons Newly Diagnosed with HIV and AIDS

Region 3: Houma Region

Assumption, Lafourche, St. Charles, St. James, St. John the Baptist, St. Mary, Terrebonne Parishes

HIV CASES DIAGNOSED January - December 2015

- 66 persons were newly diagnosed with HIV in Region 3.

| <u>Gender</u> | <u>Cases</u> | <u>Percent</u> | <u>Race/Ethnicity</u> | <u>Cases</u> | <u>Percent</u> |
|---------------|--------------|----------------|-----------------------|--------------|----------------|
| Men | 46 | 70 % | Black, Non-Hispanic | 41 | 62 % |
| Women | 20 | 30 % | White, Non-Hispanic | 19 | 29 % |
| Total | 66 | 100 % | Hispanic | 5 | 8 % |
| | | | American Indian | 1 | 2 % |
| | | | Total | 66 | 100 % |

| <u>Age at Diagnosis</u> | <u>Cases</u> | <u>Percent</u> | <u>Cases with a reported risk (by birth sex):</u> | <u>Cases</u> | <u>Percent</u> |
|-------------------------|--------------|----------------|---|--------------|----------------|
| 0 - 12 | 1 | 2 % | Men who have sex with men | 28 | 62 % |
| 13 - 24 | 19 | 29 % | Injection drug use | 4 | 9 % |
| 25 - 34 | 20 | 30 % | Men who have sex with men and inject drugs | 1 | 2 % |
| 35 - 44 | 12 | 18 % | Heterosexual contact | 11 | 24 % |
| 45+ | 14 | 21 % | Perinatal | 1 | 2 % |
| Total | 66 | 100 % | Total | 45 | 100 % |
| | | | <i>No Identified Risk</i> | <i>21</i> | |

AIDS CASES DIAGNOSED January - December 2015

- 26 persons were newly diagnosed with AIDS in Region 3.

| <u>Gender</u> | <u>Cases</u> | <u>Percent</u> | <u>Race/Ethnicity</u> | <u>Cases</u> | <u>Percent</u> |
|-------------------|--------------|----------------|-----------------------|--------------|----------------|
| Men | 23 | 88 % | Black, Non-Hispanic | 16 | 64 % |
| Women | 3 | 12 % | White, Non-Hispanic | 5 | 29 % |
| Transgender Women | 1 | 4 % | Hispanic | 4 | 7 % |
| Total | 26 | 100 % | American Indian | 1 | 4 % |
| | | | Total | 26 | 100 % |

| <u>Age at Diagnosis</u> | <u>Cases</u> | <u>Percent</u> | <u>Cases with a reported risk (by birth sex):</u> | <u>Cases</u> | <u>Percent</u> |
|-------------------------|--------------|----------------|---|--------------|----------------|
| 13 - 24 | 5 | 19 % | Men who have sex with men | 13 | 76 % |
| 25 - 34 | 8 | 31 % | Injection drug use | 1 | 6 % |
| 35 - 44 | 5 | 19 % | Heterosexual contact | 3 | 18 % |
| 45+ | 8 | 31 % | Total | 17 | 100 % |
| Total | 26 | 100 % | <i>No Identified Risk</i> | <i>9</i> | |

Persons Diagnosed with Early Syphilis**Region 3: Houma Region***Assumption, Lafourche, St. Charles, St. James, St. John the Baptist, St. Mary, Terrebonne Parishes***PERSONS DIAGNOSED WITH P&S SYPHILIS January – March 2016**

- 9 persons were diagnosed with P&S Syphilis in Region 3 from January 1 through March 31, 2016.

| <u>Gender</u> | <u>Cases</u> | <u>Percent</u> | <u>Race/Ethnicity</u> | <u>Cases</u> | <u>Percent</u> |
|---------------|--------------|----------------|------------------------|--------------|----------------|
| Men | 6 | 67 % | American Indian | 0 | 0 % |
| Women | 3 | 33 % | Asian/Pacific Islander | 0 | 0 % |
| Total | 9 | 100 % | Black, Non-Hispanic | 7 | 78 % |
| | | | Hispanic | 1 | 11 % |
| | | | White, Non-Hispanic | 1 | 11 % |
| | | | Other/Multi-race | 0 | 0 % |
| | | | Unknown | 0 | 0 % |
| | | | Total | 9 | 100 % |

| <u>Age at Diagnosis</u> | <u>Cases</u> | <u>Percent</u> | <u>Cases with a reported risk (Can have >1 risk)</u> | <u>Cases</u> | <u>Percent</u> |
|-------------------------|--------------|----------------|---|--------------|----------------|
| 0 - 12 | 0 | 0 % | Men who have sex with men and inject drugs | 0 | 0 % |
| 13 - 19 | 0 | 0 % | Men who have sex with men | 4 | 50 % |
| 20 – 29 | 7 | 78 % | Injection drug use | 0 | 0 % |
| 30 - 39 | 2 | 22 % | Transactional sex | 0 | 0 % |
| 40+ | 0 | 0 % | High risk heterosexual contact | 2 | 25 % |
| Total | 9 | 100 % | Heterosexual contact | 3 | 38 % |
| | | | Incarceration | 2 | 25 % |
| | | | Illegal drug use (non-injection) | 0 | 0 % |
| | | | <i>No Reported Risk</i> | <i>1</i> | |

PERSONS DIAGNOSED WITH EARLY LATENT SYPHILIS January – March 2016

- 4 persons were diagnosed with Early Latent Syphilis in Region 3 from January 1 through March 31, 2016.

| <u>Gender</u> | <u>Cases</u> | <u>Percent</u> | <u>Race/Ethnicity</u> | <u>Cases</u> | <u>Percent</u> |
|---------------|--------------|----------------|------------------------|--------------|----------------|
| Men | 2 | 50 % | American Indian | 0 | 0 % |
| Women | 2 | 50 % | Asian/Pacific Islander | 0 | 0 % |
| Total | 4 | 100 % | Black, Non-Hispanic | 4 | 100 % |
| | | | Hispanic | 0 | 0 % |
| | | | White, Non-Hispanic | 0 | 0 % |
| | | | Other/Multi-race | 0 | 0 % |
| | | | Unknown | 0 | 0 % |
| | | | Total | 4 | 100 % |

| <u>Age at Diagnosis</u> | <u>Cases</u> | <u>Percent</u> | <u>Cases with a reported risk (Can have >1 risk)</u> | <u>Cases</u> | <u>Percent</u> |
|-------------------------|--------------|----------------|---|--------------|----------------|
| 0 - 12 | 0 | 0 % | Men who have sex with men and inject drugs | 0 | 0 % |
| 13 - 19 | 1 | 25 % | Men who have sex with men | 1 | 25 % |
| 20 – 29 | 3 | 75 % | Injection drug use | 0 | 0 % |
| 30 - 39 | 0 | 0 % | Transactional sex | 0 | 0 % |
| 40+ | 0 | 0 % | High risk heterosexual contact | 0 | 0 % |
| Total | 4 | 100 % | Heterosexual contact | 3 | 75 % |
| | | | Incarceration | 0 | 0 % |
| | | | Illegal drug use (non-injection) | 0 | 0 % |
| | | | <i>No Reported Risk</i> | <i>0</i> | |

**Persons Living with HIV Infection
Region 4: Lafayette Region
as of March 31, 2016**

Acadia, Evangeline, Iberia, Lafayette, St. Landry, St. Martin, Vermilion Parishes

- **2,422** cumulative cases have been detected in Region 4, including **18** pediatric cases.
- **833** cumulative deaths among persons in Region 4 with HIV infection have occurred, including **5** pediatric deaths.
- **1,579** persons are currently living with HIV infection in Region 4.
Persons living with HIV: **761**
Persons living with AIDS: **818**

| <u>Gender</u> | <u>Cases</u> | <u>Percent</u> |
|-------------------|--------------|----------------|
| Women | 401 | 25 % |
| Men | 1,174 | 74 % |
| Transgender Women | 4 | <1 % |
| Total | 1,579 | 100 % |

| <u>Age Category</u> | | | <u>Race/Ethnicity by Sex at Birth</u> | | | | |
|---------------------|--------------|----------------|---------------------------------------|--------------|--------------|--------------|----------------|
| <u>Current Age</u> | <u>Cases</u> | <u>Percent</u> | | <u>Men</u> | <u>Women</u> | <u>Total</u> | <u>Percent</u> |
| 0 - 12 | 1 | <1 % | Black, Non-Hispanic | 625 | 275 | 900 | 57 % |
| 13 - 19 | 5 | <1 % | White, Non-Hispanic | 470 | 112 | 582 | 37 % |
| 20 - 24 | 73 | 5 % | Hispanic | 62 | 6 | 68 | 4 % |
| 25 - 29 | 145 | 9 % | Asian/Pacific Islander | 3 | 2 | 5 | <1 % |
| 30 - 34 | 141 | 9 % | American Indian | 3 | 3 | 6 | <1 % |
| 35 - 39 | 163 | 10 % | Multi-race | 15 | 3 | 18 | 1 % |
| 40 - 44 | 141 | 9 % | Total | 1,178 | 401 | 1,579 | 100 % |
| 45 - 49 | 218 | 14 % | | | | | |
| 50 - 54 | 278 | 18 % | | | | | |
| 55 - 59 | 205 | 13 % | | | | | |
| 60+ | 209 | 13 % | | | | | |
| Total | 1,579 | 100 % | | | | | |

Exposure Category by Sex at Birth

| <u>Cases with a reported risk:</u> | <u>Men</u> | <u>Women</u> | <u>Total</u> | <u>Percent</u> |
|--|------------|--------------|--------------|----------------|
| Men who have sex with men | 592 | 0 | 592 | 57 % |
| Injection drug use | 68 | 50 | 118 | 11 % |
| Men who have sex with men and inject drugs | 57 | 0 | 57 | 6 % |
| Heterosexual contact | 72 | 173 | 245 | 24 % |
| Transfusion/Hemophilia | 5 | 1 | 6 | 1 % |
| Confirmed Other | 0 | 1 | 1 | <1 % |
| Perinatal | 6 | 5 | 11 | 1 % |
| Total | 800 | 230 | 1,030 | 100 % |
| <i>No Identified Risk</i> | <i>378</i> | <i>171</i> | <i>549</i> | |

Persons Newly Diagnosed with HIV and AIDS**Region 4: Lafayette Region***Acadia, Evangeline, Iberia, Lafayette, St. Landry, St. Martin, Vermilion Parishes***HIV CASES DIAGNOSED January - December 2015**

- 96 persons were newly diagnosed with HIV in Region 4.

| <u>Gender</u> | Cases | Percent | <u>Race/Ethnicity</u> | Cases | Percent |
|---------------|--------------|----------------|------------------------|--------------|----------------|
| Men | 73 | 78 % | Black, Non-Hispanic | 54 | 70 % |
| Women | 23 | 22 % | White, Non-Hispanic | 32 | 24 % |
| Total | 96 | 100 % | Hispanic | 6 | 4 % |
| | | | Asian/Pacific Islander | 1 | 1 % |
| | | | American Indian | 2 | 2 % |
| | | | Multi-race | 1 | 1 % |
| | | | Total | 96 | 100 % |

| <u>Age at Diagnosis</u> | Cases | Percent | <u>Cases with a reported risk (by birth sex):</u> | Cases | Percent |
|-------------------------|--------------|----------------|---|--------------|----------------|
| 13 - 24 | 25 | 26 % | Men who have sex with men | 28 | 70 % |
| 25 - 34 | 38 | 40 % | Injection drug use | 1 | 2 % |
| 35 - 44 | 11 | 11 % | Heterosexual contact | 11 | 28 % |
| 45+ | 22 | 23 % | Total | 40 | 100 % |
| Total | 96 | 100 % | <i>No Identified Risk</i> | <i>56</i> | |

AIDS CASES DIAGNOSED January - December 2015

- 37 persons were newly diagnosed with AIDS in Region 4.

| <u>Gender</u> | Cases | Percent | <u>Race/Ethnicity</u> | Cases | Percent |
|---------------|--------------|----------------|-----------------------|--------------|----------------|
| Men | 30 | 81 % | Black, Non-Hispanic | 19 | 51 % |
| Women | 7 | 19 % | White, Non-Hispanic | 15 | 41 % |
| Total | 37 | 100 % | Hispanic | 1 | 3 % |
| | | | American Indian | 1 | 3 % |
| | | | Multi-race | 1 | 3 % |
| | | | Total | 37 | 100 % |

| <u>Age at Diagnosis</u> | Cases | Percent | <u>Cases with a reported risk (by birth sex):</u> | Cases | Percent |
|-------------------------|--------------|----------------|---|--------------|----------------|
| 13 - 24 | 4 | 11 % | Men who have sex with men | 13 | 65 % |
| 25 - 34 | 9 | 24 % | Heterosexual contact | 6 | 35 % |
| 35 - 44 | 6 | 16 % | Total | 20 | 100 % |
| 45+ | 18 | 49 % | <i>No Identified Risk</i> | <i>17</i> | |
| Total | 37 | 100 % | | | |

Persons Diagnosed with Early Syphilis**Region 4: Lafayette Region***Acadia, Evangeline, Iberia, Lafayette, St. Landry, St. Martin, Vermilion Parishes***PERSONS DIAGNOSED WITH P&S SYPHILIS January – March 2016**

- 18 persons were diagnosed with P&S Syphilis in Region 4 from January 1 through March 31, 2016.

| <u>Gender</u> | <u>Cases</u> | <u>Percent</u> | <u>Race/Ethnicity</u> | <u>Cases</u> | <u>Percent</u> |
|---------------|--------------|----------------|------------------------|--------------|----------------|
| Men | 11 | 61 % | American Indian | 0 | 0 % |
| Women | 7 | 39 % | Asian/Pacific Islander | 0 | 0 % |
| Total | 18 | 100 % | Black, Non-Hispanic | 12 | 67 % |
| | | | Hispanic | 1 | 6 % |
| | | | White, Non-Hispanic | 5 | 28 % |
| | | | Other/Multi-race | 0 | 0 % |
| | | | Unknown | 0 | 0 % |
| | | | Total | 18 | 100 % |

| <u>Age at Diagnosis</u> | <u>Cases</u> | <u>Percent</u> | <u>Cases with a reported risk (Can have >1 risk)</u> | <u>Cases</u> | <u>Percent</u> |
|-------------------------|--------------|----------------|---|--------------|----------------|
| 0 - 12 | 0 | 0 % | Men who have sex with men and inject drugs | 0 | 0 % |
| 13 - 19 | 0 | 0 % | Men who have sex with men | 4 | 25 % |
| 20 – 29 | 13 | 72 % | Injection drug use | 0 | 0 % |
| 30 - 39 | 2 | 11 % | Transactional sex | 1 | 6 % |
| 40+ | 3 | 17 % | High risk heterosexual contact | 3 | 19 % |
| Total | 18 | 100 % | Heterosexual contact | 10 | 63 % |
| | | | Incarceration | 1 | 6 % |
| | | | Illegal drug use (non-injection) | 2 | 13 % |
| | | | <i>No Reported Risk</i> | 2 | |

PERSONS DIAGNOSED WITH EARLY LATENT SYPHILIS January – March 2016

- 9 persons were diagnosed with Early Latent Syphilis in Region 4 from January 1 through March 31, 2016.

| <u>Gender</u> | <u>Cases</u> | <u>Percent</u> | <u>Race/Ethnicity</u> | <u>Cases</u> | <u>Percent</u> |
|---------------|--------------|----------------|------------------------|--------------|----------------|
| Men | 5 | 56 % | American Indian | 0 | 0 % |
| Women | 4 | 44 % | Asian/Pacific Islander | 0 | 0 % |
| Total | 9 | 100 % | Black, Non-Hispanic | 5 | 56 % |
| | | | Hispanic | 0 | 0 % |
| | | | White, Non-Hispanic | 4 | 44 % |
| | | | Other/Multi-race | 0 | 0 % |
| | | | Unknown | 0 | 0 % |
| | | | Total | 9 | 100 % |

| <u>Age at Diagnosis</u> | <u>Cases</u> | <u>Percent</u> | <u>Cases with a reported risk (Can have >1 risk)</u> | <u>Cases</u> | <u>Percent</u> |
|-------------------------|--------------|----------------|---|--------------|----------------|
| 0 - 12 | 0 | 0 % | Men who have sex with men and inject drugs | 0 | 0 % |
| 13 - 19 | 1 | 11 % | Men who have sex with men | 4 | 44 % |
| 20 – 29 | 4 | 44 % | Injection drug use | 1 | 11 % |
| 30 - 39 | 4 | 44 % | Transactional sex | 2 | 22 % |
| 40+ | 0 | 0 % | High risk heterosexual contact | 0 | 0 % |
| Total | 9 | 100 % | Heterosexual contact | 5 | 56 % |
| | | | Incarceration | 4 | 44 % |
| | | | Illegal drug use (non-injection) | 3 | 33 % |
| | | | <i>No Reported Risk</i> | 0 | |

Persons Living with HIV Infection

Region 5: Lake Charles Region

as of March 31, 2016

Allen, Beauregard, Calcasieu, Cameron, Jefferson Davis Parishes

- **1,451** cumulative cases have been detected in Region 5, including **10** pediatric cases.
- **544** cumulative deaths among persons in Region 5 with HIV infection have occurred, including **2** pediatric deaths.
- **1,084** persons are currently living with HIV infection in Region 5.
 - Persons living with HIV: **514**
 - Persons living with AIDS: **570**

| <u>Gender</u> | <u>Cases</u> | <u>Percent</u> |
|-------------------|--------------|----------------|
| Women | 253 | 23 % |
| Men | 827 | 76 % |
| Transgender Women | 4 | <1% |
| Total | 1,084 | 100% |

| <u>Age Category</u> | | | <u>Race/Ethnicity by Sex at Birth</u> | | | | |
|---------------------|--------------|----------------|---------------------------------------|--------------|--------------|----------------|--------------|
| <u>Current Age</u> | <u>Cases</u> | <u>Percent</u> | <u>Men</u> | <u>Women</u> | <u>Total</u> | <u>Percent</u> | |
| 0 - 12 | 2 | <1 % | Black, Non-Hispanic | 445 | 171 | 616 | 57 % |
| 13 - 19 | 5 | <1 % | White, Non-Hispanic | 283 | 74 | 357 | 33 % |
| 20 - 24 | 27 | 2 % | Hispanic | 86 | 4 | 90 | 8 % |
| 25 - 29 | 80 | 7 % | Asian/Pacific Islander | 5 | 0 | 5 | <1 % |
| 30 - 34 | 99 | 9 % | American Indian | 3 | 0 | 3 | <1 % |
| 35 - 39 | 125 | 12 % | Unknown | 1 | 0 | 1 | <1 % |
| 40 - 44 | 147 | 14 % | Multi-race | 8 | 4 | 12 | 1 % |
| 45 - 49 | 156 | 14 % | Total | 831 | 253 | 1,084 | 100 % |
| 50 - 54 | 181 | 17 % | | | | | |
| 55 - 59 | 141 | 13 % | | | | | |
| 60+ | 121 | 11 % | | | | | |
| Total | 1,084 | 100 % | | | | | |

Exposure Category by Sex at Birth

| <u>Cases with a reported risk:</u> | <u>Men</u> | <u>Women</u> | <u>Total</u> | <u>Percent</u> |
|--|------------|--------------|--------------|----------------|
| Men who have sex with men | 375 | 0 | 375 | 47 % |
| Injection drug use | 89 | 33 | 122 | 15 % |
| Men who have sex with men and inject drugs | 77 | 0 | 77 | 10 % |
| Heterosexual contact | 92 | 110 | 202 | 25 % |
| Transfusion/Hemophilia | 2 | 0 | 2 | <1 % |
| Perinatal | 6 | 9 | 15 | 2 % |
| Total | 641 | 152 | 793 | 100 % |
| <i>No Identified Risk</i> | <i>190</i> | <i>101</i> | <i>291</i> | |

Persons Newly Diagnosed with HIV and AIDS

Region 5: Lake Charles Region

Allen, Beauregard, Calcasieu, Cameron, Jefferson Davis Parishes

HIV CASES DIAGNOSED January - December 2015

- 52 persons were newly diagnosed with HIV in Region 5.

| <u>Gender</u> | <u>Cases</u> | <u>Percent</u> | <u>Race/Ethnicity</u> | <u>Cases</u> | <u>Percent</u> |
|---------------|--------------|----------------|-----------------------|--------------|----------------|
| Men | 40 | 77 % | Black, Non-Hispanic | 26 | 50 % |
| Women | 12 | 23 % | White, Non-Hispanic | 23 | 44 % |
| Total | 52 | 100 % | Hispanic | 2 | 4 % |
| | | | Unknown | 1 | 2 % |
| | | | Total | 52 | 100 % |

| <u>Age at Diagnosis</u> | <u>Cases</u> | <u>Percent</u> | <u>Cases with a reported risk (by birth sex):</u> | <u>Cases</u> | <u>Percent</u> |
|-------------------------|--------------|----------------|---|--------------|----------------|
| 13 - 24 | 10 | 19 % | Men who have sex with men | 15 | 65 % |
| 25 - 34 | 15 | 29 % | Injection drug use | 3 | 13 % |
| 35 - 44 | 12 | 23 % | Men who have sex with men and inject drugs | 1 | 4 % |
| 45+ | 15 | 29 % | Heterosexual contact | 4 | 17 % |
| Total | 52 | 100 % | Total | 23 | 100 % |

No Identified Risk 29

AIDS CASES DIAGNOSED January - December 2015

- 23 persons were newly diagnosed with AIDS in Region 5.

| <u>Gender</u> | <u>Cases</u> | <u>Percent</u> | <u>Race/Ethnicity</u> | <u>Cases</u> | <u>Percent</u> |
|---------------|--------------|----------------|-----------------------|--------------|----------------|
| Men | 17 | 74 % | Black, Non-Hispanic | 13 | 57 % |
| Women | 6 | 26 % | White, Non-Hispanic | 8 | 35 % |
| Total | 23 | 100 % | Hispanic | 2 | 9 % |
| | | | Total | 23 | 100 % |

| <u>Age at Diagnosis</u> | <u>Cases</u> | <u>Percent</u> | <u>Cases with a reported risk (by birth sex):</u> | <u>Cases</u> | <u>Percent</u> |
|-------------------------|--------------|----------------|---|--------------|----------------|
| 13 - 24 | 2 | 9 % | Men who have sex with men | 7 | 47 % |
| 25 - 34 | 8 | 35 % | Injection drug use | 3 | 20 % |
| 35 - 44 | 8 | 35 % | Men who have sex with men and inject drugs | 1 | 7 % |
| 45+ | 5 | 22 % | Heterosexual contact | 4 | 27 % |
| Total | 23 | 100 % | Total | 15 | 100 % |

No Identified Risk 8

Persons Diagnosed with Early Syphilis**Region 5: Lake Charles Region***Allen, Beauregard, Calcasieu, Cameron, Jefferson Davis Parishes***PERSONS DIAGNOSED WITH P&S SYPHILIS January – March 2016**

- 5 persons were diagnosed with P&S Syphilis in Region 5 from January 1 through March 31, 2016.

| <u>Gender</u> | <u>Cases</u> | <u>Percent</u> | <u>Race/Ethnicity</u> | <u>Cases</u> | <u>Percent</u> |
|---------------|--------------|----------------|------------------------|--------------|----------------|
| Men | 3 | 60 % | American Indian | 0 | 0 % |
| Women | 2 | 40 % | Asian/Pacific Islander | 0 | 0 % |
| Total | 5 | 100 % | Black, Non-Hispanic | 4 | 80 % |
| | | | Hispanic | 0 | 0 % |
| | | | White, Non-Hispanic | 1 | 20 % |
| | | | Other/Multi-race | 0 | 0 % |
| | | | Unknown | 0 | 0 % |
| | | | Total | 5 | 100 % |

| <u>Age at Diagnosis</u> | <u>Cases</u> | <u>Percent</u> | <u>Cases with a reported risk (Can have >1 risk)</u> | <u>Cases</u> | <u>Percent</u> |
|-------------------------|--------------|----------------|---|--------------|----------------|
| 0 - 12 | 0 | 0 % | Men who have sex with men and inject drugs | 0 | 0 % |
| 13 - 19 | 1 | 20 % | Men who have sex with men | 2 | 40 % |
| 20 – 29 | 2 | 40 % | Injection drug use | 1 | 20 % |
| 30 - 39 | 2 | 40 % | Transactional sex | 0 | 0 % |
| 40+ | 0 | 0 % | High risk heterosexual contact | 2 | 40 % |
| Total | 5 | 100 % | Heterosexual contact | 2 | 40 % |
| | | | Incarceration | 0 | 0 % |
| | | | Illegal drug use (non-injection) | 2 | 40 % |
| | | | <i>No Reported Risk</i> | 0 | |

PERSONS DIAGNOSED WITH EARLY LATENT SYPHILIS January – December 2016

- 2 persons were diagnosed with Early Latent Syphilis in Region 5 from January 1 through March 31, 2016.

| <u>Gender</u> | <u>Cases</u> | <u>Percent</u> | <u>Race/Ethnicity</u> | <u>Cases</u> | <u>Percent</u> |
|---------------|--------------|----------------|------------------------|--------------|----------------|
| Men | 1 | 50 % | American Indian | 0 | 0 % |
| Women | 1 | 50 % | Asian/Pacific Islander | 0 | 0 % |
| Total | 2 | 100 % | Black, Non-Hispanic | 1 | 50 % |
| | | | Hispanic | 0 | 0 % |
| | | | White, Non-Hispanic | 1 | 50 % |
| | | | Other/Multi-race | 0 | 0 % |
| | | | Unknown | 0 | 0 % |
| | | | Total | 2 | 100 % |

| <u>Age at Diagnosis</u> | <u>Cases</u> | <u>Percent</u> | <u>Cases with a reported risk (Can have >1 risk)</u> | <u>Cases</u> | <u>Percent</u> |
|-------------------------|--------------|----------------|---|--------------|----------------|
| 0 - 12 | 0 | 0 % | Men who have sex with men and inject drugs | 0 | 0 % |
| 13 - 19 | 1 | 50 % | Men who have sex with men | 1 | 50 % |
| 20 – 29 | 1 | 50 % | Injection drug use | 0 | 0 % |
| 30 - 39 | 0 | 0 % | Transactional sex | 0 | 0 % |
| 40+ | 0 | 0 % | High risk heterosexual contact | 0 | 0 % |
| Total | 2 | 100 % | Heterosexual contact | 1 | 50 % |
| | | | Incarceration | 0 | 0 % |
| | | | Illegal drug use (non-injection) | 0 | 0 % |
| | | | <i>No Reported Risk</i> | 0 | |

**Persons Living with HIV Infection
Region 6: Alexandria Region
as of March 31, 2016**

Avoyelles, Catahoula, Concordia, Grant, La Salle, Rapides, Vernon, Winn Parishes

- **1,609** cumulative cases have been detected in Region 6, including **15** pediatric cases.
- **526** cumulative deaths among persons in Region 6 with HIV infection have occurred, including **5** pediatric deaths.
- **958** persons are currently living with HIV infection in Region 6.
Persons living with HIV: **493**
Persons living with AIDS: **465**

Gender

| | Cases | Percent |
|-------------------|--------------|----------------|
| Women | 271 | 28 % |
| Men | 683 | 71 % |
| Transgender Women | 4 | <1 % |
| Total | 958 | 100% |

Age Category

| Current Age | Cases | Percent |
|--------------------|--------------|----------------|
| 0 - 12 | 1 | <1 % |
| 13 - 19 | 5 | 1 % |
| 20 - 24 | 33 | 3 % |
| 25 - 29 | 70 | 7 % |
| 30 - 34 | 104 | 11 % |
| 35 - 39 | 130 | 14 % |
| 40 - 44 | 114 | 12 % |
| 45 - 49 | 120 | 13 % |
| 50 - 54 | 151 | 16 % |
| 55 - 59 | 105 | 11 % |
| 60+ | 125 | 13 % |
| Total | 958 | 100 % |

Race/Ethnicity by Sex at Birth

| | Men | Women | Total | Percent |
|------------------------|------------|--------------|--------------|----------------|
| Black, Non-Hispanic | 425 | 214 | 639 | 67 % |
| White, Non-Hispanic | 211 | 47 | 258 | 27 % |
| Hispanic | 44 | 5 | 49 | 5 % |
| Asian/Pacific Islander | 3 | 0 | 3 | <1 % |
| American Indian | 0 | 2 | 2 | <1 % |
| Multi-race | 4 | 3 | 7 | 1 % |
| Total | 687 | 271 | 958 | 100 % |

Exposure Category by Sex at Birth

| Cases with a reported risk: | Men | Women | Total | Percent |
|--|------------|--------------|--------------|----------------|
| Men who have sex with men | 294 | 0 | 294 | 46 % |
| Injection drug use | 65 | 22 | 87 | 14 % |
| Men who have sex with men and inject drugs | 39 | 0 | 39 | 6 % |
| Heterosexual contact | 71 | 131 | 202 | 32 % |
| Transfusion/Hemophilia | 4 | 2 | 6 | 1 % |
| Perinatal | 3 | 2 | 5 | 1 % |
| Total | 496 | 157 | 633 | 100 % |

| | | | |
|---------------------------|------------|------------|------------|
| <i>No Identified Risk</i> | <i>211</i> | <i>114</i> | <i>325</i> |
|---------------------------|------------|------------|------------|

Persons Newly Diagnosed with HIV and AIDS**Region 6: Alexandria Region***Avoyelles, Catahoula, Concordia, Grant, La Salle, Rapides, Vernon, Winn Parishes***HIV CASES DIAGNOSED January - December 2015**

- 60 persons were newly diagnosed with HIV in Region 6.

| <u>Gender</u> | Cases | Percent | <u>Race/Ethnicity</u> | Cases | Percent |
|----------------------|--------------|----------------|------------------------------|--------------|----------------|
| Men | 45 | 77 % | Black, Non-Hispanic | 36 | 49 % |
| Women | 15 | 21 % | White, Non-Hispanic | 15 | 32 % |
| Transgender Woman | 1 | 1 % | Hispanic | 8 | 19 % |
| Total | 60 | 100 % | American Indian | 1 | 2 % |
| | | | Total | 60 | 100 % |

| <u>Age at Diagnosis</u> | Cases | Percent | <u>Cases with a reported risk (by birth sex):</u> | Cases | Percent |
|--------------------------------|--------------|----------------|--|--------------|----------------|
| 13 - 24 | 13 | 22 % | Men who have sex with men | 15 | 62 % |
| 25 - 34 | 20 | 33 % | Injection drug use | 2 | 8 % |
| 35 - 44 | 15 | 25 % | Heterosexual contact | 7 | 29 % |
| 45+ | 12 | 20 % | Total | 24 | 100 % |
| Total | 60 | 100 % | <i>No Identified Risk</i> | <i>36</i> | |

AIDS CASES DIAGNOSED January - December 2015

- 25 persons were newly diagnosed with AIDS in Region 6.

| <u>Gender</u> | Cases | Percent | <u>Race/Ethnicity</u> | Cases | Percent |
|----------------------|--------------|----------------|------------------------------|--------------|----------------|
| Men | 13 | 52 % | Black, Non-Hispanic | 17 | 64 % |
| Women | 11 | 44 % | White, Non-Hispanic | 5 | 20 % |
| Transgender Women | 1 | 4 % | Hispanic | 2 | 16 % |
| Total | 25 | 100 % | American Indian | 1 | 4 % |
| | | | Total | 25 | 100 % |

| <u>Age at Diagnosis</u> | Cases | Percent | <u>Cases with a reported risk (by birth sex):</u> | Cases | Percent |
|--------------------------------|--------------|----------------|--|--------------|----------------|
| 13 - 24 | 1 | 4 % | Men who have sex with men | 5 | 73 % |
| 25 - 34 | 10 | 40 % | Injection drug use | 2 | 9 % |
| 35 - 44 | 7 | 28 % | Heterosexual contact | 6 | 18 % |
| 45+ | 7 | 28 % | Total | 13 | 100 % |
| Total | 25 | 100 % | <i>No Identified Risk</i> | <i>12</i> | |

Persons Diagnosed with Early Syphilis**Region 6: Alexandria Region**

Avoyelles, Catahoula, Concordia, Grant, La Salle, Rapides, Vernon, Winn Parishes

PERSONS DIAGNOSED WITH P&S SYPHILIS January – March 2016

- 8 persons were diagnosed with P&S Syphilis in Region 6 from January 1 through March 31, 2016.

| <u>Gender</u> | <u>Cases</u> | <u>Percent</u> | <u>Race/Ethnicity</u> | <u>Cases</u> | <u>Percent</u> |
|---------------|--------------|----------------|------------------------|--------------|----------------|
| Men | 8 | 100 % | American Indian | 0 | 0 % |
| Women | 0 | 0 % | Asian/Pacific Islander | 0 | 0 % |
| Total | 8 | 100 % | Black, Non-Hispanic | 6 | 75 % |
| | | | Hispanic | 1 | 13 % |
| | | | White, Non-Hispanic | 1 | 13 % |
| | | | Other/Multi-race | 0 | 0 % |
| | | | Unknown | 0 | 0 % |
| | | | Total | 8 | 100 % |

| <u>Age at Diagnosis</u> | <u>Cases</u> | <u>Percent</u> | <u>Cases with a reported risk (Can have >1 risk)</u> | <u>Cases</u> | <u>Percent</u> |
|-------------------------|--------------|----------------|---|--------------|----------------|
| 0 - 12 | 0 | 0 % | Men who have sex with men and inject drugs | 0 | 0 % |
| 13 - 19 | 1 | 13 % | Men who have sex with men | 4 | 57 % |
| 20 – 29 | 4 | 50 % | Injection drug use | 0 | 0 % |
| 30 - 39 | 3 | 38 % | Transactional sex | 3 | 43 % |
| 40+ | 0 | 0 % | High risk heterosexual contact | 5 | 71 % |
| Total | 8 | 100 % | Heterosexual contact | 1 | 14 % |
| | | | Incarceration | 1 | 14 % |
| | | | Illegal drug use (non-injection) | 3 | 43 % |
| | | | <i>No Reported Risk</i> | <i>1</i> | |

PERSONS DIAGNOSED WITH EARLY LATENT SYPHILIS January – March 2016

- 7 persons were diagnosed with Early Latent Syphilis in Region 6 from January 1 through March 31, 2016.

| <u>Gender</u> | <u>Cases</u> | <u>Percent</u> | <u>Race/Ethnicity</u> | <u>Cases</u> | <u>Percent</u> |
|---------------|--------------|----------------|------------------------|--------------|----------------|
| Men | 4 | 57 % | American Indian | 0 | 0 % |
| Women | 3 | 43 % | Asian/Pacific Islander | 0 | 0 % |
| Total | 7 | 100 % | Black, Non-Hispanic | 3 | 43 % |
| | | | Hispanic | 0 | 0 % |
| | | | White, Non-Hispanic | 4 | 57 % |
| | | | Other/Multi-race | 0 | 0 % |
| | | | Unknown | 0 | 0 % |
| | | | Total | 7 | 100 % |

| <u>Age at Diagnosis</u> | <u>Cases</u> | <u>Percent</u> | <u>Cases with a reported risk (Can have >1 risk)</u> | <u>Cases</u> | <u>Percent</u> |
|-------------------------|--------------|----------------|---|--------------|----------------|
| 0 - 12 | 0 | 0 % | Men who have sex with men and inject drugs | 1 | 14 % |
| 13 - 19 | 1 | 14 % | Men who have sex with men | 2 | 29 % |
| 20 – 29 | 3 | 43 % | Injection drug use | 0 | 0 % |
| 30 - 39 | 2 | 29 % | Transactional sex | 0 | 0 % |
| 40+ | 1 | 14 % | High risk heterosexual contact | 3 | 43 % |
| Total | 7 | 100 % | Heterosexual contact | 3 | 43 % |
| | | | Incarceration | 1 | 14 % |
| | | | Illegal drug use (non-injection) | 3 | 43 % |
| | | | <i>No Reported Risk</i> | <i>0</i> | |

Persons Living with HIV Infection

Region 7: Shreveport Region

as of March 31, 2016

Bienville, Bossier, Caddo, Claiborne, DeSoto, Natchitoches, Red River, Sabine, Webster Parishes

- **2,650** cumulative cases have been detected in Region 7, including **27** pediatric cases.
- **1,034** cumulative deaths among persons in Region 7 with HIV infection have occurred, including **10** pediatric deaths.
- **1,793** persons are currently living with HIV infection in Region 7.
Persons living with HIV: **906**
Persons living with AIDS: **887**

| <u>Gender</u> | <u>Cases</u> | <u>Percent</u> |
|-------------------|--------------|----------------|
| Women | 526 | 29 % |
| Men | 1,259 | 70 % |
| Transgender Women | 8 | <1 % |
| Total | 1,793 | 100 % |

Age Category

| <u>Current Age</u> | <u>Cases</u> | <u>Percent</u> |
|--------------------|--------------|----------------|
| 0 - 12 | 2 | <1 % |
| 13 - 19 | 14 | 1 % |
| 20 - 24 | 103 | 6 % |
| 25 - 29 | 219 | 12 % |
| 30 - 34 | 202 | 11 % |
| 35 - 39 | 237 | 13 % |
| 40 - 44 | 192 | 11 % |
| 45 - 49 | 217 | 12 % |
| 50 - 54 | 235 | 13 % |
| 55 - 59 | 177 | 10 % |
| 60+ | 195 | 11 % |
| Total | 1,793 | 100 % |

Race/Ethnicity by Sex at Birth

| | <u>Men</u> | <u>Women</u> | <u>Total</u> | <u>Percent</u> |
|------------------------|--------------|--------------|--------------|----------------|
| Black, Non-Hispanic | 886 | 430 | 1316 | 73 % |
| White, Non-Hispanic | 332 | 84 | 416 | 23 % |
| Hispanic | 24 | 7 | 31 | 2 % |
| Asian/Pacific Islander | 5 | 0 | 5 | <1 % |
| American Indian | 11 | 2 | 13 | 1 % |
| Unknown | 1 | 0 | 1 | <1 % |
| Multi-race | 8 | 3 | 11 | 1 % |
| Total | 1,267 | 526 | 1,793 | 100 % |

Exposure Category by Sex at Birth

| <u>Cases with a reported risk:</u> | <u>Men</u> | <u>Women</u> | <u>Total</u> | <u>Percent</u> |
|--|------------|--------------|--------------|----------------|
| Men who have sex with men | 712 | 0 | 712 | 58 % |
| Injection drug use | 50 | 49 | 99 | 8 % |
| Men who have sex with men and inject drugs | 87 | 0 | 87 | 7 % |
| Heterosexual contact | 85 | 225 | 310 | 25 % |
| Transfusion/Hemophilia | 3 | 3 | 6 | <1 % |
| Perinatal | 6 | 9 | 15 | 1 % |
| Total | 943 | 286 | 1,229 | 100 % |
| <i>No Identified Risk</i> | <i>324</i> | <i>240</i> | <i>564</i> | |

Persons Newly Diagnosed with HIV and AIDS

Region 7: Shreveport Region

Bienville, Bossier, Caddo, Claiborne, DeSoto, Natchitoches, Red River, Sabine, Webster Parishes

HIV CASES DIAGNOSED January - December 2015

- 138 persons were newly diagnosed with HIV in Region 7.

| <u>Gender</u> | <u>Cases</u> | <u>Percent</u> | <u>Race/Ethnicity</u> | <u>Cases</u> | <u>Percent</u> |
|-------------------|--------------|----------------|-----------------------|--------------|----------------|
| Men | 105 | 76 % | Black, Non-Hispanic | 105 | 76 % |
| Women | 33 | 24 % | White, Non-Hispanic | 24 | 17 % |
| Transgender Women | 2 | 1 % | Hispanic | 4 | 3 % |
| Total | 138 | 100 % | American Indian | 5 | 4 % |
| | | | Total | 138 | 100 % |

| <u>Age at Diagnosis</u> | <u>Cases</u> | <u>Percent</u> | <u>Cases with a reported risk (by birth sex):</u> | <u>Cases</u> | <u>Percent</u> |
|-------------------------|--------------|----------------|---|--------------|----------------|
| 13 - 24 | 39 | 28 % | Men who have sex with men | 43 | 69 % |
| 25 - 34 | 39 | 28 % | Injection drug use | 4 | 6 % |
| 35 - 44 | 21 | 15 % | Heterosexual contact | 15 | 24 % |
| 45+ | 39 | 28 % | Total | 62 | 100 % |
| Total | 138 | 100 % | <i>No Identified Risk</i> | <i>76</i> | |

AIDS CASES DIAGNOSED January - December 2015

- 51 persons were newly diagnosed with AIDS in Region 7.

| <u>Gender</u> | <u>Cases</u> | <u>Percent</u> | <u>Race/Ethnicity</u> | <u>Cases</u> | <u>Percent</u> |
|-------------------|--------------|----------------|-----------------------|--------------|----------------|
| Men | 37 | 73 % | Black, Non-Hispanic | 42 | 82 % |
| Women | 14 | 27 % | White, Non-Hispanic | 6 | 12 % |
| Transgender Women | 1 | 2 % | American Indian | 2 | 4 % |
| Total | 51 | 100 % | Multi-race | 1 | 2 % |
| | | | Total | 51 | 100 % |

| <u>Age at Diagnosis</u> | <u>Cases</u> | <u>Percent</u> | <u>Cases with a reported risk (by birth sex):</u> | <u>Cases</u> | <u>Percent</u> |
|-------------------------|--------------|----------------|---|--------------|----------------|
| 13 - 24 | 4 | 8 % | Men who have sex with men | 16 | 65 % |
| 25 - 34 | 15 | 29 % | Heterosexual contact | 8 | 19 % |
| 35 - 44 | 14 | 27 % | Total | 24 | 100 % |
| 45+ | 18 | 35 % | <i>No Identified Risk</i> | <i>27</i> | |
| Total | 51 | 100 % | | | |

Persons Diagnosed with Early Syphilis**Region 7: Shreveport Region***Bienville, Bossier, Caddo, Claiborne, DeSoto, Natchitoches, Red River, Sabine, Webster Parishes***PERSONS DIAGNOSED WITH P&S SYPHILIS January – March 2016**

- 32 persons were diagnosed with P&S Syphilis in Region 7 from January 1 through March 31, 2016.

| <u>Gender</u> | <u>Cases</u> | <u>Percent</u> | <u>Race/Ethnicity</u> | <u>Cases</u> | <u>Percent</u> |
|---------------|--------------|----------------|------------------------|--------------|----------------|
| Men | 21 | 66 % | American Indian | 0 | 0 % |
| Women | 11 | 34 % | Asian/Pacific Islander | 0 | 0 % |
| Total | 32 | 100 % | Black, Non-Hispanic | 26 | 81 % |
| | | | Hispanic | 1 | 3 % |
| | | | White, Non-Hispanic | 5 | 16 % |
| | | | Other/Multi-race | 0 | 0 % |
| | | | Unknown | 0 | 0 % |
| | | | Total | 32 | 100 % |

| <u>Age at Diagnosis</u> | <u>Cases</u> | <u>Percent</u> | <u>Cases with a reported risk (Can have >1 risk)</u> | <u>Cases</u> | <u>Percent</u> |
|-------------------------|--------------|----------------|---|--------------|----------------|
| 0 - 12 | 0 | 0 % | Men who have sex with men and inject drugs | 0 | 0 % |
| 13 - 19 | 5 | 16 % | Men who have sex with men | 6 | 21 % |
| 20 – 29 | 12 | 38 % | Injection drug use | 1 | 3 % |
| 30 - 39 | 12 | 38 % | Transactional sex | 0 | 0 % |
| 40+ | 3 | 9 % | High risk heterosexual contact | 6 | 21 % |
| Total | 32 | 100 % | Heterosexual contact | 16 | 55 % |
| | | | Incarceration | 6 | 21 % |
| | | | Illegal drug use (non-injection) | 2 | 7 % |
| | | | <i>No Reported Risk</i> | 3 | |

PERSONS DIAGNOSED WITH EARLY LATENT SYPHILIS January – March 2016

- 15 persons were diagnosed with Early Latent Syphilis in Region 7 from January 1 through March 31, 2016.

| <u>Gender</u> | <u>Cases</u> | <u>Percent</u> | <u>Race/Ethnicity</u> | <u>Cases</u> | <u>Percent</u> |
|---------------|--------------|----------------|------------------------|--------------|----------------|
| Men | 6 | 40 % | American Indian | 0 | 0 % |
| Women | 9 | 60 % | Asian/Pacific Islander | 0 | 0 % |
| Total | 15 | 100 % | Black, Non-Hispanic | 13 | 87 % |
| | | | Hispanic | 0 | 0 % |
| | | | White, Non-Hispanic | 2 | 13 % |
| | | | Other/Multi-race | 0 | 0 % |
| | | | Unknown | 0 | 0 % |
| | | | Total | 15 | 100 % |

| <u>Age at Diagnosis</u> | <u>Cases</u> | <u>Percent</u> | <u>Cases with a reported risk (Can have >1 risk)</u> | <u>Cases</u> | <u>Percent</u> |
|-------------------------|--------------|----------------|---|--------------|----------------|
| 0 - 12 | 0 | 0 % | Men who have sex with men and inject drugs | 0 | 0 % |
| 13 - 19 | 4 | 27 % | Men who have sex with men | 2 | 13 % |
| 20 – 29 | 7 | 47 % | Injection drug use | 0 | 0 % |
| 30 - 39 | 2 | 13 % | Transactional sex | 1 | 7 % |
| 40+ | 2 | 13 % | High risk heterosexual contact | 3 | 20 % |
| Total | 15 | 100 % | Heterosexual contact | 10 | 67 % |
| | | | Incarceration | 3 | 20 % |
| | | | Illegal drug use (non-injection) | 0 | 0 % |
| | | | <i>No Reported Risk</i> | 0 | |

Persons Living with HIV Infection

Region 8: Monroe Region

as of March 31, 2016

Caldwell, E. Carroll, Franklin, Jackson, Lincoln, Madison, Morehouse, Ouachita, Richland, Tensas, Union, W. Carroll Parishes

- **1,893** cumulative cases have been detected in Region 8, including **18** pediatric cases.
- **717** cumulative deaths among persons in Region 8 with HIV infection have occurred, including **3** pediatric deaths.
- **1,089** persons are currently living with HIV infection in Region 8.
Persons living with HIV: **511**
Persons living with AIDS: **578**

| Gender | Cases | Percent |
|-------------------|--------------|----------------|
| Women | 373 | 34 % |
| Men | 711 | 65 % |
| Transgender Women | 5 | <1 % |
| Total | 1,089 | 100 % |

Age Category

| Current Age | Cases | Percent |
|--------------------|--------------|----------------|
| 0 - 12 | 5 | <1 % |
| 13 - 19 | 9 | 1 % |
| 20 - 24 | 55 | 5 % |
| 25 - 29 | 89 | 8 % |
| 30 - 34 | 89 | 8 % |
| 35 - 39 | 145 | 13 % |
| 40 - 44 | 141 | 13 % |
| 45 - 49 | 130 | 12 % |
| 50 - 54 | 159 | 15 % |
| 55 - 59 | 123 | 11 % |
| 60+ | 144 | 13 % |
| Total | 1,089 | 100 % |

Race/Ethnicity by Sex at Birth

| | Men | Women | Total | Percent |
|------------------------|------------|--------------|--------------|----------------|
| Black, Non-Hispanic | 517 | 321 | 838 | 77 % |
| White, Non-Hispanic | 172 | 41 | 213 | 20 % |
| Hispanic | 12 | 4 | 16 | 1 % |
| Asian/Pacific Islander | 1 | 1 | 2 | <1 % |
| American Indian | 2 | 3 | 5 | <1 % |
| Unknown | 1 | 0 | 1 | <1 % |
| Multi-race | 11 | 3 | 14 | 1 % |
| Total | 716 | 373 | 1,089 | 100 % |

Exposure Category by Sex at Birth

| Cases with a reported risk: | Men | Women | Total | Percent |
|--|------------|--------------|--------------|----------------|
| Men who have sex with men | 387 | 0 | 387 | 53 % |
| Injection drug use | 36 | 28 | 64 | 9 % |
| Men who have sex with men and inject drugs | 35 | 0 | 35 | 5 % |
| Heterosexual contact | 72 | 162 | 234 | 32 % |
| Transfusion/Hemophilia | 1 | 3 | 4 | 1 % |
| Perinatal | 2 | 10 | 12 | 2 % |
| Total | 533 | 203 | 736 | 100 % |
| <i>No Identified Risk</i> | <i>183</i> | <i>170</i> | <i>353</i> | |

Persons Newly Diagnosed with HIV and AIDS

Region 8: Monroe Region

Caldwell, E. Carroll, Franklin, Jackson, Lincoln, Madison, Morehouse, Ouachita, Richland, Tensas, Union, W. Carroll Parishes

HIV CASES DIAGNOSED January - December 2015

- 68 persons were newly diagnosed with HIV in Region 8.

| <u>Gender</u> | <u>Cases</u> | <u>Percent</u> | <u>Race/Ethnicity</u> | <u>Cases</u> | <u>Percent</u> |
|---------------|--------------|----------------|-----------------------|--------------|----------------|
| Men | 52 | 76 % | Black, Non-Hispanic | 56 | 82 % |
| Women | 16 | 24 % | White, Non-Hispanic | 12 | 18 % |
| Total | 68 | 100 % | Total | 68 | 100 % |

| <u>Age at Diagnosis</u> | <u>Cases</u> | <u>Percent</u> | <u>Cases with a reported risk (by birth sex):</u> | <u>Cases</u> | <u>Percent</u> |
|-------------------------|--------------|----------------|---|--------------|----------------|
| 13 - 24 | 19 | 28 % | Men who have sex with men | 20 | 69 % |
| 25 - 34 | 18 | 26 % | Men who have sex with men and inject drugs | 1 | 3 % |
| 35 - 44 | 8 | 12 % | Heterosexual contact | 8 | 28 % |
| 45+ | 23 | 34 % | Total | 29 | 100 % |
| Total | 68 | 100 % | <i>No Identified Risk</i> | 39 | |

AIDS CASES DIAGNOSED January - December 2015

- 35 persons were newly diagnosed with AIDS in Region 8.

| <u>Gender</u> | <u>Cases</u> | <u>Percent</u> | <u>Race/Ethnicity</u> | <u>Cases</u> | <u>Percent</u> |
|---------------|--------------|----------------|-----------------------|--------------|----------------|
| Men | 19 | 54 % | Black, Non-Hispanic | 31 | 89 % |
| Women | 16 | 46 % | White, Non-Hispanic | 4 | 11 % |
| Total | 35 | 100 % | Total | 35 | 100 % |

| <u>Age at Diagnosis</u> | <u>Cases</u> | <u>Percent</u> | <u>Cases with a reported risk (by birth sex):</u> | <u>Cases</u> | <u>Percent</u> |
|-------------------------|--------------|----------------|---|--------------|----------------|
| 13 - 24 | 2 | 6 % | Men who have sex with men | 4 | 40 % |
| 25 - 34 | 11 | 31 % | Heterosexual contact | 6 | 60 % |
| 35 - 44 | 6 | 17 % | Total | 10 | 100 % |
| 45+ | 16 | 39 % | <i>No Identified Risk</i> | 25 | |
| Total | 35 | 100 % | | | |

Persons Diagnosed with Early Syphilis**Region 8: Monroe Region**

Caldwell, E. Carroll, Franklin, Jackson, Lincoln, Madison, Morehouse, Ouachita, Richland, Tensas, Union, W. Carroll Parishes

PERSONS DIAGNOSED WITH P&S SYPHILIS January – March 2016

- 9 persons were diagnosed with P&S Syphilis in Region 8 from January 1 through March 31, 2016.

| <u>Gender</u> | <u>Cases</u> | <u>Percent</u> | <u>Race/Ethnicity</u> | <u>Cases</u> | <u>Percent</u> |
|---------------|--------------|----------------|------------------------|--------------|----------------|
| Men | 5 | 56 % | American Indian | 0 | 0 % |
| Women | 4 | 44 % | Asian/Pacific Islander | 0 | 0 % |
| Total | 9 | 100 % | Black, Non-Hispanic | 8 | 89 % |
| | | | Hispanic | 0 | 0 % |
| | | | White, Non-Hispanic | 1 | 11 % |
| | | | Other/Multi-race | 0 | 0 % |
| | | | Unknown | 0 | 0 % |
| | | | Total | 9 | 100 % |

| <u>Age at Diagnosis</u> | <u>Cases</u> | <u>Percent</u> | <u>Cases with a reported risk (Can have >1 risk)</u> | <u>Cases</u> | <u>Percent</u> |
|-------------------------|--------------|----------------|---|--------------|----------------|
| 0 - 12 | 0 | 0 % | Men who have sex with men and inject drugs | 0 | 0 % |
| 13 - 19 | 1 | 11 % | Men who have sex with men | 2 | 22 % |
| 20 – 29 | 5 | 56 % | Injection drug use | 1 | 11 % |
| 30 - 39 | 2 | 22 % | Transactional sex | 1 | 11 % |
| 40+ | 1 | 11 % | High risk heterosexual contact | 2 | 22 % |
| Total | 9 | 100 % | Heterosexual contact | 6 | 67 % |
| | | | Incarceration | 0 | 0 % |
| | | | Illegal drug use (non-injection) | 0 | 0 % |
| | | | <i>No Reported Risk</i> | 0 | |

PERSONS DIAGNOSED WITH EARLY LATENT SYPHILIS January – March 2016

- 3 persons were diagnosed with Early Latent Syphilis in Region 8 from January 1 through March 31, 2016.

| <u>Gender</u> | <u>Cases</u> | <u>Percent</u> | <u>Race/Ethnicity</u> | <u>Cases</u> | <u>Percent</u> |
|---------------|--------------|----------------|------------------------|--------------|----------------|
| Men | 1 | 33 % | American Indian | 0 | 0 % |
| Women | 2 | 67 % | Asian/Pacific Islander | 0 | 0 % |
| Total | 3 | 100 % | Black, Non-Hispanic | 3 | 100 % |
| | | | Hispanic | 0 | 0 % |
| | | | White, Non-Hispanic | 0 | 0 % |
| | | | Other/Multi-race | 0 | 0 % |
| | | | Unknown | 0 | 0 % |
| | | | Total | 3 | 100 % |

| <u>Age at Diagnosis</u> | <u>Cases</u> | <u>Percent</u> | <u>Cases with a reported risk (Can have >1 risk)</u> | <u>Cases</u> | <u>Percent</u> |
|-------------------------|--------------|----------------|---|--------------|----------------|
| 0 - 12 | 0 | 0 % | Men who have sex with men and inject drugs | 0 | 0 % |
| 13 - 19 | 1 | 33 % | Men who have sex with men | 0 | 0 % |
| 20 – 29 | 1 | 33 % | Injection drug use | 0 | 0 % |
| 30 - 39 | 1 | 33 % | Transactional sex | 0 | 0 % |
| 40+ | 0 | 0 % | High risk heterosexual contact | 0 | 0 % |
| Total | 3 | 100 % | Heterosexual contact | 3 | 100 % |
| | | | Incarceration | 0 | 0 % |
| | | | Illegal drug use (non-injection) | 0 | 0 % |
| | | | <i>No Reported Risk</i> | 0 | |

**Persons Living with HIV Infection
Region 9: Hammond/Slidell Region
as of March 31, 2016**

Livingston, St. Helena, St. Tammany, Tangipahoa, Washington Parishes

- **1,695** cumulative cases have been detected in Region 9, including **19** pediatric cases.
- **499** cumulative deaths among persons in Region 9 with HIV infection have occurred, Including **5** pediatric deaths.
- **1,302** persons are currently living with HIV infection in Region 9.
Persons living with HIV: **596**
Persons living with AIDS: **706**

| <u>Gender</u> | <u>Cases</u> | <u>Percent</u> |
|-------------------|--------------|----------------|
| Women | 331 | 25 % |
| Men | 963 | 74 % |
| Transgender Women | 8 | 1% |
| Total | 1,302 | 100 % |

Age Category

| <u>Current Age</u> | <u>Cases</u> | <u>Percent</u> |
|--------------------|--------------|----------------|
| 0 - 12 | 4 | <1 % |
| 13 - 19 | 4 | <1 % |
| 20 - 24 | 33 | 3 % |
| 25 - 29 | 93 | 7 % |
| 30 - 34 | 120 | 9 % |
| 35 - 39 | 133 | 10 % |
| 40 - 44 | 144 | 11 % |
| 45 - 49 | 198 | 15 % |
| 50 - 54 | 210 | 16 % |
| 55 - 59 | 201 | 15 % |
| 60+ | 162 | 12 % |
| Total | 1,302 | 100 % |

Race/Ethnicity by Sex at Birth

| | <u>Men</u> | <u>Women</u> | <u>Total</u> | <u>Percent</u> |
|------------------------|------------|--------------|--------------|----------------|
| Black, Non-Hispanic | 451 | 208 | 659 | 51 % |
| White, Non-Hispanic | 480 | 109 | 589 | 45 % |
| Hispanic | 31 | 12 | 43 | 3 % |
| Asian/Pacific Islander | 3 | 2 | 5 | <1 % |
| American Indian | 1 | 0 | 1 | <1 % |
| Multi-race | 5 | 0 | 5 | <1 % |
| Total | 971 | 331 | 1,302 | 100 % |

Exposure Category by Sex at Birth

| <u>Cases with a reported risk:</u> | <u>Men</u> | <u>Women</u> | <u>Total</u> | <u>Percent</u> |
|--|------------|--------------|--------------|----------------|
| Men who have sex with men | 515 | 0 | 515 | 58 % |
| Injection drug use | 56 | 46 | 102 | 12 % |
| Men who have sex with men and inject drugs | 60 | 0 | 60 | 7 % |
| Heterosexual contact | 72 | 118 | 190 | 21 % |
| Transfusion/Hemophilia | 5 | 0 | 5 | 1 % |
| Perinatal | 9 | 5 | 14 | 2 % |
| Total | 717 | 169 | 886 | 100 % |

| | | | |
|---------------------------|------------|------------|------------|
| <i>No Identified Risk</i> | <i>254</i> | <i>162</i> | <i>416</i> |
|---------------------------|------------|------------|------------|

Persons Newly Diagnosed with HIV and AIDS

Region 9: Hammond/Slidell Region

Livingston, St. Helena, St. Tammany, Tangipahoa, Washington Parishes

HIV CASES DIAGNOSED January - December 2015

- 65 persons were newly diagnosed with HIV in Region 9.

| <u>Gender</u> | <u>Cases</u> | <u>Percent</u> | <u>Race/Ethnicity</u> | <u>Cases</u> | <u>Percent</u> |
|-------------------|--------------|----------------|------------------------|--------------|----------------|
| Men | 45 | 69 % | Black, Non-Hispanic | 41 | 63 % |
| Women | 19 | 29 % | White, Non-Hispanic | 19 | 29 % |
| Transgender Women | 1 | 2 % | Hispanic | 4 | 6 % |
| Total | 65 | 100 % | Asian/Pacific Islander | 1 | 2 % |
| | | | Total | 65 | 100 % |

| <u>Age at Diagnosis</u> | <u>Cases</u> | <u>Percent</u> | <u>Cases with a reported risk (by birth sex):</u> | <u>Cases</u> | <u>Percent</u> |
|-------------------------|--------------|----------------|---|--------------|----------------|
| 13 - 24 | 4 | 6 % | Men who have sex with men | 17 | 68 % |
| 25 - 34 | 23 | 35 % | Men who have sex with men and inject drugs | 1 | 4 % |
| 35 - 44 | 18 | 28 % | Heterosexual contact | 7 | 28 % |
| 45+ | 20 | 31 % | Total | 25 | 100 % |
| Total | 65 | 100 % | <i>No Identified Risk</i> | <i>40</i> | |

AIDS CASES DIAGNOSED January - December 2015

- 42 persons were newly diagnosed with AIDS in Region 9.

| <u>Gender</u> | <u>Cases</u> | <u>Percent</u> | <u>Race/Ethnicity</u> | <u>Cases</u> | <u>Percent</u> |
|---------------|--------------|----------------|------------------------|--------------|----------------|
| Men | 32 | 76 % | Black, Non-Hispanic | 26 | 62 % |
| Women | 10 | 24 % | White, Non-Hispanic | 14 | 33 % |
| Total | 42 | 100 % | Hispanic | 1 | 2 % |
| | | | Asian/Pacific Islander | 1 | 2 % |
| | | | Total | 42 | 100 % |

| <u>Age at Diagnosis</u> | <u>Cases</u> | <u>Percent</u> | <u>Cases with a reported risk (by birth sex):</u> | <u>Cases</u> | <u>Percent</u> |
|-------------------------|--------------|----------------|---|--------------|----------------|
| 13 - 24 | 3 | 7 % | Men who have sex with men | 11 | 55 % |
| 25 - 34 | 10 | 24 % | Injection drug use | 2 | 10 % |
| 35 - 44 | 8 | 19 % | Heterosexual contact | 7 | 35 % |
| 45+ | 21 | 50 % | Total | 20 | 100 % |
| Total | 42 | 100 % | <i>No Identified Risk</i> | <i>22</i> | |

Persons Diagnosed with Early Syphilis**Region 9: Hammond/Slidell Region***Livingston, St. Helena, St. Tammany, Tangipahoa, Washington Parishes***PERSONS DIAGNOSED WITH P&S SYPHILIS January – March 2016**

- 1 persons were diagnosed with P&S Syphilis in Region 9 from January 1 through March 31, 2016.

| <u>Gender</u> | <u>Cases</u> | <u>Percent</u> | <u>Race/Ethnicity</u> | <u>Cases</u> | <u>Percent</u> |
|---------------|--------------|----------------|------------------------|--------------|----------------|
| Men | 1 | 100 % | American Indian | 0 | 0 % |
| Women | 0 | 0 % | Asian/Pacific Islander | 0 | 0 % |
| Total | 1 | 100 % | Black, Non-Hispanic | 0 | 0 % |
| | | | Hispanic | 0 | 0 % |
| | | | White, Non-Hispanic | 1 | 100 % |
| | | | Other/Multi-race | 0 | 0 % |
| | | | Unknown | 0 | 0 % |
| | | | Total | 1 | 100 % |

| <u>Age at Diagnosis</u> | <u>Cases</u> | <u>Percent</u> | <u>Cases with a reported risk (Can have >1 risk)</u> | <u>Cases</u> | <u>Percent</u> |
|-------------------------|--------------|----------------|---|--------------|----------------|
| 0 - 12 | 0 | 0 % | Men who have sex with men and inject drugs | 0 | 0 % |
| 13 - 19 | 1 | 100 % | Men who have sex with men | 0 | 0 % |
| 20 – 29 | 0 | 0 % | Injection drug use | 0 | 0 % |
| 30 - 39 | 0 | 0 % | Transactional sex | 0 | 0 % |
| 40+ | 0 | 0 % | High risk heterosexual contact | 0 | 0 % |
| Total | 1 | 100 % | Heterosexual contact | 0 | 0 % |
| | | | Incarceration | 0 | 0 % |
| | | | Illegal drug use (non-injection) | 0 | 0 % |
| | | | <i>No Reported Risk</i> | <i>1</i> | |



PERSONS DIAGNOSED WITH EARLY LATENT SYPHILIS January – March 2016

- 5 persons were diagnosed with Early Latent Syphilis in Region 9 from January 1 through March 31, 2016.


| <u>Gender</u> | <u>Cases</u> | <u>Percent</u> | <u>Race/Ethnicity</u> | <u>Cases</u> | <u>Percent</u> |
|---------------|--------------|----------------|------------------------|--------------|----------------|
| Men | 4 | 80 % | American Indian | 0 | 0 % |
| Women | 1 | 20 % | Asian/Pacific Islander | 0 | 0 % |
| Total | 5 | 100 % | Black, Non-Hispanic | 4 | 80 % |
| | | | Hispanic | 1 | 20 % |
| | | | White, Non-Hispanic | 0 | 0 % |
| | | | Other/Multi-race | 0 | 0 % |
| | | | Unknown | 0 | 0 % |
| | | | Total | 5 | 100 % |

| <u>Age at Diagnosis</u> | <u>Cases</u> | <u>Percent</u> | <u>Cases with a reported risk (Can have >1 risk)</u> | <u>Cases</u> | <u>Percent</u> |
|-------------------------|--------------|----------------|---|--------------|----------------|
| 0 - 12 | 0 | 0 % | Men who have sex with men and inject drugs | 0 | 0 % |
| 13 - 19 | 0 | 0 % | Men who have sex with men | 3 | 60 % |
| 20 – 29 | 2 | 40 % | Injection drug use | 0 | 0 % |
| 30 - 39 | 3 | 60 % | Transactional sex | 0 | 0 % |
| 40+ | 0 | 0 % | High risk heterosexual contact | 0 | 0 % |
| Total | 5 | 100 % | Heterosexual contact | 1 | 20 % |
| | | | Incarceration | 1 | 20 % |
| | | | Illegal drug use (non-injection) | 1 | 20 % |
| | | | <i>No Reported Risk</i> | <i>0</i> | |


10-Year Trends in New HIV Diagnoses and New AIDS Diagnoses Louisiana, (2006-2015)



10-Year Trends in Persons Living With HIV Infection Louisiana (2006-2015)



7-Year Trends in Persons Diagnosed with P&S Syphilis Louisiana (2009-2015)



7-Year Trends in Persons Diagnosed with Early Syphilis Louisiana (2009-2015)

