STI/HIV/Hepatitis B & C Update

2022 Data Tables

Monroe Region 8



Reported Cases of HIV Region 8 - Monroe, 2022

Part			2022 New Diagnoses ^a		2022	2022 Prevalence ^b		
Race/Ethnicity		'	Cases	%	Rated	Cases	%	Rated
Black		Total	67	100.0%	19.3	1,212	100.0%	348.7
Hispanic/Latinx 5 7.5% 53.3 56 4.6% 597.3 White 18 26.9% 9.0 267 22.0% 133.3 20ther 0 0 0.0% n/a 26 2.1% n/a 27 20.0% 29.4 826 68.3% 489.0 27 27 28 28 28 28 28 28	Race/Ethr	nicitye						
White		Black	44	65.7%	33.9	862	71.2%	664.4
Other O O O O O O O O O		Hispanic/Latinx	5	7.5%	53.3	56	4.6%	597.3
Female		White	18	26.9%	9.0	267	22.0%	133.3
Female		Other	0	0.0%	n/a	26	2.1%	n/a
Female	Gender							
Transgender Male to Female 1 1.5% n/a 15 1.2% n/a 1 1.5% n/a 0 0.0% 0.0 0.0% 0.		Female	16	23.9%	9.0	371	30.4%	209.3
Ages Transgender Female to Male 0 0.0% n/a 0 0.0% n/a 0-14 0 0.0% 0.0 1 0.1% n/a 15-19 4 6.0% n/a 11 0.9% 44.0 20-29 30 44.8% 60.1 167 13.8% 334.6 30-39 15 22.4% 34.4 244 20.1% 560.2 40-49 4 6.0% n/a 301 24.8% 745.6 50+ 14 20.9% 11.5 488 40.3% 399.3 Riskh GBM 40 59.7% n/a 577 48.0% n/a PVID 6 9.0% n/a 58 4.8% n/a PWID 6 9.0% n/a 467 38.9% n/a High Risk Heterosexual 19 28.4% n/a 41 1.1% n/a Porinatal 0 0.0% <t< td=""><td></td><td>Male</td><td>50</td><td>74.6%</td><td>29.4</td><td>826</td><td>68.3%</td><td>489.0</td></t<>		Male	50	74.6%	29.4	826	68.3%	489.0
Ages		Transgender Male to Female	1	1.5%	n/a	15	1.2%	n/a
0-14		Transgender Female to Male	0	0.0%	n/a	0	0.0%	n/a
0-14	Ageg							
20-29 30 44.8% 60.1 167 13.8% 334.6	- 3	0-14	0	0.0%	0.0	1	0.1%	n/a
30-39		15-19	4	6.0%	n/a	11	0.9%	44.0
A0-49		20-29	30	44.8%	60.1	167	13.8%	334.6
Riskh		30-39	15	22.4%	34.4	244	20.1%	560.2
Caldwell Caldwell		40-49	4	6.0%	n/a	301	24.8%	745.6
GBM		50+	14	20.9%	11.5	488	40.3%	399.3
PWID 6 9.0% n/a 82 6.8% n/a GBM/PWID 2 3.0% n/a 58 4.8% n/a High Risk Heterosexual 19 28.4% n/a 467 38.9% n/a Perinatal 0 0.0% n/a 13 1.1% n/a Transfusion/Hemophilia 0 0.0% n/a 4 0.3% n/a Other 0 0.0% n/a 0 0.0% n/a 0 0.0% n/a 13 1.1% n/a East Carroll 1 1.5% n/a 31 2.6% 429.4 Franklin 5 7.5% 25.4 57 4.7% 289.8 Jackson 5 7.5% 33.6 48 4.0% 322.7 Lincoln 10 14.9% 20.8 110 9.1% 228.4 Madison 2 3.0% n/a 35 2.9% 357.2 Morehouse 1 1.5% n/a 57 4.7% 227.8 Ouachita 33 49.3% 20.8 70.3 58.0% 442.8 Richland 1 1.5% n/a 34 2.8% 841 Union 4 6.0% n/a 32 2.6% 151.7	Riskh							
GBM/PWID 2 3.0% n/a 58 4.8% n/a High Risk Heterosexual 19 28.4% n/a 467 38.9% n/a Perinatal 0 0.0% n/a 13 1.1% n/a Transfusion/Hemophilia 0 0.0% n/a 4 0.3% n/a Other 0 0.0% n/a 0 0.0% n/a 0 0.0% n/a Other 0 0.0% n/a 0 0.0% n/a 0 0.0% n/a Other 0 0.0% n/a 0 0.0% 0.0% n/a 0 0.0% n		GBM	40	59.7%	n/a	577	48.0%	n/a
High Risk Heterosexual 19 28.4% n/a 467 38.9% n/a Perinatal 0 0.0% n/a 13 1.1% n/a 17 17 18 19 19 19 19 19 19 19		PWID	6	9.0%	n/a	82	6.8%	n/a
Perinatal 0 0.0% n/a 13 1.1% n/a Transfusion/Hemophilia 0 0.0% n/a 4 0.3% n/a 0ther 0 0.0% n/a 0 0.0% n/a 0 0.0% n/a 0 0.0% n/a 1.7% 219.4 East Carroll 1 1.5% n/a 31 2.6% 429.4 Franklin 5 7.5% 25.4 57 4.7% 289.8 Jackson 5 7.5% 33.6 48 4.0% 322.7 Lincoln 10 14.9% 20.8 110 9.1% 228.4 Madison 2 3.0% n/a 35 2.9% 357.2 Morehouse 1 1.5% n/a 57 4.7% 227.8 Ouachita 33 49.3% 20.8 703 58.0% 442.8 Richland 1 1.5% n/a 63 5.2% 318.1 Tensas 1 1.5% n/a 34 2.8% 841 Union 4 6.0% n/a 32 2.6% 151.7		GBM/PWID	2	3.0%	n/a	58	4.8%	n/a
Transfusion/Hemophilia 0 0.0% n/a 0 0.3% n/a 0 0.0% n/a 0 0.0% n/a 0 0.0% n/a n/a 0 0.0% n/a n/a 0 0.0% n/a n/a		High Risk Heterosexual	19	28.4%	n/a	467	38.9%	n/a
Parish Other 0 0.0% n/a 0 0.0% n/a Caldwell 2 3.0% n/a 21 1.7% 219.4 East Carroll 1 1.5% n/a 31 2.6% 429.4 Franklin 5 7.5% 25.4 57 4.7% 289.8 Jackson 5 7.5% 33.6 48 4.0% 322.7 Lincoln 10 14.9% 20.8 110 9.1% 228.4 Madison 2 3.0% n/a 35 2.9% 357.2 Morehouse 1 1.5% n/a 57 4.7% 227.8 Ouachita 33 49.3% 20.8 703 58.0% 442.8 Richland 1 1.5% n/a 63 5.2% 318.1 Tensas 1 1.5% n/a 34 2.8% 841 Union 4 6.0% n/a 32		Perinatal	0	0.0%	n/a	13	1.1%	n/a
Parish Caldwell East Carroll 1 1.5% n/a Franklin 5 7.5% 25.4 Jackson 5 7.5% 33.6 48 4.0% 322.7 Lincoln 10 14.9% 20.8 Madison 2 3.0% n/a 35 2.9% 357.2 Morehouse 1 1.5% n/a Ouachita 33 49.3% 20.8 Richland 1 1.5% n/a Richland 1 1.5% n/a 1 1.5% n/a Richland 1 1.5% n/a		Transfusion/Hemophilia	0	0.0%	n/a	4	0.3%	n/a
Caldwell 2 3.0% n/a 21 1.7% 219.4 East Carroll 1 1.5% n/a 31 2.6% 429.4 Franklin 5 7.5% 25.4 57 4.7% 289.8 Jackson 5 7.5% 33.6 48 4.0% 322.7 Lincoln 10 14.9% 20.8 110 9.1% 228.4 Madison 2 3.0% n/a 35 2.9% 357.2 Morehouse 1 1.5% n/a 57 4.7% 227.8 Ouachita 33 49.3% 20.8 703 58.0% 442.8 Richland 1 1.5% n/a 63 5.2% 318.1 Tensas 1 1.5% n/a 34 2.8% 841 Union 4 6.0% n/a 32 2.6% 151.7		Other	0	0.0%	n/a	0	0.0%	n/a
East Carroll 1 1.5% n/a 31 2.6% 429.4 Franklin 5 7.5% 25.4 57 4.7% 289.8 Jackson 5 7.5% 33.6 48 4.0% 322.7 Lincoln 10 14.9% 20.8 110 9.1% 228.4 Madison 2 3.0% n/a 35 2.9% 357.2 Morehouse 1 1.5% n/a 57 4.7% 227.8 Ouachita 33 49.3% 20.8 703 58.0% 442.8 Richland 1 1.5% n/a 63 5.2% 318.1 Tensas 1 1.5% n/a 34 2.8% 841 Union 4 6.0% n/a 32 2.6% 151.7	Parish [,]							
Franklin 5 7.5% 25.4 57 4.7% 289.8 Jackson 5 7.5% 33.6 48 4.0% 322.7 Lincoln 10 14.9% 20.8 110 9.1% 228.4 Madison 2 3.0% n/a 35 2.9% 357.2 Morehouse 1 1.5% n/a 57 4.7% 227.8 Ouachita 33 49.3% 20.8 703 58.0% 442.8 Richland 1 1.5% n/a 63 5.2% 318.1 Tensas 1 1.5% n/a 34 2.8% 841 Union 4 6.0% n/a 32 2.6% 151.7		Caldwell	2	3.0%	n/a	21	1.7%	219.4
Jackson 5 7.5% 33.6 48 4.0% 322.7 Lincoln 10 14.9% 20.8 110 9.1% 228.4 Madison 2 3.0% n/a 35 2.9% 357.2 Morehouse 1 1.5% n/a 57 4.7% 227.8 Ouachita 33 49.3% 20.8 703 58.0% 442.8 Richland 1 1.5% n/a 63 5.2% 318.1 Tensas 1 1.5% n/a 34 2.8% 841 Union 4 6.0% n/a 32 2.6% 151.7		East Carroll	1	1.5%	n/a	31	2.6%	429.4
Lincoln 10 14.9% 20.8 110 9.1% 228.4 Madison 2 3.0% n/a 35 2.9% 357.2 Morehouse 1 1.5% n/a 57 4.7% 227.8 Ouachita 33 49.3% 20.8 703 58.0% 442.8 Richland 1 1.5% n/a 63 5.2% 318.1 Tensas 1 1.5% n/a 34 2.8% 841 Union 4 6.0% n/a 32 2.6% 151.7		Franklin	5	7.5%	25.4	57	4.7%	289.8
Madison 2 3.0% n/a 35 2.9% 357.2 Morehouse 1 1.5% n/a 57 4.7% 227.8 Ouachita 33 49.3% 20.8 703 58.0% 442.8 Richland 1 1.5% n/a 63 5.2% 318.1 Tensas 1 1.5% n/a 34 2.8% 841 Union 4 6.0% n/a 32 2.6% 151.7		Jackson	5	7.5%	33.6	48	4.0%	322.7
Morehouse 1 1.5% n/a 57 4.7% 227.8 Ouachita 33 49.3% 20.8 703 58.0% 442.8 Richland 1 1.5% n/a 63 5.2% 318.1 Tensas 1 1.5% n/a 34 2.8% 841 Union 4 6.0% n/a 32 2.6% 151.7		Lincoln	10	14.9%	20.8	110	9.1%	228.4
Ouachita 33 49.3% 20.8 703 58.0% 442.8 Richland 1 1.5% n/a 63 5.2% 318.1 Tensas 1 1.5% n/a 34 2.8% 841 Union 4 6.0% n/a 32 2.6% 151.7		Madison	2	3.0%	n/a	35	2.9%	357.2
Richland 1 1.5% n/a 63 5.2% 318.1 Tensas 1 1.5% n/a 34 2.8% 841 Union 4 6.0% n/a 32 2.6% 151.7		Morehouse	1	1.5%	n/a	57	4.7%	227.8
Tensas 1 1.5% n/a 34 2.8% 841 Union 4 6.0% n/a 32 2.6% 151.7		Ouachita	33	49.3%	20.8	703	58.0%	442.8
Union 4 6.0% n/a 32 2.6% 151.7		Richland	1	1.5%	n/a	63	5.2%	318.1
		Tensas	1	1.5%	n/a	34	2.8%	841
West Carroll 2 3.0% n/a 21 1.7% 218.9		Union	4	6.0%	n/a	32	2.6%	151.7
		West Carroll	2	3.0%	n/a	21	1.7%	218.9

New diagnoses include only individuals whose initial HIV diagnosis was made in Louisiana.
 Prevalent cases include all cases presumed to be alive and living in Louisiana, regardless of the state of initial HIV report.
 Cases with unknown/missing values for a demographic variable have been excluded from the counts and percentages for the demographic category. Total number of new diagnoses and prevalence includes all cases, regardless of demographic variable completeness.
 Cases per 100,000 population. Rates not available (n/a) for case counts less than 5. Rates not available for other/unknown race groups, transgender groups, and risk exposure groups.
 Race/ethnicity "Other" category does not include missing or unknown race/ethnicity.
 Cases are presented by current gender identity rather than sex at birth.
 Age when first diagnosed with HIV is used for new HIV diagnoses. Current age is used for HIV prevalence. Age categories align with 2021 census age groups.
 Risk shows imputed risk, not reported risk. Persons with no imputed risk are not included in the total percentage. GBM = Gay, bisexual, & other men who have sex with men; PWID = Persons who inject drugs.
 Parish where first diagnosed with HIV is used for new HIV diagnoses. Parish of current residence is used for HIV prevalence. Persons without a parish of residence are not included in the total percentage.

Reported Cases of Primary and Secondary Syphilis Region 8 - Monroe, 2022

New Diagnosesa

31.0

35.2

47.8%

52.2%

	Cases ^b	%	Ratec
Total	115	100.0%	33.1
ce/Ethnicity ^d			
Black	84	73.7%	64.7
Hispanic/Latinx	1	0.9%	n/a
White	24	21.1%	12.0
Other	5	4.4%	n/a

55

60

Affected Subpopulations Femaleh	Cases ^b	%
Black Female 10-19	5	4.4%
Black Female 20-29	23	20.2%
Black Female 30-39	8	7.0%
White Female 30-39	6	5.3%
White Female 40-49	5	4.4%

Agef

Gendere

Female

Male

1	0-19	9	7.8%	18.6
2	20-29	56	48.7%	112.2
3	30-39	33	28.7%	75.8
4	10-49	13	11.3%	32.2
5	50+	4	3.5%	n/a

Affected Subpopulations Maleh

Black Male 20-29	26	22.8%
Black Male 30-39	13	11.4%

Parish^g

Caldwell	3	2.6%	n/a
East Carroll	2	1.7%	n/a
Franklin	4	3.5%	n/a
Jackson	5	4.3%	33.6
Lincoln	12	10.4%	24.9
Madison	5	4.3%	51.0
Morehouse	8	7.0%	32.0
Ouachita	70	60.9%	44.1
Richland	4	3.5%	n/a
Tensas	0	0.0%	0.0
Union	2	1.7%	n/a
West Carroll	0	0.0%	0.0

<sup>Row diagnoses include only individuals whose initial primary and secondary syphillis diagnosis was made in Louisiana.
Cases with unknown/missing values for a demographic variable have been excluded from the counts and percentages for the demographic category. Total number of new diagnoses includes all cases, regardless of demographic variable completeness.
Cases per 100,000 population. Rates not available (n/a) for case counts less than 5. Rates not available for "Other" race group.
Race/ethnicity "Other" category does not include missing or unknown race/ethnicity.
Cases are presented by current gender identity rather than sex at birth.
Age when first diagnosed with primary and secondary syphillis is used for new primary and secondary syphillis diagnoses. Age categories align with 2021 census age groups.</sup>

Page with his diagnosed with primary and secondary syphillis is used for new primary and secondary syphillis diagnoses. Age categories and with 2021 census age groups.

Parish where first diagnosed with primary and secondary syphillis is used for new primary and secondary syphillis diagnoses. Persons without a parish of residence are not included in the total percentage.

Pall subpopulations under 5 were excluded.

Reported Cases of Early Non-Primary Non-Secondary Syphilis Region 8 - Monroe, 2022

	New Diagnoses ^a						
		Cases ^b	%	Ratec			_
To	otal	70	100.0%	20.1			
Race/Ethn	icity ^d				Affected Subpopulations Female	h Casesb	%
BI	ack	51	72.9%	39.3	Black Female 20-29	9	12.9%
Hi	spanic/Latinx	0	0.0%	0.0			
W	hite	17	24.3%	8.5			
Ot	ther	2	2.9%	n/a			
Gender ^e							
Fe	emale	25	35.7%	14.1			
Ma	ale	44	62.9%	25.8			
Tr	ansgender Male to Female	1	1.4%	n/a			
Agef					Affected Subpopulations Maleh		
10)-19	5	7.1%	10.4	Black Male 20-29	10	14.3%
20)-29	27	38.6%	54.1	Black Male 30-39	14	20.0%
30)-39	21	30.0%	48.2	Black Male 50+	5	7.1%
40)-49	7	10.0%	17.3			
50)+	10	14.3%	8.2			
Parish							
Ca	aldwell	2	2.9%	n/a			
Ea	ast Carroll	0	0.0%	0.0			
Fr	anklin	3	4.3%	n/a			
Ja	ackson	0	0.0%	0.0			
Lir	ncoln	3	4.3%	n/a			
Ma	adison	5	7.1%	51.0			
Me	orehouse	3	4.3%	n/a			
O	uachita	46	65.7%	29.0			
Ri	chland	6	8.6%	30.3			
Te	ensas	0	0.0%	0.0			
Ur	nion	1	1.4%	n/a			
W	est Carroll	1	1.4%	n/a			

a New diagnoses include only individuals whose initial non-primary non-secondary syphillis diagnosis was made in Louisiana.
b Cases with unknown/missing values for a demographic variable have been excluded from the counts and percentages for the demographic category. Total number of new diagnoses includes all cases, regardless of demographic variable completeness.
c Cases per 100,000 population. Rates not available (na) for case counts less than 5. Rates not available for "Other" race group.
Race/ethnicity "Other" category does not include missing or unknown race/ethnicity.
c Cases are presented by current gender identity rather than sex at birth.
Age when first diagnosed with non-primary non-secondary syphillis is used for new non-primary non-secondary syphillis diagnoses. Age categories align with 2021 census age groups.
Parish where first diagnosed with non-primary non-secondary syphillis is used for new non-primary non-secondary syphillis diagnoses. Persons without a parish of residence are not included in the total percentage.
All subpopulations under 5 were excluded.

Reported Cases of Congenital Syphilis Louisiana, 2022

Region specific table not available due to low case counts.

Infants and Mothers Linked to New Congenital Syphilis Cases^a

	intants and	wotners	Linkea to
	Cases	%	Rateb
Total Infant CS Cases	115	100.0%	204.0
Total Mothers Linked to New Cases	114	100.0%	4.8
Maternal Race/Ethnicity			
Black	71	62.3%	9.0
Hispanic/Latinx	6	5.3%	4.9
White	29	24.6%	2.1
Other	9	7.9%	n/a
Maternal Age			
15-19	7	6.1%	4.8
20-24	22	19.3%	14.8
25-29	37	32.5%	24.5
30-34	25	21.9%	15.3
35+	23	20.2%	1.8
Maternal Region of Residenced			
Alexandria - Region 6	11	31.0%	7.7
Baton Rouge - Region 2	21	18.4%	5.9
Hammond - Region 9	6	5.3%	1.9
Houma - Region 3	5	4.4%	2.5
Lafayette - Region 4	17	14.9%	5.6
Lake Charles - Region 5	8	7.0%	5.3
Monroe - Region 8	15	13.2%	8.5
New Orleans - Region 1	15	13.2%	3.3
Shreveport - Region 7	16	14.0%	6.0
Infant Region of Residenced			
Alexandria - Region 6	11	9.6%	293.2
Baton Rouge - Region 2	21	18.3%	246.2
Hammond - Region 9	6	5.2%	81.6
Houma - Region 3	5	4.3%	113.3
Lafayette - Region 4	17	14.8%	215.7
Lake Charles - Region 5	9	7.8%	237.8
Monroe - Region 8	15	13.0%	373.9
New Orleans - Region 1	15	13.0%	145.5
Shreveport - Region 7	16	13.9%	252.5
Maternal Stage of Syphilis			
Primary	2	1.8%	n/a
Secondary	8	7.0%	n/a
Early non-primary non-secondary	57	50.0%	n/a
Unknown/Late Latent	42	36.8%	n/a
Other	5	4.4%	n/a

Positive Toxicology Screense Cases					
Amphetamines/Meth	18				
Cocaine	1				
Heroin	4				
Marijuana	22				

 ^a New cases only include those who indicated a Louisiana residence at the time of delivery.
 ^b Congenital syphilis rates are calculated per 100,000 live births. Maternal rates are calculated using yearly census data, per 100,000 population. Percentages for maternal variables reflect distinct count of mothers. Rates for case counts under 20 are unreliable. Rates for case counts under 5 are not available.
 ^c Reflects maternal age at delivery.
 ^d Reflects maternal and infant region of residence at delivery.

e Reflects substances detected during maternal toxicology screens conducted during pregnancy or delivery. Mothers can have positive toxicology screens for more than one substance.

Reported Cases of Gonorrhea Region 8 - Monroe, 2022

New	Diagnoses ^a

Ratec

%

Casesb

	Total	1,298	100.0%	373.4	
Race/E	Ethnicity ^d				Affect
	Black	1,031	80.8%	794.6	
	Hispanic/Latinx	17	1.3%	181.3	
	White	187	14.7%	93.3	
	Other	41	3.2%	n/a	
Sex at	Birth				
	Female	642	49.5%	362.1	
	Male	656	50.5%	385.1	
Agee					
	Under 10	0	0.0%	0.0	
	10-19	422	32.5%	874.1	
	20-29	597	46.0%	1,196.2	
	30-39	185	14.3%	424.8	
	40-49	65	5.0%	161.0	
	50+	29	2.2%	23.7	
Paris	h ^f				
	Caldwell	14	1.1%	146.3	Affec
	East Carroll	39	3.0%	540.2	Alle
	Franklin	61	4.7%	310.1	
	Jackson	57	4.4%	383.2	
	Lincoln	180	13.9%	373.8	
	Madison	45	3.5%	459.2	
	Morehouse	105	8.1%	419.6	
	Ouachita	632	48.7%	398.1	
	Richland	79	6.1%	398.9	
	Tensas	9	0.7%	222.6	
	Union	63	4.9%	298.7	
	West Carroll	14	1.1%	145.9	

ected Subpopulations Female ^g	Casesb	%
Black Female 10-19	189	14.8%
Black Female 20-29	229	17.9%
Black Female 30-39	39	3.1%
Black Female 40-49	12	0.9%
Black Female 50+	7	0.5%
Other Female 10-19	7	0.5%
Other Female 20-29	13	1.0%
White Female 10-19	23	1.8%
White Female 20-29	47	3.7%
White Female 30-39	37	2.9%
White Female 40-49	12	0.9%
White Female 50+	6	0.5%

cted Subpopulations Maleg

Black Male 10-19	182	14.3%
Black Male 20-29	266	20.8%
Black Male 30-39	72	5.6%
Black Male 40-49	26	2.0%
Black Male 50+	9	0.7%
Other Male 20-29	6	0.5%
Other Male 30-39	8	0.6%
White Male 10-19	5	0.4%
White Male 20-29	20	1.6%
White Male 30-39	20	1.6%
White Male 40-49	12	0.9%
White Male 50+	5	0.4%

<sup>a New diagnoses include only individuals whose initial gonorrhea diagnosis was made in Louisiana.
b Cases with unknown/missing values for a demographic variable have been excluded from the counts and percentages for the demographic category. Total number of new diagnoses includes all cases, regardless of demographic variable completeness.
c Cases per 100,000 population. Rates not available (n/a) for case counts less than 5. Rates not available for "Other" race group.
d Race/ethnicity "Other" category does not include missing or unknown race/ethnicity.
d Age when first diagnosed with gonorrhea is used for new gonorrhea diagnoses. Age categories align with 2021 census age groups.
I Parish where first diagnosed with gonorrhea is used for new gonorrhea diagnoses. Persons without a parish of residence are not included in the total percentage.

Affected subpopulations with counts less than 5 are excluded.</sup>

Reported Cases of Chlamydia Region 8 - Monroe, 2022

			New Diagnose	
	Casesb	%	Rate	
Total	2 932	100.0%	843.5	

Race/Ethnicity^d 1,771.1 Black 2,298 80.4% 298.6 Hispanic/Latinx 28 1.0% White 218.6 438 15.3% Other 94 3.3% n/a

Sex at Birth				
Female	1,960	66.8%	1,105.6	
Male	972	33.2%	570.6	

	Maio	012	00.270	0.0.0
Age				
	Under 10	1	0.0%	n/a
	10-19	1,124	38.3%	2,328.2
	20-29	1,419	48.4%	2,843.3
	30-39	298	10.2%	684.2
	40-49	61	2.1%	151.1

Parish				
	Caldwell	37	1.3%	386.6
	East Carroll	65	2.2%	900.3
	Franklin	154	5.3%	783.0

Franklin	154	5.3%	783.0
Jackson	84	2.9%	564.7
Lincoln	435	14.8%	903.4
Madison	149	5.1%	1,520.6
Morehouse	251	8.6%	1,003.0
Ouachita	1,342	45.8%	845.3
Richland	190	6.5%	959.4
Tensas	24	0.8%	593.6
Union	163	5.6%	772.8

38

29

1.0%

23.7

396.1

1.3%

Affected Subpopulations Femaleg	Casesb	%
Black Female 10-19	607	21.2%
Black Female 20-29	750	26.2%
Black Female 30-39	125	4.4%
Black Female 40-49	19	0.7%
Black Female 50+	12	0.4%
Hispanic Female 10-19	5	0.2%
Hispanic Female 20-29	9	0.3%
Hispanic Female 30-39	5	0.2%
Other Female 10-19	25	0.9%
Other Female 20-29	34	1.2%
White Female 10-19	120	4.2%
White Female 20-29	152	5.3%
White Female 30-39	38	1.3%

Affected Subpopulations Maleg

White Female 40-49

Black Male 10-19	304	10.6%
Black Male 20-29	360	12.6%
Black Male 30-39	92	3.2%
Black Male 40-49	19	0.7%
Black Male 50+	9	0.3%
Other Male 10-19	7	0.2%
Other Male 20-29	13	0.5%
Other Male 30-39	8	0.3%
White Male 10-19	27	0.9%
White Male 20-29	58	2.0%
White Male 30-39	18	0.6%
White Male 40-49	8	0.3%
White Male 50+	6	0.2%

0.4%

11

West Carroll

50+

<sup>a New diagnoses include only individuals whose initial chlamydia diagnosis was made in Louisiana.
b Cases with unknown/missing values for a demographic variable have been excluded from the counts and percentages for the demographic category. Total number of new diagnoses includes all cases, regardless of demographic variable completeness.
c Cases per 100,000 population. Rates not available (n/a) for case counts less than 5. Rates not available for "Other" race group.
d Race/ethnicity "Other" category does not include missing or unknown race/ethnicity.
e Age when first diagnosed with chlamydia is used for new chlamydia diagnoses. Age categories align with 2021 census age groups.</sup>

Age when this troughosed with chiarmydia is used for new chiarmydia diagnoses. Age categories angli with 2021 census age groups.
 Parish where first diagnosed with chlamydia is used for new chlamydia diagnoses. Persons without a parish of residence are not included in the total percentage.
 Affected subpopulations with counts less than 5 are excluded.

Reported Cases of Acute Hepatitis B Louisiana, 2022

Region specific table not available due to low case counts.

New Diagnoses^a

	Cases ^b	%	Ratec			
Total	29	100.0%	0.6			
ce/Ethnicity ^d				Affected Subpopulations Femaleg	Casesb	%
Asian	0	0.0%	0.0	White Female	9	33.3%
Black	6	22.2%	0.4			
Hispanic/ Latinx	0	0.0%	0.0			
White	21	77.8%	0.7			
x at Birth						
Female	15	51.7%	0.6	•		
Male	14	48.3%	0.6			
le _e						
18-29	1	3.4%	n/a			
30-39	4	13.8%	n/a			
40-49	10	34.5%	1.8			
50-59	9	31.0%	1.5			
60-69	3	10.3%	n/a			
70+	2	6.9%	n/a			
gion ^f				Affected Subpopulations Male		
Region 1 - New Orleans	2	6.9%	n/a	White Male	12	44.49
Region 2 - Baton Rouge	4	13.8%	n/a			
Region 3 - Houma	6	20.7%	1.5			
Region 4 - Lafayette	2	6.9%	n/a			
Region 5 - Lake Charles	5	17.2%	1.6			
Region 6 - Alexandria	5	17.2%	1.7			

0

0

5

0.0%

0.0%

17.2%

0.0 0.0

8.0

Region 7 - Shreveport

Region 9 - Hammond/Slidell

Region 8 - Monroe



a New diagnoses include only individuals whose initial acute hepatitis B diagnosis was made in Louisiana.
 b Cases with unknown/missing values for a demographic variable have been excluded from the counts and percentages for the demographic category. Total number of new diagnoses includes all cases, regardless of demographic variable completeness.
 c Cases per 100,000 population. Rates not available (n/a) for case counts less than 5. Rates not available for "Other" race

Cases per 100,000 population. Rates not available (n/a) for case counts less than 5. Rates not available for "Other" race groups.
 Race/ethnicity "Other" category does not include missing or unknown race/ethnicity.
 Age when first diagnosed with acute hepatitis B is used for new diagnoses. Age categories align with 2021 census age groups.
 Region where first diagnosed with acute hepatitis B is used for new diagnoses. Persons without a region of residence are not included in the total percentage.
 Affected subpopulations with counts less than 5 are excluded.

Reported Cases of Chronic Hepatitis B Region 8 - Monroe, 2022

New Diagnosesa % Casesb Ratec Total 57 100.0% 16.4 Casesb % Race/Ethnicity^d Affected Subpopulations Female⁹ 10.2% 5 2 4.3% n/a Black Male 30-39 Asian 28 21.5 5 10.2% Black 59.6% Black Male 60-69 Hispanic/Latinx 3 6.4% n/a White 14 29.8% 6.9 Sex at Birth 19 Female 33.3% 10.7 Male 38 66.7% 22.3 Agee 0-17 0 0.0% 0.0 18-29 12 21.1% 19.8 12 21.1% 27.6 30-39 10 17.5% 24.8 40-49 50-59 9 15.8% 21.9 60-69 6 10.5% 14.5 8 70+ 14.0% 20.2 **Parishf** Affected Subpopulations Maleg Caldwell 0 0.0% 0.0 Black Male 30-39 5 10.2% East Carroll 3 5.3% n/a Black Male 60-69 5 10.2% 2 n/a Franklin 3.5% 33.6 5 Jackson 8.8% Jefferson 0 0.0% 0.0 9 18.6 Lincoln 15.8% Madison 0 0.0% 0.0 Morehouse 1 1.8% n/a Ouachita 56.1% 20.1 32 Richland 1.8% n/a 1 Tensas 2 n/a 3.5% Union 2 3.5% n/a

0.0

0

0.0%

West Carroll



New diagnoses include only individuals whose initial chronic hepatitis B diagnosis was made in Louisiana.
 Cases with unknown/missing values for a demographic variable have been excluded from the counts and percentages for the demographic category. Total number of new diagnoses includes all cases, regardless of demographic variable

c Cases per 100,000 population. Rates not available (n/a) for case counts less than 5. Rates not available for "Other" race

groups.

d Race/ethnicity "Other" category does not include missing or unknown race/ethnicity.

e Age when first diagnosed with chronic hepatitis B is used for new diagnoses. Age categories align with 2021 census age

groups.

Farish where first diagnosed with chronic hepatitis B is used for new diagnoses. Persons without a parish of residence are not included in the total percentage.

Affected subpopulations with counts less than 5 are excluded.

Reported Cases of Acute Hepatitis C Region 8 - Monroe, 2022

New	Diagi	noses
-----	-------	-------

		Casesb	%	Ratec
	Total	8	100.0%	2.3
Race/E	Ethnicity ^d			
	Asian	0	0.0%	0.0
	Black	3	50.0%	n/a
	Hispanic/ Latinx	1	16.7%	n/a
	White	2	33.3%	n/a
Sex at	Birth			
	Female	4	50.0%	n/a
	Male	4	50.0%	n/a
Agee				
	0-17	0	0.0%	0.0
	18-29	2	25.0%	n/a
	30-39	4	50.0%	n/a
	40-49	0	0.0%	0.0
	50-59	1	12.5%	n/a
	60-69	1	12.5%	n/a
	70+	0	0.0%	0.0

Affected Subpopulations Female⁹ Cases^b %

All female subpopulations had a case count less than 5.

Parishf

Caldwell	0	0.0%	0.0
East Carroll	0	0.0%	0.0
Franklin	0	0.0%	0.0
Jackson	1	16.7%	n/a
Lincoln	1	16.7%	n/a
Madison	0	0.0%	0.0
Ouachita	3	50.0%	n/a
Richland	0	0.0%	0.0
Tensas	0	0.0%	0.0
Union	1	16.7%	n/a
West Carroll	0	0.0%	0.0

Affected Subpopulations Maleg

All male subpopulations had a case count less than 5.



a New diagnoses include only individuals whose initial acute hepatitis C diagnosis was made in Louisiana.
 b Cases with unknown/missing values for a demographic variable have been excluded from the counts and percentages for the demographic category. Total number of new diagnoses includes all cases, regardless of demographic variable completeness.
 c Cases per 100,000 population. Rates not available (n/a) for case counts less than 5. Rates not available for "Other" race

Cases per 100,000 population. Rates not available (n/a) for case counts less than 5. Rates not available for "Other" race groups.

Race/ethnicity "Other" category does not include missing or unknown race/ethnicity.

Age when first diagnosed with acute hepatitis C is used for new diagnoses. Age categories align with 2021 census age groups.

Parish where first diagnosed with acute hepatitis C is used for new diagnoses. Persons without a parish of residence are not

included in the total percentage.

9 Affected subpopulations with counts less than 5 are excluded.

Reported Cases of Chronic Hepatitis C Region 8 - Monroe, 2022

New Diagnosesa

70.8

	Casesb	%	Ratec			
Total	232	100.0%	66.7	· -		
Race/Ethnicity ^d				Affected Subpopulations Female	Casesb	%
Asian	3	1.3%	n/a	Black Female 60-69	8	3.5%
Black	65	29.1%	50.0	White Female 18-29	5	2.2%
Hispanic/ Latinx	13	5.8%	138.6	White Female 30-39	12	5.2%

White Female 40-49 White Female 50-59

White Female 60-69

11

11

7

4.8%

4.8%

3.1%

Sex at Birth				
Female	79	34.1%	44.5	
Male	153	65.9%	89.8	

Agee

White

0-17	0	0.0%	0.0
18-29	18	7.8%	29.6
30-39	47	20.3%	107.9
40-49	49	21.1%	121.3
50-59	47	20.3%	114.4
60-69	56	24.1%	134.9
70+	15	6.5%	37.8

Parishf Affected

63.7%

142

Caldwell	9	4.0%	94.0	_
East Carroll	10	4.4%	138.5	
Franklin	11	4.9%	55.9	
Jackson	14	6.2%	94.1	
Lincoln	16	7.1%	33.2	
Madison	8	3.6%	81.6	
Morehouse	17	7.6%	67.9	
Ouachita	110	48.9%	69.2	
Richland	20	8.9%	100.9	
Tensas	2	0.9%	n/a	
West Carroll	8	3.6%	83.3	
				_

Subpopulations Maleg		
Black Male 30-39	7	3.1%
Black Male 50-59	10	4.4%
Black Male 60-69	22	9.6%
Black Male 70+	5	2.2%
White Male 18-29	9	3.9%
White Male 30-39	21	9.2%
White Male 40-49	28	12.2%
White Male 50-59	17	7.4%
White Male 60-69	14	6.1%



a New diagnoses include only individuals whose initial chronic hepatitis C diagnosis was made in Louisiana.
 b Cases with unknown/missing values for a demographic variable have been excluded from the counts and percentages for the demographic category. Total number of new diagnoses includes all cases, regardless of demographic variable completeness.
 c Cases per 100,000 population. Rates not available (n/a) for case counts less than 5. Rates not available for "Other" race

Cases per 100,000 population. Rates not available (n/a) for case counts less than 5. Rates not available for "Other" race groups.

d Race/ethnicity "Other" category does not include missing or unknown race/ethnicity.
de Age when first diagnosed with chronic hepatitis C is used for new diagnoses. Age categories align with 2021 census age groups.

Parisk where first diagnosed with chronic hepatitis C is used for new diagnoses. Persons without a parish of residence are

not included in the total percentage.

9 Affected subpopulations with counts less than 5 are excluded