

# Louisiana Department of Health and Hospitals Office of Public Health

STD/HIV Program

### DeAnn Gruber, PhD, LCSW

Administrative Director

### Debbie Wendell, PhD, MPH

Data Management/Analysis Manager

### Jessica Fridge, MSPH

Surveillance Program Manager

### Stephanie Taylor, MD

STD Medical Director

#### **Amy Busby**

Regional Operations Manager

### Joy Ewell

CDC Lead PHA-Louisiana

#### Editor/Production:

### Demerial Banks, MPH Catherine Desmarais, MPH

#### Data Management/Analysis and Surveillance:

Rashad Arcement, MSPH, Incidence & Resistance Coordinator

Demerial Banks, MPH, HIV Surveillance Epidemiologist

Tania Bechtel, HIV Data Manager

Antoine Brantley, MPH, Data Analyst

Catherine Desmarais, MPH, STD Surveillance Epidemiologist and Data Manager

Mohammad Rahman, MBBS, MPH, PhD, STD Epidemiologist Supervisor

William T. Robinson, Ph.D., Biostatistician

Denise Robertson, ICCR Coordinator

Javone Davis, Field Operations Supervisor

Josette Russell, HIV Data Entry Specialist

Lakeesha Brumfield, HIV Data Entry Specialist

Carmen Washington, HIV Data Manager

#### **HIV Field Epidemiologists:**

#### Betsy Shepard, PAC, MPH, Epidemiologist Supervisor

James Hubbard	Region 1	Cheryl Edney, BS MT(ASCP)	<b>Regions 4, 5, 6</b>
Monica Pendelton	Regions 1, 3	Gayle Norris, MA	Regions 7, 8
		Gale Terry, LPN	Regions 2, 9

#### Regional STD Supervisors:

Michael Carter	Region 1	Patrick Williams	Region 5
Megan King	Region 2	Amy Busby	Region 6
<b>Javone Davis</b>	Regions 3, 8	Charles Jones	Region 7
Ken May	Region 4	John Lyons	Region 9

# TABLE OF CONTENTS

Contents	Page
Surveillance Summary	1
Louisiana: Persons Living with HIV Infection	3
Persons newly diagnosed with HIV and AIDS	4
Persons diagnosed with Early Syphilis	5
Parish: Persons Living with HIV Infection	6
Persons diagnosed with Early Syphilis	7
Metro New Orleans: Persons Living with HIV Infection	8
Persons newly diagnosed with HIV and AIDS	9
Persons diagnosed with Early Syphilis	10
Metro Baton Rouge: Persons Living with HIV Infection	11
Persons newly diagnosed with HIV and AIDS	12
Persons diagnosed with Early Syphilis	13
Region 1: Persons Living with HIV Infection	14
Persons newly diagnosed with HIV and AIDS	15
Persons diagnosed with Early Syphilis	16
Region 2: Persons Living with HIV Infection	17
Persons newly diagnosed with HIV and AIDS	18
Persons diagnosed with Early Syphilis	19
Region 3: Persons Living with HIV Infection	20
Persons newly diagnosed with HIV and AIDS	21
Persons diagnosed with Early Syphilis	22
Region 4: Persons Living with HIV Infection	23
Persons newly diagnosed with HIV and AIDS	24
Persons diagnosed with Early Syphilis	25
Region 5: Persons Living with HIV Infection	26
Persons newly diagnosed with HIV and AIDS	27
Persons diagnosed with Early Syphilis	28
Region 6: Persons Living with HIV Infection	29
Persons newly diagnosed with HIV and AIDS	30
Persons diagnosed with Early Syphilis	31
Region 7: Persons Living with HIV Infection	32
Persons newly diagnosed with HIV and AIDS	33
Persons diagnosed with Early Syphilis	34
Region 8: Persons Living with HIV Infection	35
Persons newly diagnosed with HIV and AIDS	36
Persons diagnosed with Early Syphilis	37
Region 9: Persons Living with HIV Infection	38
Persons newly diagnosed with HIV and AIDS	39
Persons diagnosed with Early Syphilis	40
List of Figures	4.1
Louisiana: 10-Year Trends New HIV and New AIDS Diagnoses (2005-2014)	41
New HIV Diagnoses	
New AIDS Diagnoses	10
Louisiana: 10-Year Trends P&S Syphilis Diagnoses (2005-2014)	42
Louisiana: 5-Year Trends Early Latent Syphilis Diagnoses (2010-2014)	42
Louisiana: 10-Year Persons Living with HIV Infection (2005-2014)	43
2014 HIV & Syphilis: Region 1- New Orleans	

# **Surveillance Summary**

### Data in this report includes cases diagnosed with HIV, AIDS or Early Syphilis

- In 2013, Louisiana ranked **3<sup>rd</sup>** in the nation for estimated HIV case rates (30.3 per 100,000 population) and 11<sup>th</sup> in the estimated number of HIV cases. The New Orleans MSA ranked **2<sup>nd</sup>** and the Baton Rouge MSA ranked **4<sup>th</sup>** for estimated HIV case rates (43.4 and 38.1 per 100,000, respectively), among the large metropolitan areas in the nation. The most recent *CDC HIV Surveillance Report* (vol. 25) was released in February 2015 and provided the 2013 rankings.
- According to the same report, Louisiana ranked 3<sup>rd</sup> highest in estimated state AIDS case rates (16.9 per 100,000) and 11<sup>th</sup> in the number of estimated AIDS cases in 2013. The Baton Rouge metropolitan area ranked 3<sup>rd</sup> in estimated AIDS case rates (25.4 per 100,000) and the New Orleans metropolitan area ranked 5<sup>th</sup> in estimated AIDS case rates (33.0 per 100,000) in 2013 among the large metropolitan areas in the nation.
- In 2013, Louisiana ranked 3<sup>rd</sup> in the nation for primary and secondary (P&S) syphilis case rates (9.1 per 100,000 population) according to the 2013 *CDC STD Surveillance Report*.
- As of March 31, 2015, a total of **20,112** persons are living with HIV infection (PLWH) in Louisiana; of these individuals, **10,666** persons (53%) have an AIDS diagnosis.
- In 2014, **1,422** new HIV cases were diagnosed in Louisiana. Of these cases, **30%** (n=430) occurred in Region 1 (New Orleans) and **24%** (n=344) occurred in Region 2 (Baton Rouge).
- In the first quarter of 2015 (January March), a total of **177** persons were diagnosed with early syphilis\* in Louisiana; **100** persons were diagnosed with P&S syphilis, and **77** persons were diagnosed with early latent syphilis. Of these cases, **21%** (n=37) occurred in Region 1 (New Orleans) and **18%** (n=32) each occurred in Region 2 (Baton Rouge) and Region 7 (Shreveport).
- The primary risk factor for new HIV diagnoses in Louisiana is men who have sex with men (MSM). In 2014, of the persons who reported a risk factor, 73% reported being MSM, and an additional 2% reported being MSM and an injection drug user (IDU). As of March 31, 2015, 54% of all people living with HIV infection in Louisiana were MSM.
- The primary risk factor for early syphilis diagnoses in Louisiana is MSM. In the first quarter of 2015, of the persons who reported a risk factor, 43% reported being MSM, and 41% reported engaging in heterosexual activity. *Persons could report multiple risk factors for syphilis diagnoses*.
- In Louisiana, 25% of new HIV cases, 31% of new AIDS cases, and 31% of the early syphilis diagnoses were among women.
- African Americans in Louisiana are experiencing a severe health disparity in regards to HIV and STDs. In 2014, 70% of newly diagnosed HIV cases and 74% of newly diagnosed AIDS cases were among African Americans. In the first quarter of 2015, 81% of early syphilis cases were among African Americans. African Americans make up only 32% of Louisiana's overall population.
- A large percentage of new HIV and syphilis diagnoses occurred among adolescents and young adults. In 2014, 25% of all new HIV diagnoses occurred among persons 13-24 years old and an additional 33% occurred among persons 25-34 years old. In the first quarter of 2015, 48% of all early syphilis diagnoses occurred among persons 13-24 years old and an additional 34% occurred among persons 25-34 years old.

<sup>\*</sup> A syphilis diagnosis with less than one year duration is considered to be "early syphilis." This designation covers the primary, secondary, and early latent stages of syphilis.

### **New Quarterly Report Format:**

SHP's new quarterly report format includes P&S syphilis and early latent syphilis diagnoses for Louisiana, the New Orleans and Baton Rouge MSAs, and all nine of the public health regions. It is important to note that the time periods for the HIV and syphilis data are not equivalent. Please pay special attention to the titles above each table. There are different reporting delays and methods for HIV and syphilis which impact the timeliness of those data in this report. The 'Persons Living with AIDS' excerpts have been removed.

### AIDS Case Definition Change – Effective for all 2014 and future AIDS Diagnoses:

Prior to 2014, the surveillance case definition for an AIDS diagnosis was: a CD4 cell count <200, a CD4 percentage <14%, or the diagnosis of an opportunistic infection (OI) such as Kaposi Sarcoma. The CDC has changed the case surveillance definition for AIDS. For all AIDS diagnoses, starting January 1, 2014, the case definition is: a CD4 cell count <200 or the diagnosis of an OI. A CD4 percent <14% is no longer part of the AIDS case definition, unless the CD4 count is not available. Previously, a person with a CD4 count of 300 but a percent of 10%, would have been defined as an AIDS case. Because that CD4 count is not below 200, that case is no longer considered a new AIDS diagnosis. If the CD4 count of 300 was not available and only the percent of 10% was available, then the case would be newly diagnosed as AIDS. This change in surveillance case definition will reduce the number of AIDS diagnoses that occur each year, starting with 2014 data.

### **Transgender Data:**

SHP will be incorporating more transgender and gender identity data into the quarterly report. For this first quarter, the addition of transgender women and transgender men was incorporated into the Gender breakdown for PLWH and new diagnoses of HIV and AIDS. In subsequent reports we will add transgender data to the early syphilis reports and account for transgender identity in the race and exposure category tables.

### Time Periods for this Quarterly Report (First Quarter 2015 Report):

Persons Living with HIV Infection\* – As of March 31, 2015

New HIV Diagnoses\* – Calendar Year 2014

New AIDS Diagnoses\* - Calendar Year 2014

New P&S Syphilis Diagnoses – January to March 31, 2015

New Early Latent Syphilis Diagnoses – January to March 31, 2015

\*Transgender diagnoses have been reported for Persons Living with HIV infection, New HIV Diagnoses, New AIDS Diagnoses

### **Future Quarterly Report:**

Second Quarter 2015 Report: Release in August 2015

Persons Living with HIV Infection, As of June 30, 2015

HIV and AIDS Diagnoses, Calendar Year 2014 (Updated)

P&S Syphilis and Early Latent Syphilis Diagnoses, First and Second Quarter 2015 (January-June 2015)

Third Quarter 2015 Report: Release in November 2015

Persons Living with HIV Infection, As of September 30, 2015

HIV and AIDS Diagnoses, Calendar Year 2014 (Updated)

P&S Syphilis and Early Latent Syphilis Diagnoses, First-Third Quarter 2015 (January-September 2015)

Fourth Quarter 2015 Report: Release in February 2016

Persons Living with HIV Infection, As of December 31, 2015

HIV and AIDS Diagnoses, Calendar Year 2014 (Updated)

P&S Syphilis and Early Latent Syphilis Diagnoses, Calendar Year 2015

Please note: This report reflects only information reported to Louisiana's HIV/AIDS and STD surveillance systems.

### Persons Living with HIV Infection Louisiana as of March 31, 2015

- **35,922** cumulative cases have been detected in Louisiana, including **351** pediatric cases.
- 15,575 cumulative deaths among persons with HIV infection have occurred in Louisiana, including 121 pediatric deaths.
- 20,112 persons are currently living with HIV infection in Louisiana.

Persons living with HIV: 9,446
Persons living with AIDS: 10,666

### Gender

	Cases	Percent
Women	5,883	29 %
Men	14,064	70 %
Transgender Women	163	1 %
Transgender Men	2	<1 %
Total	20,112	100 %

Age Category			Race/Ethnicity by Birth Sex				
<b>Current Age</b>	Cases	Percent		Men	Women	Total	Percent
0 - 12	52	<1 %	Black, Non-Hispanic	8,902	4,846	13,748	68 %
13 - 19	133	1 %	White, Non-Hispanic	4,475	819	5,294	26 %
20 - 24	815	4 %	Hispanic	655	156	811	4 %
25 - 29	1,763	9 %	Asian/Pacific Islander	68	16	84	<1 %
30 - 34	2,098	10 %	American Indian	28	9	37	<1 %
35 - 39	2,408	12 %	Unknown	5	4	9	<1 %
40 - 44	2,336	12 %	Multi-race	94	35	129	1 %
45 - 49	2,695	13 %	Total	14,227	5,885	20,112	100 %
50 - 54	3,109	15 %					
55 - 59	2,339	12 %					
60+	2,364	12 %					
Total	20,112	100 %					

### **Exposure Category by Birth Sex**

Cases with a reported risk:	Men	Women	Total	Percent
Men who have sex with men	7,253	0	7,253	54 %
Injection drug use	1,051	639	1,690	13 %
Men who have sex with men and inject drugs	829	0	829	6 %
Heterosexual contact	1,096	2,355	3,451	26 %
Transfusion/Hemophilia	40	31	71	1 %
Confirmed Other	0	1	1	<1 %
Perinatal	95	109	204	2 %
Total	10,364	3,135	13,499	100 %
No Identified Risk	3,863	2,750	6,613	

# Persons Newly Diagnosed with HIV and AIDS Louisiana

### **HIV CASES DIAGNOSED January - December 2014**

• 1,422 persons were newly diagnosed with HIV in Louisiana.

Gender	Cases	Percent	Race/Ethnicity	Cases	Percent
Men	1,051	74 %	Black, Non-Hispanic	990	70 %
Women	357	25 %	White, Non-Hispanic	317	22 %
Transgender V	Women 14	1 %	Hispanic	86	6 %
Total	1,422	100 %	Asian/Pacific Islander	14	1 %
			American Indian	4	<1 %
			Unknown	1	<1 %
			Multi-race	10	1 %
			Total	1,422	100 %

Age at Diagnosis	Cases	Percent	Cases with a reported risk (by birth sex):	Cases	Percent
0 - 12	2	<1 %	Men who have sex with men	576	73 %
13 - 24	352	25 %	Injection drug use	47	6 %
25 - 34	463	33 %	Men who have sex with men and inject drugs	18	2 %
35 - 44	285	20 %	Heterosexual contact	148	19 %
45+	320	23 %	Perinatal	4	1 %
Total	1,422	100 %	Total	793	100 %

No Identified Risk 629

# AIDS CASES DIAGNOSED January - December 2014

• 617 persons were newly diagnosed with AIDS in Louisiana.

Gender	Cases	Percent	Race/Ethnicity	Cases	Percent
Men	418	68 %	Black, Non-Hispanic	458	74 %
Women	192	31 %	White, Non-Hispanic	116	19 %
Transgender Women	7	1 %	Hispanic	32	5 %
Total	617	100 %	Asian/Pacific Islander	3	<1 %
			American Indian	1	<1 %
			Unknown	1	<1 %
			Multi-race	6	1 %
			Total	617	100 %

Age at Diagnosis	Cases	Percent	Cases with a reported risk (by birth sex):	Cases	Percent
13 - 24	47	8 %	Men who have sex with men	188	56 %
25 - 34	176	29 %	Injection drug use	43	13 %
35 - 44	164	27 %	Men who have sex with men and inject drugs	6	2 %
45+	230	37 %	Heterosexual contact	97	29 %
Total	617	100 %	Transfusion/Hemophilia	1	<1 %
			Total	335	100 %
			No Identified Risk	282	

# Persons Diagnosed with Early Syphilis Louisiana

### PERSONS DIAGNOSED WITH P&S SYPHILIS January - March 2015

• 100 persons were diagnosed with P&S Syphilis in Louisiana between January 1 and March 31, 2015.

Sex at Birth	Cases	Percent	Race/Ethnicity	Cases	Percent
Female	28	28 %	American Indian	0	0 %
Male	72	72 %	Asian/Pacific Islander	0	0 %
Total	100	100 %	Black, Non-Hispanic	83	83 %
			Hispanic	0	0 %
			White, Non-Hispanic	17	17 %
			Other/Multi-race	0	0 %
			Unknown	0	0 %
			Total	100	100 %

Age at Diagnosis	Cases	Percent	Cases with a reported risk (Can have >1 risk):	Cases	Percent
0 - 12	0	0 %	Men who have sex with men and inject drugs	0	0 %
13 - 19	11	11 %	Men who have sex with men	46	48 %
20 - 29	62	62 %	Injection drug use	0	0 %
30 - 39	12	12 %	Transactional sex	5	5 %
40+	15	15 %	High risk heterosexual contact	19	20 %
Total	100	100 %	Heterosexual contact	37	39 %
			Incarceration	12	13 %
			Illegal drug use (non-injection)	31	32 %
			No Reported Risk	4	

### PERSONS DIAGNOSED WITH EARLY LATENT SYPHILIS January - March 2015

• 77 persons were diagnosed with Early Latent Syphilis in Louisiana between January 1 and March 31, 2015.

Sex at Birth	Cases	Percent	Race/Ethnicity	Cases	Percent
Female	27	35 %	American Indian	2	3 %
Male	50	65 %	Asian/Pacific Islander	0	0 %
Total	77	100 %	Black, Non-Hispanic	60	78 %
			Hispanic	2	3 %
			White, Non-Hispanic	13	17 %
			Other/Multi-race	0	0 %
			Unknown	0	0 %
			Total	77	100 %

Age at Diagnosis	Cases	Percent	Cases with a reported risk (Can have >1 risk):	Cases	Percent
0 - 12	0	0 %	Men who have sex with men and inject drugs	0	0 %
13 - 19	11	14 %	Men who have sex with men	29	38 %
20 - 29	38	49 %	Injection drug use	0	0 %
30 - 39	19	25 %	Transactional sex	1	1 %
40+	9	12 %	High risk heterosexual contact	15	19 %
Total	77	100 %	Heterosexual contact	34	44 %
			Incarceration	7	9 %
			Illegal drug use (non-injection)	19	25 %
			No Reported Risk	0	

<sup>\*</sup> A syphilis diagnosis with less than one year duration is considered to be "early syphilis." This designation covers the primary, secondary, and early latent stages of syphilis.

# Persons Living with HIV Infection by Parish as of March 31, 2015

# First Quarter 2015

<u>Parish</u>	Living Cases	<u>Parish</u>	Living Cases
Acadia	108	Sabine	21
Allen	262	St. Bernard	171
Ascension	233	St. Charles	109
Assumption	27	St. Helena	16
Avoyelles	184	St. James	76
Beauregard	46	St. John the Baptist	168
Bienville	31	St. Landry	274
Bossier	200	St. Martin	106
Caddo	1,083	St. Mary	81
Calcasieu	654	St. Tammany	424
Caldwell	18	Tangipahoa	369
Cameron	2	Tensas	33
Catahoula	28	Terrebonne	206
Claiborne	54	Union	34
Concordia	44	Vermilion	96
De Soto	54	Vernon	68
East Baton Rouge	3,884	Washington	184
East Carroll	28	Webster	81
East Feliciana	156	West Baton Rouge	95
Evangeline	73	West Carroll	10
Franklin	50	West Feliciana	173
Grant	45	Winn	46
Iberia	116		
Iberville	356		
Jackson	21		
Jefferson	1,859		
Jefferson Davis	61		
La Salle	31		
Lafayette	745		
Lafourche	143		
Lincoln	100		
Livingston	210		
Madison	32		
Morehouse	61		
Natchitoches	114		
Orleans	4,930		
Ouachita	616		
Plaquemines	38		
Pointe Coupee	48		
Rapides	465		
Red River	14		
Richland	47		

Total 20,112

# Persons Diagnosed with Early Syphilis\* by Parish January - March 2015

Parish_	Diagnoses	<b>Parish</b>	Diagnoses
Acadia	2	Sabine	0
Allen	1	St. Bernard	1
Ascension	0	St. Charles	0
Assumption	0	St. Helena	0
Avoyelles	1	St. James	0
Beauregard	0	St. John the Baptist	
Bienville	0	St. Landry	5
Bossier	6	St. Martin	2
Caddo	22	St. Mary	0
Calcasieu	5	St. Tammany	1
Caldwell	0	Tangipahoa	2
Cameron	0	Tensas	0
Catahoula	0	Terrebonne	12
Claiborne	0	Union	0
Concordia	0	Vermilion	4
De Soto	3	Vernon	0
East Baton Rouge	31	Washington	0
East Carroll	2	Webster	1
East Feliciana	0	West Baton Rouge	0
Evangeline	0	West Carroll	0
Franklin	0	West Feliciana	0
Grant	0	Winn	0
Iberia	5		
Iberville	1		
Jackson	0		
Jefferson	12		
Jefferson Davis	0		
La Salle	0		
Lafayette	8		
Lafourche	0		
Lincoln	4		
Livingston	0		
Madison	0		
Morehouse	2		
Natchitoches	0		
Orleans	24		
Ouachita	10		
Plaquemines	0		
Pointe Coupee	0		
Rapides	6		
Red River	0		
Richland	1		

Total 177

<sup>\*</sup> A syphilis diagnosis with less than one year duration is considered to be "early syphilis." This designation covers the primary, secondary, and early latent stages of syphilis.

### Persons Living with HIV Infection Metro New Orleans as of March 31, 2015

Jefferson, Orleans, St. Tammany, St. John the Baptist, Plaquemines, St. Bernard, St. James, St. Charles Parishes

- 16,409 cumulative cases have been detected in the New Orleans MSA, including 168 pediatric cases.
- 7,146 cumulative deaths among persons with HIV infection have occurred in the New Orleans MSA, including 60 pediatric deaths
- 7,775 persons are currently living with HIV infection in the New Orleans MSA.

Persons living with HIV: **3,610** Persons living with AIDS: **4,165** 

#### Gender

	Cases	Percent
Women	1,986	26 %
Men	5,692	73 %
Transgender Women	95	1%
Transgender Men	2	<1%
Total	7.775	100 %

Age Category			Race/Ethnicity by Birth S	Sex			
Current Age	Cases	Percent		Men	Women	Total	Percent
0 - 12	19	<1 %	Black, Non-Hispanic	3,203	1,615	4,818	62 %
13 - 19	49	1 %	White, Non-Hispanic	2,135	248	2,383	31 %
20 - 24	283	4 %	Hispanic	363	107	470	6 %
25 - 29	653	8 %	Asian/Pacific Islander	39	7	46	1 %
30 - 34	766	10 %	American Indian	9	0	9	<1 %
35 - 39	910	12 %	Unknown	4	2	6	<1 %
40 - 44	858	11 %	Multi-race	34	9	43	1 %
45 - 49	1,049	13 %	Total	5,787	1,988	7,775	100 %
50 - 54	1,221	16 %					
55 - 59	948	12 %					
60+	1,019	13 %					
Total	7,775	100 %					

Exposure	<u>Category</u>	by Birth Sex

Cases with a reported risk:	Men	Women	Total	Percent
Men who have sex with men	3,287	0	3,287	61 %
Injection drug use	318	215	533	10 %
Men who have sex with men and inject drugs	313	0	313	6 %
Heterosexual contact	357	797	1,154	21 %
Transfusion/Hemophilia	18	14	32	1 %
Perinatal	30	43	73	1 %
Total	4,323	1,069	5,392	100 %
No Identified Risk	1,464	919	2,383	

### Persons Newly Diagnosed with HIV and AIDS Metro New Orleans

Jefferson, Orleans, St. Tammany, St. John the Baptist, Plaquemines, St. Bernard, St. James, St. Charles Parishes

# **HIV CASES DIAGNOSED January - December 2014**

• 471 persons were newly diagnosed with HIV in the New Orleans MSA.

<u>Gender</u>	Cases	Percent	Race/Ethnicity	Cases	Percent
Men	380	81 %	Black, Non-Hispanic	301	64 %
Women	91	19 %	White, Non-Hispanic	117	25 %
Total	471	100 %	Hispanic	43	9 %
			Asian/Pacific Islander	6	1 %
			Unknown	1	<1 %
			Multi-race	3	1 %
			Total	471	100 %

Age at Diagnosis	Cases	Percent	Cases with a reported risk (by birth sex):	Cases	Percent
13 - 24	126	27 %	Men who have sex with men	246	79 %
25 - 34	149	32 %	Injection drug use	12	4 %
35 - 44	94	20 %	Men who have sex with men and inject drugs	8	3 %
45+	102	22 %	Heterosexual contact	46	15 %
Total	471	100 %	Total	312	100 %
			No Identified Risk	159	

# **AIDS CASES DIAGNOSED January - December 2014**

• 202 persons were newly diagnosed with AIDS in the New Orleans MSA.

<b>Gender</b>	Cases	Percent	Race/Ethnicity Cases	Perce	nt
Men	136	67 %	Black, Non-Hispanic 144	71	%
Women	62	31%	White, Non-Hispanic 41	20	%
Transgender Women	4	2 %	Hispanic 14	7	%
Total	202	100 %	Asian/Pacific Islander 1	<1	%
			Unknown 1	<1	%
			Multi-race 1	<1	%
			Total 202	100	%
Age at Diagnosis	Cases	s Percent	Cases with a reported risk (by birth sex):	Cases	Percent
13 - 24	13	6 %	Men who have sex with men	71	60 %
25 - 34	60	30 %	Injection drug use	14	12 %
35 - 44	56	5 28 %	Men who have sex with men and inject drugs	2	2 %
45+	73	36 %	Heterosexual contact	31	26 %
Total	202	2 100 %	Transfusion/Hemophilia	1	1 %
			Total	119	100 %

No Identified Risk

83

### Persons Diagnosed with Early Syphilis Metro New Orleans

Jefferson, Orleans, St. Tammany, St. John the Baptist, Plaquemines, St. Bernard, St. James, St. Charles Parishes

### PERSONS DIAGNOSED WITH P&S SYPHILIS January - March 2015

• 21 persons were diagnosed with P&S Syphilis in the New Orleans MSA from January through March 2015.

Sex at Birth	Cases	Percent	Race/Ethnicity	Cases	Percent
Female	1	5 %	American Indian	0	0 %
Male	20	95 %	Asian/Pacific Islander	0	0 %
Total	21	100 %	Black, Non-Hispanic	16	76 %
			Hispanic	0	0 %
			White, Non-Hispanic	5	24 %
			Other/Multi-race	0	0 %
			Unknown	0	0 %
			Total	21	100 %

Age at Diagnosis	Cases	Percent	Cases with a reported risk (Can have >1 risk):	Cases	Percent
0 - 12	0	0 %	Men who have sex with men and inject drugs	0	0 %
13 - 19	1	5 %	Men who have sex with men	15	75 %
20 - 29	13	62 %	Injection drug use	0	0 %
30 - 39	3	14 %	Transactional sex	0	0 %
40+	4	19 %	High risk heterosexual contact	2	10 %
Total	21	100 %	Heterosexual contact	4	20 %
			Incarceration	3	15 %
			Illegal drug use (non-injection)	7	35 %
			No Reported Risk	1	

### PERSONS DIAGNOSED WITH EARLY LATENT SYPHILIS January - March 2015

• 17 persons were diagnosed with Early Latent Syphilis in the New Orleans MSA from January through March 2015.

Sex at Birth	Cases	Percent	Race/Ethnicity	Cases	Percent
Female	4	24 %	American Indian	1	6 %
Male	13	76 %	Asian/Pacific Islander	0	0 %
Total	17	100 %	Black, Non-Hispanic	12	71 %
			Hispanic	1	6 %
			White, Non-Hispanic	3	18 %
			Other/Multi-race	0	0 %
			Unknown	0	0 %
			Total	17	100 %

Age at Diagnosis	Cases	Percent	Cases with a reported risk (Can have >1 risk):	Cases	Percent
0 - 12	0	0 %	Men who have sex with men and inject drugs	0	0 %
13 - 19	0	0 %	Men who have sex with men	12	71 %
20 - 29	4	24 %	Injection drug use	0	0 %
30 - 39	12	71 %	Transactional sex	0	0 %
40+	1	6 %	High risk heterosexual contact	1	6 %
Total	17	100 %	Heterosexual contact	4	24 %
			Incarceration	0	0 %
			Illegal drug use (non-injection)	7	41 %
			No Reported Risk	0	

### Persons Living with HIV Infection Metro Baton Rouge as of March 31, 2015

Ascension, East Baton Rouge, East Feliciana, Iberville, Livingston, Pointe Coupee, St. Helena, West Baton Rouge, West Feliciana Parishes

- **8,273** cumulative cases have been detected in the Baton Rouge MSA, including **80** pediatric cases.
- 3,584 cumulative deaths among persons with HIV infection have occurred in the Baton Rouge MSA, including 20 pediatric deaths.
- 5,171 persons are currently living with HIV infection in the Baton Rouge MSA.

Persons living with HIV: 2,386 Persons living with AIDS: 2,785

### **Gender**

	Cases	Percent
Women	1,836	36 %
Men	3,296	64 %
Transgender Women	39	<1%
Total	5,171	100 %

Age Category			Race/Ethnicity by Birth Sex	<u> </u>			
Current Age	Cases	Percent		Men	Women	Total	Percent
0 - 12	18	<1 %	Black, Non-Hispanic	2,605	1,644	4,249	82 %
13 - 19	34	1 %	White, Non-Hispanic	646	159	805	16 %
20 - 24	219	4 %	Hispanic	53	17	70	1 %
25 - 29	451	9 %	Asian/Pacific Islander	10	6	16	<1 %
30 - 34	606	12 %	American Indian	3	1	4	<1 %
35 - 39	641	12 %	Multi-race	18	9	27	1 %
40 - 44	647	13 %	Total	3,335	1,836	5,171	100 %
45 - 49	690	13 %					
50 - 54	751	15 %					
55 - 59	551	11 %					
60+	563	11 %					
Total	5,171	100 %					

Exposure Category by Birth Sex				
Cases with a reported risk:	Men	Women	Total	Percent
Men who have sex with men	1,372	0	1,372	43 %
Injection drug use	378	208	586	18 %
Men who have sex with men and inject drugs	194	0	194	6 %
Heterosexual contact	323	666	989	31 %
Transfusion/Hemophilia	6	8	14	<1 %
Perinatal	31	28	59	2 %
Total	2,304	910	3,214	100 %
No Identified Risk	1,031	926	1,957	

# Persons Newly Diagnosed with HIV and AIDS Metro Baton Rouge

Ascension, East Baton Rouge, East Feliciana, Iberville, Livingston, Pointe Coupee, St. Helena, West Baton Rouge, West Feliciana Parishes

# **HIV CASES DIAGNOSED January - December 2014**

• 278 persons were newly diagnosed with HIV in the Baton Rouge MSA.

<b>Gender</b>	Cases	Percent	Race/Ethnicity	Cases	Percent
Men	194	70 %	Black, Non-Hispanic	233	84 %
Women	80	29 %	White, Non-Hispanic	40	14 %
Transgender Women	4	1 %	Hispanic	3	1 %
Total	278	100 %	Asian/Pacific Islander	1	<1 %
			Multi-race	1	<1 %
			Total	278	100 %

Age at Diagnosis	Cases	Percent	Cases with a reported risk (by birth sex):	Cases	Percent
0 - 12	1	<1 %	Men who have sex with men	108	66 %
13 - 24	78	28 %	Injection drug use	15	9 %
25 - 34	74	27 %	Men who have sex with men and inject drugs	4	3 %
35 - 44	61	22 %	Heterosexual contact	34	21 %
45+	64	24 %	Perinatal	1	1 %
Total	278	100 %	Total	163	100 %
			No Identified Risk	115	

# **AIDS CASES DIAGNOSED January - December 2014**

• 196 persons were newly diagnosed with AIDS in the Baton Rouge MSA.

<b>Gender</b>	Cases	Percent	Race/Ethnicity	Cases	Percent
Men	138	70 %	Black, Non-Hispanic	175	89 %
Women	56	29 %	White, Non-Hispanic	20	10 %
Transgender Women	2	1 %	Multi-race	1	1 %
Total	196	100 %	Total	196	100 %

Age at Diagnosis	Cases	Percent	Cases with a reported risk (by birth sex):	Cases	Percent
0 - 12	2	1 %	Men who have sex with men	52	45 %
13 - 24	17	9 %	Injection drug use	21	18 %
25 - 34	57	29 %	Men who have sex with men and inject drugs	5	4 %
35 - 44	57	29 %	Heterosexual contact	35	30 %
45+	63	32 %	Perinatal	3	3 %
Total	196	100 %	Total	116	100 %

No Identified Risk 80

### Persons Diagnosed with Early Syphilis Metro Baton Rouge

Ascension, East Baton Rouge, East Feliciana, Iberville, Livingston, Pointe Coupee, St. Helena, West Baton Rouge, West Feliciana Parishes

### PERSONS DIAGNOSED WITH P&S SYPHILIS January – March 2015

• 22 persons were diagnosed with P&S Syphilis in the Baton Rouge MSA from January through March 2015.

Sex at Birth	Cases	Percent	Race/Ethnicity	Cases	Percent
Female	8	36 %	American Indian	0	0 %
Male	14	64 %	Asian/Pacific Islander	0	0 %
Total	22	100 %	Black, Non-Hispanic	18	82 %
			Hispanic	0	0 %
			White, Non-Hispanic	4	18 %
			Other/Multi-race	0	0 %
			Unknown	0	0 %
			Total	22	100 %

Age at Diagnosis	Cases	Percent	Cases with a reported risk (Can have >1 risk):	Cases	Percent
0 - 12	0	0 %	Men who have sex with men and inject drugs	0	0 %
13 - 19	1	5 %	Men who have sex with men	13	62 %
20 - 29	16	73 %	Injection drug use	0	0 %
30 - 39	2	9 %	Transactional sex	2	10 %
40+	3	14 %	High risk heterosexual contact	6	29 %
Total	22	100 %	Heterosexual contact	6	29 %
			Incarceration	3	14 %
			Illegal drug use (non-injection)	5	24 %
			No Reported Risk	1	

### PERSONS DIAGNOSED WITH EARLY LATENT SYPHILIS January – March 2015

• 10 persons were diagnosed with Early Latent Syphilis in the Baton Rouge MSA from January through March 2015.

Sex at Birth	Cases	Percent	Race/Ethnicity	Cases	Percent
Female	3	30 %	American Indian	0	0 %
Male	7	70 %	Asian/Pacific Islander	0	0 %
Total	10	100 %	Black, Non-Hispanic	9	90 %
			Hispanic	0	0 %
			White, Non-Hispanic	1	10 %
			Other/Multi-race	0	0 %
			Unknown	0	0 %
			Total	10	100 %

Age at Diagnosis	Cases	Percent	Cases with a reported risk (Can have >1 risk):	Cases	Percent
0 - 12	0	0 %	Men who have sex with men and inject drugs	0	0 %
13 - 19	0	7 %	Men who have sex with men	5	50 %
20 - 29	6	60 %	Injection drug use	0	0 %
30 - 39	3	30 %	Transactional sex	1	10 %
40+	1	10 %	High risk heterosexual contact	0	0 %
Total	10	100 %	Heterosexual contact	4	40 %
			Incarceration	2	20 %
			Illegal drug use (non-injection)	0	0 %
			No Reported Risk	0	

### Persons Living with HIV Infection Region 1: New Orleans Region as of March 31, 2015

Jefferson, Orleans, Plaquemines, St. Bernard Parishes

- 15,344 cumulative cases have been detected in Region 1, including 150 pediatric cases.
- 6,695 cumulative deaths among persons in Region 1 with HIV infection have occurred, including 55 pediatric deaths.
- 6,998 persons are currently living with HIV infection in Region 1.

Persons living with HIV: **3,250** Persons living with AIDS: **3,748** 

### **Gender**

	Cases	Percent
Women	1,777	25 %
Men	5,039	72 %
Transgender Women	90	1 %
Transgender Men	2	<1 %
Total	6,998	100 %

Age Category			Race/Ethnicity by Sex at I	<u> Birth</u>			
Current Age	Cases	Percent		Men	Women	Total	Percent
0 - 12	14	<1 %	Black, Non-Hispanic	2,951	1,477	4,428	63 %
13 - 19	43	1 %	White, Non-Hispanic	1,843	188	2,031	29 %
20 - 24	249	4 %	Hispanic	346	98	444	6 %
25 - 29	582	8 %	Asian/Pacific Islander	37	5	42	1 %
30 - 34	697	10 %	American Indian	8	0	8	<1 %
35 - 39	822	12 %	Unknown	4	2	6	<1 %
40 - 44	772	11 %	Multi-race	30	9	39	1 %
45 - 49	947	14 %	Total	5,219	1,779	6,998	100 %
50 - 54	1,085	16 %					
55 - 59	852	12 %					
60+	935	13 %					
Total	6,998	100 %					

Cases with a reported risk:	Men	Women	Total	Percent
Men who have sex with men	2,949	0	2,949	61 %
Injection drug use	293	192	485	10 %
Men who have sex with men and inject drugs	289	0	289	6 %
Heterosexual contact	329	722	1,051	22 %
Transfusion/Hemophilia	16	14	30	1 %
Perinatal	23	35	58	1 %
Total	3,899	963	4,862	100 %
No Identified Risk	1,320	816	2,136	

# Persons Newly Diagnosed with HIV and AIDS Region 1: New Orleans Region

Jefferson, Orleans, Plaquemines, St. Bernard Parishes

### **HIV CASES DIAGNOSED January - December 2014**

• 430 persons were newly diagnosed with HIV in Region 1.

<b>Gender</b>	Cases	Percent	Race/Ethnicity	Cases	Percent
Men	341	79 %	Black, Non-Hispanic	278	65 %
Women	83	19 %	White, Non-Hispanic	104	24 %
Transgender Women	6	1 %	Hispanic	38	9 %
Total	430	100 %	Asian/Pacific Islander	6	1 %
			Unknown	1	<1 %
			Multi-race	3	1 %
			Total	430	100 %

Age at Diagnosis	Cases	Percent	Cases with a reported risk (by birth sex):	Cases	Percent
13 - 24	112	26 %	Men who have sex with men	225	78 %
25 - 34	140	33 %	Injection drug use	11	4 %
35 - 44	84	20 %	Men who have sex with men and inject drugs	8	3 %
45+	94	22 %	Heterosexual contact	44	15 %
Total	430	100 %	Total	288	100 %

No Identified Risk 142

# **AIDS CASES DIAGNOSED January - December 2014**

• 184 persons were newly diagnosed with AIDS in Region 1.

<u>Gender</u>	Cases	Percent	Race/Ethnicity	Cases	Percent
Men	126	68 %	Black, Non-Hispanic	131	71 %
Women	55	30 %	White, Non-Hispanic	38	21 %
Transgender Women	1 3	2 %	Hispanic	12	7 %
Total	184	100 %	Asian/Pacific Islander	1	1 %
			Unknown	1	1 %
			Multi-race	1	1 %
			Total	184	100 %

Age at Diagnosis	Cases	Percent	Cases with a reported risk (by birth sex):	Cases	Percent
13 - 24	11	6 %	Men who have sex with men	66	59 %
25 - 34	56	30 %	Injection drug use	14	13 %
35 - 44	48	26 %	Men who have sex with men and inject drugs	2	2 %
45+	69	38 %	Heterosexual contact	28	25 %
Total	184	100 %	Transfusion/Hemophilia	1	1 %
			Total	111	100 %

No Identified Risk

73

### Persons Diagnosed with Early Syphilis Region 1: New Orleans Region

Jefferson, Orleans, Plaquemines, St. Bernard Parishes

### PERSONS DIAGNOSED WITH P&S SYPHILIS January – March 2015

• 21 persons were diagnosed with P&S Syphilis in Region 1 from January through March 2015.

Sex at Birth	Cases	Percent	Race/Ethnicity	Cases	Percent
Female	1	5 %	American Indian	0	0 %
Male	20	95 %	Asian/Pacific Islander	0	0 %
Total	21	100 %	Black, Non-Hispanic	16	76 %
			Hispanic	0	0 %
			White, Non-Hispanic	5	24 %
			Other/Multi-race	0	0 %
			Unknown	0	0 %
			Total	21	100 %

Age at Diagnosis	Cases	Percent	Cases with a reported risk (Can have >1 risk):	Cases	Percent
0 - 12	0	0 %	Men who have sex with men and inject drugs	0	0 %
13 - 19	1	5 %	Men who have sex with men	15	75 %
20 - 29	13	62 %	Injection drug use	0	0 %
30 - 39	3	14 %	Transactional sex	0	0 %
40+	4	19 %	High risk heterosexual contact	2	10 %
Total	21	100 %	Heterosexual contact	4	20 %
			Incarceration	3	15 %
			Illegal drug use (non-injection)	7	35 %
			No Reported Risk	1	

# PERSONS DIAGNOSED WITH EARLY LATENT SYPHILIS January – March 2015

• 16 persons were diagnosed with Early Latent Syphilis in Region 1 from January through March 2015.

Sex at Birth	Cases	Percent	Race/Ethnicity	Cases	Percent
Female	3	19 %	American Indian	1	6 %
Male	13	81 %	Asian/Pacific Islander	0	0 %
Total	16	100 %	Black, Non-Hispanic	12	75 %
			Hispanic	1	6 %
			White, Non-Hispanic	2	13 %
			Other/Multi-race	0	0 %
			Unknown	0	0 %
			Total	16	100 %

Age at Diagnosis	Cases	Percent	Cases with a reported risk (Can have >1 risk):	Cases	Percent
0 - 12	0	0 %	Men who have sex with men and inject drugs	0	0 %
13 - 19	0	0 %	Men who have sex with men	12	75 %
20 - 29	3	19 %	Injection drug use	0	0 %
30 - 39	12	75 %	Transactional sex	0	0 %
40+	1	6 %	High risk heterosexual contact	1	6 %
Total	16	100 %	Heterosexual contact	3	19 %
			Incarceration	0	0 %
			Illegal drug use (non-injection)	7	44 %
			No Reported Risk	0	

### Persons Living with HIV Infection Region 2: Baton Rouge Region as of March 31, 2015

Ascension, East Baton Rouge, East Feliciana, Iberville, Pointe Coupee, West Baton Rouge, West Feliciana Parishes

- **7,979** cumulative cases have been detected in Region 2, including **75** pediatric cases.
- **3,456** cumulative deaths among persons in Region 2 with HIV infection have occurred, including **17** pediatric deaths.
- 4,945 persons are currently living with HIV infection in Region 2.

Persons living with HIV: 2,284 Persons living with AIDS: 2,661

### **Gender**

	Cases	Percent
Women	1,769	36 %
Men	3,140	63 %
Transgender Women	36	1 %
Total	4,945	100 %

Age Category			Race/Ethnicity by Sex at I	<u> Birth</u>			
<b>Current Age</b>	Cases	Percent		Men	Women	Total	Percent
0 - 12	17	<1 %	Black, Non-Hispanic	2,543	1,621	4,164	84 %
13 - 19	32	1 %	White, Non-Hispanic	552	118	670	14 %
20 - 24	214	4 %	Hispanic	50	15	65	1 %
25 - 29	439	9 %	Asian/Pacific Islander	10	5	15	<1 %
30 - 34	590	12 %	American Indian	3	1	4	<1 %
35 - 39	612	12 %	Multi-race	18	9	27	1 %
40 - 44	615	12 %	Total	3,176	1,769	4,945	100 %
45 - 49	649	13 %					
50 - 54	715	14 %					
55 - 59	526	11 %					
60+	536	11 %					
Total	4,945	100 %					

Cases with a reported risk:	Men	Women	Total	Percent
Men who have sex with men	1,291	0	1,291	42 %
Injection drug use	365	195	560	18 %
Men who have sex with men and inject drugs	185	0	185	6 %
Heterosexual contact	308	641	949	31 %
Transfusion/Hemophilia	4	8	12	<1 %
Perinatal	30	26	56	2 %
Total	2,183	870	3,053	100 %
No Identified Risk	993	899	1,892	

# Persons Newly Diagnosed with HIV and AIDS Region 2: Baton Rouge Region

Ascension, East Baton Rouge, East Feliciana, Iberville, Pointe Coupee, West Baton Rouge, West Feliciana Parishes

# **HIV CASES DIAGNOSED January - December 2014**

• 344 persons were newly diagnosed with HIV in Region 2.

<b>Gender</b>	Cases	Percent	Race/Ethnicity	Cases	Percent
Men	236	69 %	Black, Non-Hispanic	285	83 %
Women	104	30 %	White, Non-Hispanic	39	11 %
Transgender Women	4	1 %	Hispanic	12	3 %
Total	344	100 %	Asian/Pacific Islander	6	2 %
			American Indian	1	<1 %
			Multi-race	1	<1 %
			Total	344	100 %

Age at Diagnosis	Cases	Percent	Cases with a reported risk (by birth sex):	Cases	Percent
0 - 12	1	<1 %	Men who have sex with men	115	67 %
13 - 24	80	23 %	Injection drug use	12	7 %
25 - 34	108	31 %	Men who have sex with men and inject drugs	2	1 %
35 - 44	72	21 %	Heterosexual contact	40	23 %
45+	83	24 %	Perinatal	3	2 %
Total	344	100 %	Total	172	100 %

No Identified Risk 172

# **AIDS CASES DIAGNOSED January - December 2014**

• 161 persons were newly diagnosed with AIDS in Region 2.

<b>Gender</b>	Cases	Percent	Race/Ethnicity	Cases	Percent
Men	108	67 %	Black, Non-Hispanic	139	86 %
Women	51	32 %	White, Non-Hispanic	15	9 %
Transgender Womer	n 2	1 %	Hispanic	6	4 %
Total	161	100 %	Asian/Pacific Islander	1	1 %
			Total	161	100 %

Age at Diagnosis	Cases	Percent	Cases with a reported risk (by birth sex):	Cases	Percent
13 - 24	14	9 %	Men who have sex with men	37	47 %
25 - 34	41	25 %	Injection drug use	11	14 %
35 - 44	48	30 %	Men who have sex with men and inject drugs	1	1 %
45+	58	36 %	Heterosexual contact	29	37 %
Total	161	100 %	Total	78	100 %

No Identified Risk

83

### Persons Diagnosed with Early Syphilis Region 2: Baton Rouge Region

Ascension, East Baton Rouge, East Feliciana, Iberville, Pointe Coupee, West Baton Rouge, West Feliciana Parishes

### PERSONS DIAGNOSED WITH P&S SYPHILIS January – March 2015

• 22 persons were diagnosed with P&S Syphilis in Region 2 from January through March 2015.

Sex at Birth	Cases	Percent	Race/Ethnicity	Cases	Percent
Female	8	36 %	American Indian	0	0 %
Male	14	64 %	Asian/Pacific Islander	0	0 %
Total	22	100 %	Black, Non-Hispanic	18	82 %
			Hispanic	0	0 %
			White, Non-Hispanic	4	18 %
			Other/Multi-race	0	0 %
			Unknown	0	0 %
			Total	22	100 %

Age at Diagnosis	Cases	Percent	Cases with a reported risk (Can have >1 risk):	Cases	Percent
0 - 12	0	0 %	Men who have sex with men and inject drugs	0	0 %
13 - 19	1	5 %	Men who have sex with men	13	62 %
20 - 29	16	73 %	Injection drug use	0	0 %
30 - 39	2	9 %	Transactional sex	2	10 %
40+	3	14 %	High risk heterosexual contact	6	29 %
Total	22	100 %	Heterosexual contact	6	29 %
			Incarceration	3	14 %
			Illegal drug use (non-injection)	5	24 %
			No Reported Risk	1	

# PERSONS DIAGNOSED WITH EARLY LATENT SYPHILIS January – March 2015

• 10 persons were diagnosed with Early Latent Syphilis in Region 2 from January through March 2015.

Sex at Birth	Cases	Percent	Race/Ethnicity	Cases	Percent
Female	3	30 %	American Indian	0	0 %
Male	7	70 %	Asian/Pacific Islander	0	0 %
Total	10	100 %	Black, Non-Hispanic	9	90 %
			Hispanic	0	0 %
			White, Non-Hispanic	1	10 %
			Other/Multi-race	0	0 %
			Unknown	0	0 %
			Total	10	100 %

Age at Diagnosis	Cases	Percent	Cases with a reported risk (Can have >1 risk):	Cases	Percent
0 - 12	0	0 %	Men who have sex with men and inject drugs	0	0 %
13 - 19	0	0 %	Men who have sex with men	5	50 %
20 - 29	6	60 %	Injection drug use	0	0 %
30 - 39	3	30 %	Transactional sex	1	10 %
40+	1	10 %	High risk heterosexual contact	0	0 %
Total	10	100 %	Heterosexual contact	4	40 %
			Incarceration	2	20 %
			Illegal drug use (non-injection)	0	0 %
			No Reported Risk	0	

### Persons Living with HIV Infection Region 3: Houma Region as of March 31, 2015

Assumption, Lafourche, St. Charles, St. James, St. John the Baptist, St. Mary, Terrebonne Parishes

- 1,228 cumulative cases have been detected in Region 3, including 19 pediatric cases.
- **564** cumulative deaths among persons in Region 3 with HIV infection have occurred, including **8** pediatric deaths.
- 810 persons are currently living with HIV infection in Region 3.

Persons living with HIV: 364
Persons living with AIDS: 446

<u>Gender</u>	Cases	Percent
Women	269	33 %
Men	537	66 %
Transgender Women	4	1 %
Total	810	100 %

Age Category			Race/Ethnicity by Sex at B	<u>irth</u>			
Current Age	Cases	Percent		Men	Women	Total	Percent
0 - 12	4	<1 %	Black, Non-Hispanic	285	199	484	60 %
13 - 19	10	1 %	White, Non-Hispanic	228	59	287	35 %
20 - 24	40	5 %	Hispanic	22	7	29	4 %
25 - 29	85	10 %	Asian/Pacific Islander	2	1	3	<1 %
30 - 34	91	11 %	American Indian	1	1	2	<1 %
35 - 39	92	11 %	Multi-race	3	2	5	1 %
40 - 44	96	12 %	Total	541	269	810	100 %
45 - 49	111	14 %					
50 - 54	103	13 %					
55 - 59	86	11 %					
60+	92	11 %					
Total	810	100 %					

Cases with a reported risk:	Men	Women	Total	Percent
Men who have sex with men	323	0	323	57 %
Injection drug use	33	18	51	9 %
Men who have sex with men and inject drugs	16	0	16	3 %
Heterosexual contact	34	122	156	28 %
Perinatal	8	9	17	3 %
Total	414	149	563	100 %
No Identified Risk	127	120	247	

# Persons Newly Diagnosed with HIV and AIDS Region 3: Houma Region

Assumption, Lafourche, St. Charles, St. James, St. John the Baptist, St. Mary, Terrebonne Parishes

# **HIV CASES DIAGNOSED January - December 2014**

• 60 persons were newly diagnosed with HIV in Region 3.

<b>Gender</b>	Cases	Percent	Race/Ethnicity	Cases	Percent
Men	43	72 %	Black, Non-Hispanic	31	52 %
Women	17	28 %	White, Non-Hispanic	23	38 %
Total	60	100 %	Hispanic	6	10 %
			Total	60	100 %

Age at Diagnosis	Cases	Percent	Cases with a reported risk (by birth sex):	Cases	Percent
0 - 12	1	2 %	Men who have sex with men	28	76 %
13 - 24	19	32 %	Injection drug use	2	5 %
25 - 34	20	33 %	Heterosexual contact	6	16 %
35 - 44	7	12 %	Perinatal	1	3 %
45+	13	22 %	Total	37	100 %
Total	60	100 %			
			No Identified Risk	23	

### **AIDS CASES DIAGNOSED January - December 2014**

• 28 persons were newly diagnosed with AIDS in Region 3.

<b>Gender</b>	Cases	Percent	Race/Ethnicity	Cases	Percent
Men	17	61 %	Black, Non-Hispanic	18	64 %
Women	10	36 %	White, Non-Hispanic	8	29 %
Transgender Womer	1 1	4 %	Hispanic	2	7 %
Total	28	100 %	Total	28	100 %

Age at Diagnosis	Cases	Percent	Cases with a reported risk (by birth sex):	Cases	Percent
13 - 24	1	4 %	Men who have sex with men	12	71 %
25 - 34	9	32 %	Injection drug use	2	12 %
35 - 44	9	32 %	Heterosexual contact	3	18 %
45+	9	32 %	Total	17	100 %
Total	28	100 %			
			No Identified Risk	11	

# Persons Diagnosed with Early Syphilis Region 3: Houma Region

Assumption, Lafourche, St. Charles, St. James, St. John the Baptist, St. Mary, Terrebonne Parishes

### PERSONS DIAGNOSED WITH P&S SYPHILIS January – March 2015

• 8 persons were diagnosed with P&S Syphilis in Region 3 from January through March 2015.

Sex at Birth	Cases	Percent	Race/Ethnicity	Cases	Percent
Female	5	63 %	American Indian	0	0 %
Male	3	37 %	Asian/Pacific Islander	0	0 %
Total	8	100 %	Black, Non-Hispanic	8	100 %
			Hispanic	0	0 %
			White, Non-Hispanic	0	0 %
			Other/Multi-race	0	0 %
			Unknown	0	0 %
			Total	8	100 %

Age at Diagnosis	Cases	Percent	Cases with a reported risk (Can have >1 risk):	Cases	Percent
0 - 12	0	0 %	Men who have sex with men and inject drugs	0	0 %
13 - 19	2	25 %	Men who have sex with men	0	0 %
20 - 29	5	63 %	Injection drug use	0	0 %
30 - 39	1	13 %	Transactional sex	0	0 %
40+	0	0 %	High risk heterosexual contact	2	25 %
Total	8	100 %	Heterosexual contact	6	75 %
			Incarceration	2	25 %
			Illegal drug use (non-injection)	6	75 %
			No Reported Risk	0	

# PERSONS DIAGNOSED WITH EARLY LATENT SYPHILIS January – March 2015

• 7 persons were diagnosed with Early Latent Syphilis in Region 3 from January through March 2015.

Sex at Birth	Cases	Percent	Race/Ethnicity	Cases	Percent
Female	2	29 %	American Indian	1	14 %
Male	5	71 %	Asian/Pacific Islander	0	0 %
Total	7	100 %	Black, Non-Hispanic	6	86 %
			Hispanic	0	0 %
			White, Non-Hispanic	0	0 %
			Other/Multi-race	0	0 %
			Unknown	0	0 %
			Total	7	100 %

Age at Diagnosis	Cases	Percent	Cases with a reported risk (Can have >1 risk):	Cases	Percent
0 - 12	0	0 %	Men who have sex with men and inject drugs	0	0 %
13 - 19	2	29 %	Men who have sex with men	0	0 %
20 - 29	4	57 %	Injection drug use	0	0 %
30 - 39	0	0 %	Transactional sex	0	0 %
40+	1	14 %	High risk heterosexual contact	4	57 %
Total	7	100 %	Heterosexual contact	3	43 %
			Incarceration	1	14 %
			Illegal drug use (non-injection)	2	29 %
			No Reported Risk	0	

### Persons Living with HIV Infection Region 4: Lafayette Region as of March 31, 2015

Acadia, Evangeline, Iberia, Lafayette, St. Landry, St. Martin, Vermilion Parishes

- 2,331 cumulative cases have been detected in Region 4, including 18 pediatric cases.
- 1,078 cumulative deaths among persons in Region 4 with HIV infection have occurred, including 10 pediatric deaths.
- 1,518 persons are currently living with HIV infection in Region 4.

Persons living with HIV: 718
Persons living with AIDS: 800

<u>Gender</u>	Cases	Percent
Women	373	25 %
Men	1,138	75 %
Transgender Women	7	<1 %
Total	1,518	100 %

Age Category			Race/Ethnicity by Sex at I	<u> Birth</u>			
Current Age	Cases	Percent		Men	Women	Total	Percent
13 - 19	7	<1 %	Black, Non-Hispanic	605	263	868	57 %
20 - 24	63	4 %	White, Non-Hispanic	465	96	561	37 %
25 - 29	128	8 %	Hispanic	55	6	61	4 %
30 - 34	140	9 %	Asian/Pacific Islander	4	2	6	<1 %
35 - 39	152	10 %	American Indian	2	2	4	<1 %
40 - 44	158	10 %	Unknown	0	1	1	<1 %
45 - 49	218	14 %	Multi-race	14	3	17	1 %
50 - 54	280	18 %	Total	1,145	373	1,518	100 %
55 - 59	189	12 %					
60+	183	12 %					
Total	1,518	100 %					

Cases with a reported risk:	Men	Women	Total	Percent
Men who have sex with men	583	0	583	58 %
Injection drug use	71	51	122	12 %
Men who have sex with men and inject drugs	59	0	59	6 %
Heterosexual contact	67	154	221	22 %
Transfusion/Hemophilia	5	1	6	1 %
Confirmed Other	0	1	1	<1 %
Perinatal	4	5	9	1 %
Total	789	212	1,001	100 %
No Identified Risk	356	161	517	

# Persons Newly Diagnosed with HIV and AIDS Region 4: Lafayette Region

Acadia, Evangeline, Iberia, Lafayette, St. Landry, St. Martin, Vermilion Parishes

### **HIV CASES DIAGNOSED January - December 2014**

• 120 persons were newly diagnosed with HIV in Region 4.

<b>Gender</b>	Cases	Percent	Race/Ethnicity	Cases	Percent
Men	93	78 %	Black, Non-Hispanic	84	70 %
Women	27	22 %	White, Non-Hispanic	29	24 %
Total	120	100 %	Hispanic	5	4 %
			Asian/Pacific Islander	1	1 %
			Multi-race	1	1 %
			Total	120	100 %

Age at Diagnosis	Cases	Percent	Cases with a reported risk (by birth sex):	Cases	Percent
13 - 24	31	26 %	Men who have sex with men	43	70 %
25 - 34	45	38 %	Injection drug use	6	10 %
35 - 44	15	12 %	Men who have sex with men and inject drugs	1	2 %
45+	29	24 %	Heterosexual contact	11	18 %
Total	120	100 %	Total	61	100 %

No Identified Risk 59

# **AIDS CASES DIAGNOSED January - December 2014**

• 47 persons were newly diagnosed with AIDS in Region 4.

<b>Gender</b>	Cases	Percent	Race/Ethnicity	Cases	Percent
Men	36	77 %	Black, Non-Hispanic	33	70 %
Women	11	23 %	White, Non-Hispanic	9	19 %
Total	47	100 %	Hispanic	3	6 %
			Asian/Pacific Islander	1	2 %
			Multi-race	1	2 %
			Total	47	100 %

Age at Diagnosis	Cases	Percent	Cases with a reported risk (by birth sex):	Cases	Percent
13 - 24	3	6 %	Men who have sex with men	15	56 %
25 - 34	15	32 %	Injection drug use	5	19 %
35 - 44	7	15 %	Men who have sex with men and inject drugs	1	4 %
45+	22	47 %	Heterosexual contact	6	22 %
Total	47	100 %	Total	27	100 %

No Identified Risk

20

# Persons Diagnosed with Early Syphilis Region 4: Lafayette Region

Acadia, Evangeline, Iberia, Lafayette, St. Landry, St. Martin, Vermilion Parishes

### PERSONS DIAGNOSED WITH P&S SYPHILIS January – March 2015

• 15 persons were diagnosed with P&S Syphilis in Region 4 from January through March 2015.

Sex at Birth	Cases	Percent	Race/Ethnicity	Cases	Percent
Female	3	20 %	American Indian	0	0 %
Male	12	80 %	Asian/Pacific Islander	0	0 %
Total	15	100 %	Black, Non-Hispanic	11	73 %
			Hispanic	0	0 %
			White, Non-Hispanic	4	27 %
			Other/Multi-race	0	0 %
			Unknown	0	0 %
			Total	15	100 %

Age at Diagnosis	Cases	Percent	Cases with a reported risk (Can have >1 risk):	Cases	Percent
0 - 12	0	0 %	Men who have sex with men and inject drugs	0	0 %
13 - 19	1	7 %	Men who have sex with men	7	54 %
20 - 29	9	60 %	Injection drug use	0	0 %
30 - 39	2	13 %	Transactional sex	1	8 %
40+	3	20 %	High risk heterosexual contact	2	15 %
Total	15	100 %	Heterosexual contact	4	31 %
			Incarceration	1	8 %
			Illegal drug use (non-injection)	3	23 %
			No Reported Risk	2	

### PERSONS DIAGNOSED WITH EARLY LATENT SYPHILIS January – March 2015

• 11 persons were diagnosed with Early Latent Syphilis in Region 4 from January through March 2015.

Sex at Birth	Cases	Percent	Race/Ethnicity	Cases	Percent
Female	5	45 %	American Indian	0	0 %
Male	6	55 %	Asian/Pacific Islander	0	0 %
Total	11	100 %	Black, Non-Hispanic	5	45 %
			Hispanic	1	9 %
			White, Non-Hispanic	5	45 %
			Other/Multi-race	0	0 %
			Unknown	0	0 %
			Total	11	100 %

Age at Diagnosis	Cases	Percent	Cases with a reported risk (Can have >1 risk):	Cases	Percent
0 - 12	0	0 %	Men who have sex with men and inject drugs	0	0 %
13 - 19	1	9 %	Men who have sex with men	4	36 %
20 - 29	6	55 %	Injection drug use	0	0 %
30 - 39	2	18 %	Transactional sex	0	0 %
40+	2	18 %	High risk heterosexual contact	2	18 %
Total	11	100 %	Heterosexual contact	6	55 %
			Incarceration	1	9 %
			Illegal drug use (non-injection)	3	27 %
			No Reported Risk	0	

### Persons Living with HIV Infection Region 5: Lake Charles Region as of March 31, 2015

Allen, Beauregard, Calcasieu, Cameron, Jefferson Davis Parishes

- 1,428 cumulative cases have been detected in Region 5, including 10 pediatric cases.
- 587 cumulative deaths among persons in Region 5 with HIV infection have occurred, including 2 pediatric deaths.
- 1,025 persons are currently living with HIV infection in Region 5.

Persons living with HIV: 470
Persons living with AIDS: 555

Gender	Cases	Percent
Women	245	24 %
Men	776	76 %
Transgender Women	4	<1%
Total	1.025	100%

Age Category			Race/Ethnicity by Sex at B	<u>irth</u>			
Current Age	Cases	Percent		Men	Women	Total	Percent
0 - 12	3	<1 %	Black, Non-Hispanic	412	162	574	56 %
13 - 19	7	1 %	White, Non-Hispanic	269	73	342	33 %
20 - 24	26	3 %	Hispanic	83	5	88	9 %
25 - 29	71	7 %	Asian/Pacific Islander	4	0	4	<1 %
30 - 34	96	9 %	American Indian	2	0	2	<1 %
35 - 39	131	13 %	Multi-race	10	5	15	1 %
40 - 44	131	13 %	Total	780	245	1,025	100 %
45 - 49	147	14 %					
50 - 54	183	18 %					
55 - 59	128	12 %					
60+	102	10 %					
Total	1,025	100 %					

Cases with a reported risk:	Men	Women	Total	Percent
Men who have sex with men	343	0	343	46 %
Injection drug use	88	34	122	17 %
Men who have sex with men and inject drugs	70	0	70	9 %
Heterosexual contact	84	106	190	26 %
Transfusion/Hemophilia	2	0	2	<1 %
Perinatal	5	8	13	2 %
Total	592	148	740	100 %
No Identified Risk	188	97	285	

# Persons Newly Diagnosed with HIV and AIDS Region 5: Lake Charles Region

Allen, Beauregard, Calcasieu, Cameron, Jefferson Davis Parishes

# **HIV CASES DIAGNOSED January - December 2013**

• 57 persons were newly diagnosed with HIV in Region 5.

<b>Gender</b>	Cases	Percent	Race/Ethnicity	Cases	Percent
Men	43	75 %	Black, Non-Hispanic	22	39 %
Women	14	25 %	White, Non-Hispanic	29	51 %
Total	57	100 %	Hispanic	4	7 %
			Asian/Pacific Islander	1	2 %
			Multi-race	1	2 %
			Total	57	100 %

Age at Diagnosis	Cases	Percent	Cases with a reported risk (by birth sex):	Cases	Percent
13 - 24	9	16 %	Men who have sex with men	22	61 %
25 - 34	19	33 %	Injection drug use	5	14 %
35 - 44	13	23 %	Men who have sex with men and inject drugs	3	8 %
45+	16	28 %	Heterosexual contact	6	17 %
Total	57	100 %	Total	36	100 %

No Identified Risk 21

# **AIDS CASES DIAGNOSED January - December 2013**

• 21 persons were newly diagnosed with AIDS in Region 5.

<u>Gender</u>	Cases	Percent	Race/Ethnicity	Cases	Percent
Men	15	71 %	Black, Non-Hispanic	7	33 %
Women	6	29 %	White, Non-Hispanic	12	57 %
Total	21	100 %	Hispanic	1	5 %
			Multi-race	1	5 %
			Total	21	100 %

Age at Diagnosis	Cases	Percent	Cases with a reported risk (by birth sex):	Cases	Percent
13 - 24	2	10 %	Men who have sex with men	7	64 %
25 - 34	9	43 %	Injection drug use	3	27 %
35 - 44	3	14 %	Heterosexual contact	1	9 %
45+	7	33 %	Total	11	100 %
Total	21	100 %			
			No Identified Risk	10	

# Persons Diagnosed with Early Syphilis Region 5: Lake Charles Region

Allen, Beauregard, Calcasieu, Cameron, Jefferson Davis Parishes

### PERSONS DIAGNOSED WITH P&S SYPHILIS January – March 2015

• 3 persons were diagnosed with P&S Syphilis in Region 5 from January through March 2015.

Sex at Birth	Cases	Percent	Race/Ethnicity	Cases	Percent
Female	1	33 %	American Indian	0	0 %
Male	2	67 %	Asian/Pacific Islander	0	0 %
Total	3	100 %	Black, Non-Hispanic	2	67 %
			Hispanic	0	0 %
			White, Non-Hispanic	1	33 %
			Other/Multi-race	0	0 %
			Unknown	0	0 %
			Total	3	100 %

Age at Diagnosis	Cases	Percent	Cases with a reported risk (Can have >1 risk):	Cases	Percent
0 - 12	0	0 %	Men who have sex with men and inject drugs	0	0 %
13 - 19	0	0 %	Men who have sex with men	1	33 %
20 - 29	1	33 %	Injection drug use	0	0 %
30 - 39	1	33 %	Transactional sex	1	33 %
40+	1	33 %	High risk heterosexual contact	1	33 %
Total	3	100 %	Heterosexual contact	1	33 %
			Incarceration	1	33 %
			Illegal drug use (non-injection)	1	33 %
			No Reported Risk	0	

# PERSONS DIAGNOSED WITH EARLY LATENT SYPHILIS January – March 2015

• 3 persons were diagnosed with Early Latent Syphilis in Region 5 from January through March 2015.

Sex at Birth	Cases	Percent	Race/Ethnicity	Cases	Percent
Female	0	0 %	American Indian	0	0 %
Male	3	100 %	Asian/Pacific Islander	0	0 %
Total	3	100 %	Black, Non-Hispanic	2	67 %
			Hispanic	0	0 %
			White, Non-Hispanic	1	33 %
			Other/Multi-race	0	0 %
			Unknown	0	0 %
			Total	3	100 %

Age at Diagnosis	Cases	Percent	Cases with a reported risk (Can have >1 risk):	Cases	Percent
0 - 12	0	0 %	Men who have sex with men and inject drugs	0	0 %
13 - 19	1	33 %	Men who have sex with men	3	100 %
20 - 29	2	67 %	Injection drug use	0	0 %
30 - 39	0	0 %	Transactional sex	0	0 %
40+	0	0 %	High risk heterosexual contact	1	33 %
Total	3	100 %	Heterosexual contact	0	0 %
			Incarceration	1	33 %
			Illegal drug use (non-injection)	1	33 %
			No Reported Risk	0	

### Persons Living with HIV Infection Region 6: Alexandria Region as of March 31, 2015

Avoyelles, Catahoula, Concordia, Grant, La Salle, Rapides, Vernon, Winn Parishes

- 1,576 cumulative cases have been detected in Region 6, including 15 pediatric cases.
- 579 cumulative deaths among persons in Region 6 with HIV infection have occurred, including 6 pediatric deaths.
- 911 persons are currently living with HIV infection in Region 6.

Persons living with HIV: 470
Persons living with AIDS: 441

### **Gender**

	Cases	Percent
Women	275	30 %
Men	633	69 %
Transgender	3	<1 %
Total	911	100%

Age Category			Race/Ethnicity by Sex at B	<u>irth</u>			
Current Age	Cases	Percent		Men	Women	Total	Percent
0 - 12	2	<1 %	Black, Non-Hispanic	404	216	620	68 %
13 - 19	4	<1 %	White, Non-Hispanic	180	50	230	25 %
20 - 24	35	4 %	Hispanic	45	6	51	6 %
25 - 29	70	8 %	Asian/Pacific Islander	3	0	3	<1 %
30 - 34	91	10 %	American Indian	0	1	1	<1 %
35 - 39	124	14 %	Multi-race	4	2	6	1 %
40 - 44	107	12 %	Total	636	275	911	100 %
45 - 49	120	13 %					
50 - 54	148	16 %					
55 - 59	109	12 %					
60+	101	11 %					
Total	911	100 %					

Cases with a reported risk:	Men	Women	Total	Percent
Men who have sex with men	269	0	269	44 %
Injection drug use	60	24	84	14 %
Men who have sex with men and inject drugs	33	0	33	5 %
Heterosexual contact	67	136	203	34 %
Transfusion/Hemophilia	5	2	7	1 %
Perinatal	5	4	9	1 %
Total	439	166	605	100 %
No Identified Risk	197	109	306	

# Persons Newly Diagnosed with HIV and AIDS Region 6: Alexandria Region

Avoyelles, Catahoula, Concordia, Grant, La Salle, Rapides, Vernon, Winn Parishes

# **HIV CASES DIAGNOSED January - December 2014**

• 75 persons were newly diagnosed with HIV in Region 6.

<u>Gender</u>	Cases	Percent	Race/Ethnicity	Cases	Percent
Men	58	77 %	Black, Non-Hispanic	37	49 %
Women	16	21 %	White, Non-Hispanic	24	32 %
Transgender Woman	1	1%	Hispanic	14	19 %
Total	75	100 %	Total	75	100 %

Age at Diagnosis	Cases	Percent	Cases with a reported risk (by birth sex):	Cases	Percent
13 - 24	12	16 %	Men who have sex with men	21	66 %
25 - 34	23	31 %	Injection drug use	2	6 %
35 - 44	23	31 %	Heterosexual contact	9	28 %
45+	17	23 %	Total	32	100 %
Total	75	100 %			
			No Identified Risk	43	

# **AIDS CASES DIAGNOSED January - December 2014**

• 25 persons were newly diagnosed with AIDS in Region 6.

<b>Gender</b>	Cases	Percent	Race/Ethnicity	Cases	Percent
Men	18	72 %	Black, Non-Hispanic	16	64 %
Women	7	28 %	White, Non-Hispanic	5	20 %
Total	25	100 %	Hispanic	4	16 %
			Total	25	100 %

Age at Diagnosis	Cases	Percent	Cases with a reported risk (by birth sex):	Cases	Percent
13 - 24	1	4 %	Men who have sex with men	8	73 %
25 - 34	5	20 %	Injection drug use	1	9 %
35 - 44	6	24 %	Heterosexual contact	2	18 %
45+	13	52 %	Total	11	100 %
Total	25	100 %			
			No Identified Risk	14	

# Persons Diagnosed with Early Syphilis Region 6: Alexandria Region

Avoyelles, Catahoula, Concordia, Grant, La Salle, Rapides, Vernon, Winn Parishes

### PERSONS DIAGNOSED WITH P&S SYPHILIS January – March 2015

• 6 persons were diagnosed with P&S Syphilis in Region 6 January through March 2015.

Sex at Birth	Cases	Percent	Race/Ethnicity	Cases	Percent
Female	0	0 %	American Indian	0	0 %
Male	6	100 %	Asian/Pacific Islander	0	0 %
Total	6	100 %	Black, Non-Hispanic	6	100 %
			Hispanic	0	0 %
			White, Non-Hispanic	0	0 %
			Other/Multi-race	0	0 %
			Unknown	0	0 %
			Total	6	100 %

Age at Diagnosis	Cases	Percent	Cases with a reported risk (Can have >1 risk):	Cases	Percent
0 - 12	0	0 %	Men who have sex with men and inject drugs	0	0 %
13 - 19	1	17 %	Men who have sex with men	4	67 %
20 - 29	3	50 %	Injection drug use	0	0 %
30 - 39	1	17 %	Transactional sex	0	0 %
40+	1	17 %	High risk heterosexual contact	1	17 %
Total	6	100 %	Heterosexual contact	2	33 %
			Incarceration	1	17 %
			Illegal drug use (non-injection)	3	50 %
			No Reported Risk	0	

# PERSONS DIAGNOSED WITH EARLY LATENT SYPHILIS January – March 2015

• 1 persons were diagnosed with Early Latent Syphilis in Region 6 January through March 2015.

Sex at Birth	Cases	Percent	Race/Ethnicity	Cases	Percent
Female	0	0 %	American Indian	0	0 %
Male	1	100 %	Asian/Pacific Islander	0	0 %
Total	1	100 %	Black, Non-Hispanic	1	100 %
			Hispanic	0	0 %
			White, Non-Hispanic	0	0 %
			Other/Multi-race	0	0 %
			Unknown	0	0 %
			Total	1	100 %

Age at Diagnosis	Cases	Percent	Cases with a reported risk (Can have >1 risk):	Cases	Percent
0 - 12	0	0 %	Men who have sex with men and inject drugs	0	0 %
13 - 19	0	0 %	Men who have sex with men	0	0 %
20 - 29	0	0 %	Injection drug use	0	0 %
30 - 39	0	0 %	Transactional sex	0	0 %
40+	1	100 %	High risk heterosexual contact	0	0 %
Total	1	100 %	Heterosexual contact	1	100 %
			Incarceration	1	100 %
			Illegal drug use (non-injection)	1	100 %
			No Reported Risk	0	

### Persons Living with HIV Infection Region 7: Shreveport Region as of March 31, 2015

Bienville, Bossier, Caddo, Claiborne, DeSoto, Natchitoches, Red River, Sabine, Webster Parishes

- 2,555 cumulative cases have been detected in Region 7, including 27 pediatric cases.
- 1,136 cumulative deaths among persons in Region 7 with HIV infection have occurred, including 12 pediatric deaths.
- 1,652 persons are currently living with HIV infection in Region 7.

Persons living with HIV: **841**Persons living with AIDS: **811** 

<u>Gender</u>	Cases	Percent
Women	493	30 %
Men	1,151	70 %
Transgender Women	8	<1 %
Total	1,652	100 %

<b>Age Category</b>			Race/Ethnicity by Sex at I	<u> Birth</u>			
Current Age	Cases	Percent		Men	Women	Total	Percent
0 - 12	1	<1 %	Black, Non-Hispanic	798	404	1,202	73 %
13 - 19	12	1 %	White, Non-Hispanic	322	78	400	24 %
20 - 24	107	6 %	Hispanic	21	5	26	2 %
25 - 29	210	13 %	Asian/Pacific Islander	3	0	3	<1 %
30 - 34	194	12 %	American Indian	8	2	10	1 %
35 - 39	224	14 %	Unknown	0	1	1	<1 %
40 - 44	160	10 %	Multi-race	7	3	10	1 %
45 - 49	200	12 %	Total	1,159	493	1,652	100 %
50 - 54	233	14 %					
55 - 59	152	9 %					
60+	159	10 %					
Total	1,652	100 %					

Cases with a reported risk:	Men	Women	Total	Percent
Men who have sex with men	664	0	664	58 %
Injection drug use	53	44	97	8 %
Men who have sex with men and inject drugs	80	0	80	7 %
Heterosexual contact	76	209	285	25 %
Transfusion/Hemophilia	2	3	5	<1 %
Perinatal	6	7	13	1 %
Total	881	263	1,144	100 %
No Identified Risk	278	230	508	

# Persons Newly Diagnosed with HIV and AIDS Region 7: Shreveport Region

Bienville, Bossier, Caddo, Claiborne, DeSoto, Natchitoches, Red River, Sabine, Webster Parishes

### **HIV CASES DIAGNOSED January - December 2014**

• 156 persons were newly diagnosed with HIV in Region 7.

<u>Gender</u>	Cases	Percent	Race/Ethnicity	Cases	Percent
Men	115	74 %	Black, Non-Hispanic	122	78 %
Women	39	25 %	White, Non-Hispanic	28	18 %
Transgender Women	2	1 %	Hispanic	2	1 %
Total	156	100 %	American Indian	1	1 %
			Multi-race	3	2 %
			Total	156	100 %

Age at Diagnosis	Cases	Percent	Cases with a reported risk (by birth sex):	Cases	Percent
13 - 24	46	29 %	Men who have sex with men	70	80 %
25 - 34	55	35 %	Injection drug use	4	5 %
35 - 44	28	18 %	Men who have sex with men and inject drugs	2	2 %
45+	27	17 %	Heterosexual contact	12	14 %
Total	156	100 %	Total	88	100 %
			No Identified Risk	68	

# AIDS CASES DIAGNOSED January - December 2014

• 53 persons were newly diagnosed with AIDS in Region 7.

<b>Gender</b>	Cases	Percent	Race/Ethnicity	Cases	Percent
Men	37	70 %	Black, Non-Hispanic	41	77 %
Women	15	28 %	White, Non-Hispanic	9	17 %
Transgender Womer	1 1	2 %	Hispanic	1	2 %
Total	53	100 %	Multi-race	2	4 %
			Total	53	100 %

Age at Diagnosis	Cases	Percent	Cases with a reported risk (by birth sex):	Cases	Percent
13 - 24	5	9 %	Men who have sex with men	17	65 %
25 - 34	20	38 %	Injection drug use	2	8 %
35 - 44	13	25 %	Men who have sex with men and inject drugs	2	8 %
45+	15	28 %	Heterosexual contact	5	19 %
Total	53	100 %	Total	26	100 %

No Identified Risk

27

## Persons Diagnosed with Early Syphilis Region 7: Shreveport Region

Bienville, Bossier, Caddo, Claiborne, DeSoto, Natchitoches, Red River, Sabine, Webster Parishes

#### PERSONS DIAGNOSED WITH P&S SYPHILIS January – March 2015

• 16 persons were diagnosed with P&S Syphilis in Region 7 January through March 2015.

Sex at Birth	Cases	Percent	Race/Ethnicity	Cases	Percent
Female	9	56 %	American Indian	0	0 %
Male	7	44 %	Asian/Pacific Islander	0	0 %
Total	16	100 %	Black, Non-Hispanic	14	87 %
			Hispanic	0	0 %
			White, Non-Hispanic	2	13 %
			Other/Multi-race	0	0 %
			Unknown	0	0 %
			Total	16	100 %

Age at Diagnosis	Cases	Percent	Cases with a reported risk (Can have >1 risk):	Cases	Percent
0 - 12	0	0 %	Men who have sex with men and inject drugs	0	0 %
13 - 19	4	25 %	Men who have sex with men	2	13 %
20 - 29	9	56 %	Injection drug use	0	0 %
30 - 39	1	6 %	Transactional sex	1	6 %
40+	2	13 %	High risk heterosexual contact	4	25 %
Total	16	100 %	Heterosexual contact	10	63 %
			Incarceration	1	6 %
			Illegal drug use (non-injection)	4	25 %
			No Reported Risk	0	

## PERSONS DIAGNOSED WITH EARLY LATENT SYPHILIS January – March 2015

• 16 persons were diagnosed with Early Latent Syphilis in Region 7 January through March 2015.

Sex at Birth	Cases	Percent	Race/Ethnicity	Cases	Percent
Female	9	56 %	American Indian	0	0 %
Male	7	44 %	Asian/Pacific Islander	0	0 %
Total	16	100 %	Black, Non-Hispanic	13	81 %
			Hispanic	0	0 %
			White, Non-Hispanic	3	19 %
			Other/Multi-race	0	0 %
			Unknown	0	0 %
			Total	16	100 %

Age at Diagnosis	Cases	Percent	Cases with a reported risk (Can have >1 risk):	Cases	Percent
0 - 12	0	0 %	Men who have sex with men and inject drugs	0	0 %
13 - 19	4	25 %	Men who have sex with men	2	13 %
20 - 29	10	63 %	Injection drug use	0	0 %
30 - 39	1	6 %	Transactional sex	0	0 %
40+	1	6 %	High risk heterosexual contact	4	25 %
Total	16	100 %	Heterosexual contact	10	63 %
			Incarceration	0	0 %
			Illegal drug use (non-injection)	2	13 %
			No Reported Risk	0	

#### Persons Living with HIV Infection Region 8: Monroe Region as of March 31, 2015

Caldwell, E. Carroll, Franklin, Jackson, Lincoln, Madison, Morehouse, Ouachita, Richland, Tensas, Union, W. Carroll Parishes

- 1,842 cumulative cases have been detected in Region 8, including 18 pediatric cases.
- 772 cumulative deaths among persons in Region 8 with HIV infection have occurred, including 4 pediatric deaths.
- 1,050 persons are currently living with HIV infection in Region 8.

Persons living with HIV: 496
Persons living with AIDS: 554

<u>Gender</u>	Cases	Percent
Women	366	35 %
Men	680	65 %
Transgender Women	4	<1 %
Total	1.050	100 %

Age Category			Race/Ethnicity by Sex at B	irth			
Current Age	Cases	Percent		Men	Women	Total	Percent
0 - 12	6	1 %	Black, Non-Hispanic	490	315	805	77 %
13 - 19	11	1 %	White, Non-Hispanic	165	46	211	20 %
20 - 24	46	4 %	Hispanic	12	4	16	2 %
25 - 29	90	9 %	Asian/Pacific Islander	3	1	4	<1 %
30 - 34	101	10 %	American Indian	3	2	5	<1 %
35 - 39	132	13 %	Unknown	1	0	1	<1 %
40 - 44	153	15 %	Multi-race	6	2	8	1 %
45 - 49	119	11 %	Total	680	370	1,050	100 %
50 - 54	146	14 %					
55 - 59	126	12 %					
60+	120	11 %					
Total	1,050	100 %					

## **Exposure Category by Sex at Birth**

Cases with a reported risk:	Men	Women	Total	Perce	ent
Men who have sex with men	361	0	361	51	%
Injection drug use	34	32	66	9	%
Men who have sex with men and inject drugs	41	0	41	6	%
Heterosexual contact	65	152	217	31	%
Transfusion/Hemophilia	1	3	4	1	%
Perinatal	3	10	13	2	%
Total	505	197	702	100	%
No Identified Risk	175	173	348		

#### Persons Newly Diagnosed with HIV and AIDS Region 8: Monroe Region

Caldwell, E. Carroll, Franklin, Jackson, Lincoln, Madison, Morehouse, Ouachita, Richland, Tensas, Union, W. Carroll Parishes

#### **HIV CASES DIAGNOSED January - December 2014**

• 103 persons were newly diagnosed with HIV in Region 8.

<u>Gender</u>	Cases	Percent	Race/Ethnicity	Cases	Percent
Men	64	62 %	Black, Non-Hispanic	84	82 %
Women	38	37 %	White, Non-Hispanic	16	16 %
Transgender Women	1	1 %	American Indian	2	2 %
Total	103	100 %	Multi-race	1	1 %
			Total	103	100 %

Age at Diagnosis	Cases	Percent	Cases with a reported risk (by birth sex):	Cases	Percent
13 - 24	28	27 %	Men who have sex with men	28	62 %
25 - 34	28	27 %	Injection drug use	1	2 %
35 - 44	24	23 %	Men who have sex with men and inject drugs	1	2 %
45+	23	22 %	Heterosexual contact	15	33 %
Total	103	100 %	Total	45	100 %

No Identified Risk 58

# **AIDS CASES DIAGNOSED January - December 2014**

• 51 persons were newly diagnosed with AIDS in Region 8.

<b>Gender</b>	Cases	Percent	Race/Ethnicity	Cases	Percent
Men	27	53 %	Black, Non-Hispanic	42	82 %
Women	24	47 %	White, Non-Hispanic	7	14 %
Total	51	100 %	American Indian	1	2 %
			Multi-race	1	2 %
			Total	51	100 %

Age at Diagnosis	Cases	Percent	Cases with a reported risk (by birth sex):	Cases	Percent
13 - 24	5	10 %	Men who have sex with men	9	36 %
25 - 34	12	24 %	Injection drug use	2	8 %
35 - 44	14	27 %	Heterosexual contact	14	56 %
45+	20	39 %	Total	25	100 %
Total	51	100 %			
			No Identified Risk	26	

## Persons Diagnosed with Early Syphilis Region 8: Monroe Region

Caldwell, E. Carroll, Franklin, Jackson, Lincoln, Madison, Morehouse, Ouachita, Richland, Tensas, Union, W. Carroll Parishes

#### PERSONS DIAGNOSED WITH P&S SYPHILIS January – March 2015

• 8 persons were diagnosed with P&S Syphilis in Region 8 January through March 2015.

Sex at Birth	Cases	Percent	Race/Ethnicity	Cases	Percent
Female	1	13 %	American Indian	0	0 %
Male	7	87 %	Asian/Pacific Islander	0	0 %
Total	8	100 %	Black, Non-Hispanic	7	87 %
			Hispanic	0	0 %
			White, Non-Hispanic	1	13 %
			Other/Multi-race	0	0 %
			Unknown	0	0 %
			Total	8	100 %

Age at Diagnosis	Cases	Percent	Cases with a reported risk (Can have >1 risk):	Cases	Percent
0 - 12	0	0 %	Men who have sex with men and inject drugs	0	0 %
13 - 19	1	13 %	Men who have sex with men	3	38 %
20 - 29	5	62 %	Injection drug use	0	0 %
30 - 39	1	13 %	Transactional sex	0	0 %
40+	1	13 %	High risk heterosexual contact	1	13 %
Total	8	100 %	Heterosexual contact	4	50 %
			Incarceration	0	0 %
			Illegal drug use (non-injection)	2	25 %
			No Reported Risk	0	

# PERSONS DIAGNOSED WITH EARLY LATENT SYPHILIS January – March 2015

• 11 persons were diagnosed with Early Latent Syphilis in Region 8 January through March 2015.

Sex at Birth	Cases	Percent	Race/Ethnicity	Cases	Percent
Female	4	36 %	American Indian	0	0 %
Male	7	64 %	Asian/Pacific Islander	0	0 %
Total	11	100 %	Black, Non-Hispanic	11	100 %
			Hispanic	0	0 %
			White, Non-Hispanic	0	0 %
			Other/Multi-race	0	0 %
			Unknown	0	0 %
			Total	11	100 %

Age at Diagnosis	Cases	Percent	Cases with a reported risk (Can have >1 risk):	Cases	Percent
0 - 12	0	0 %	Men who have sex with men and inject drugs	0	0 %
13 - 19	3	27 %	Men who have sex with men	3	27 %
20 - 29	6	55 %	Injection drug use	0	0 %
30 - 39	1	9 %	Transactional sex	0	0 %
40+	1	9 %	High risk heterosexual contact	3	27 %
Total	11	100 %	Heterosexual contact	5	45 %
			Incarceration	1	9 %
			Illegal drug use (non-injection)	3	27 %
			No Reported Risk	0	

#### Persons Living with HIV Infection Region 9: Hammond/Slidell Region as of March 31, 2015

Livingston, St. Helena, St. Tammany, Tangipahoa, Washington Parishes

- 1,630 cumulative cases have been detected in Region 9, including 19 pediatric cases.
- **708** cumulative deaths among persons in Region 9 with HIV infection have occurred, including **7** pediatric deaths.
- 1,203 persons are currently living with HIV infection in Region 9.

Persons living with HIV: 553 Persons living with AIDS: 650

<u>Gender</u>	Cases	Percent
Women	312	26 %
Men	884	73 %
Transgender Women	7	<1%
Total	1,203	100 %

Age Category			Race/Ethnicity by Sex at B	<u>irth</u>			
Current Age	Cases	Percent		Men	Women	Total	Percent
0 - 12	5	<1 %	Black, Non-Hispanic	414	189	603	50 %
13 - 19	7	1 %	White, Non-Hispanic	451	111	562	47 %
20 - 24	35	3 %	Hispanic	21	10	31	3 %
25 - 29	88	7 %	Asian/Pacific Islander	2	2	4	<1 %
30 - 34	98	8 %	American Indian	1	0	1	<1 %
35 - 39	119	10 %	Multi-race	2	0	2	<1 %
40 - 44	144	12 %	Total	891	312	1,203	100 %
45 - 49	184	15 %					
50 - 54	216	18 %					
55 - 59	171	14 %					
60+	136	11 %					
Total	1,203	100 %					

#### **Exposure Category by Sex at Birth**

Cases with a reported risk:	Men	Women	Total	Percent
Men who have sex with men	470	0	470	57 %
Injection drug use	54	49	103	12 %
Men who have sex with men and inject drugs	56	0	56	7 %
Heterosexual contact	66	113	179	22 %
Transfusion/Hemophilia	5	0	5	1 %
Perinatal	11	5	16	2 %
Total	662	167	829	100 %
No Identified Risk	229	145	374	

## Persons Newly Diagnosed with HIV and AIDS Region 9: Hammond/Slidell Region

Livingston, St. Helena, St. Tammany, Tangipahoa, Washington Parishes

## **HIV CASES DIAGNOSED January - December 2014**

• 76 persons were newly diagnosed with HIV in Region 9.

<b>Gender</b>	Cases	Percent	Race/Ethnicity	Cases	Percent
Men	57	75 %	Black, Non-Hispanic	46	61 %
Women	19	25 %	White, Non-Hispanic	25	33 %
Total	76	100 %	Hispanic	5	7 %
			Total	76	100 %

Age at Diagnosis	Cases	Percent	Cases with a reported risk (by birth sex):	Cases	Percent
13 - 24	15	20 %	Men who have sex with men	24	71 %
25 - 34	25	33 %	Injection drug use	4	12 %
35 - 44	18	24 %	Men who have sex with men and inject drugs	1	3 %
45+	18	24 %	Heterosexual contact	5	15 %
Total	76	100 %	Total	34	100 %

No Identified Risk 42

#### **AIDS CASES DIAGNOSED January - December 2014**

• 47 persons were newly diagnosed with AIDS in Region 9.

<b>Gender</b>	Cases	Percent	Race/Ethnicity	Cases	Percent
Men	34	72 %	Black, Non-Hispanic	31	66 %
Women	13	28 %	White, Non-Hispanic	13	28 %
Total	47	100 %	Hispanic	3	6 %
			Total	47	100 %

Age at Diagnosis	Cases	Percent	Cases with a reported risk (by birth sex):	Cases	Percent
13 - 24	5	11 %	Men who have sex with men	17	59 %
25 - 34	9	19 %	Injection drug use	3	10 %
35 - 44	16	34 %	Heterosexual contact	9	31 %
45+	17	36 %	Total	29	100 %
Total	47	100 %			
			No Identified Risk	18	

## Persons Diagnosed with Early Syphilis Region 9: Hammond/Slidell Region

Livingston, St. Helena, St. Tammany, Tangipahoa, Washington Parishes

#### PERSONS DIAGNOSED WITH P&S SYPHILIS January - March 2015

• 1 persons were diagnosed with P&S Syphilis in Region 9 January through March 2015.

Sex at Birth	Cases	Percent	Race/Ethnicity	Cases	Percent
Female	0	0 %	American Indian	0	0 %
Male	1	100 %	Asian/Pacific Islander	0	0 %
Total	1	100 %	Black, Non-Hispanic	1	100 %
			Hispanic	0	0 %
			White, Non-Hispanic	0	0 %
			Other/Multi-race	0	0 %
			Unknown	0	0 %
			Total	1	100 %

Age at Diagnosis	Cases	Percent	Cases with a reported risk (Can have >1 risk):	Cases	Percent
0 - 12	0	0 %	Men who have sex with men and inject drugs	0	0 %
13 - 19	0	0 %	Men who have sex with men	1	100 %
20 - 29	1	100 %	Injection drug use	0	0 %
30 - 39	0	0 %	Transactional sex	0	0 %
40+	0	0 %	High risk heterosexual contact	0	0 %
Total	1	100 %	Heterosexual contact	0	0 %
			Incarceration	0	0 %
			Illegal drug use (non-injection)	0	0 %
			No Reported Risk	0	


# PERSONS DIAGNOSED WITH EARLY LATENT SYPHILIS January – March 2015


• 2 persons were diagnosed with Early Latent Syphilis in Region 9 January through March 2015.

Sex at Birth	Cases	Percent	Race/Ethnicity	Cases	Percent
Female	1	50 %	American Indian	0	0 %
Male	1	50 %	Asian/Pacific Islander	0	0 %
Total	2	100 %	Black, Non-Hispanic	1	50 %
			Hispanic	0	0 %
			White, Non-Hispanic	1	50 %
			Other/Multi-race	0	0 %
			Unknown	0	0 %
			Total	2	100 %


Age at Diagnosis	Cases	Percent	Cases with a reported risk (Can have >1 risk):	Cases	Percent
0 - 12	0	0 %	Men who have sex with men and inject drugs	0	0 %
13 - 19	0	0 %	Men who have sex with men	0	0 %
20 - 29	1	50 %	Injection drug use	0	0 %
30 - 39	0	0 %	Transactional sex	0	0 %
40+	1	50 %	High risk heterosexual contact	0	0 %
Total	2	100 %	Heterosexual contact	2	100 %
			Incarceration	0	0 %
			Illegal drug use (non-injection)	0	0 %
			No Reported Risk	0	

10-Year Trends in New HIV Diagnoses and New AIDS Diagnoses Louisiana, (2005-2014)







10-Year Trends in Primary & Secondary Syphilis Diagnoses Louisiana, (2005-2014)



5-Year Trends in Early Latent Syphilis Diagnoses Louisiana, (2010-2014)



# 10-Year Trends in Persons Living With HIV Infection Louisiana (2005-2014)





# 2014 HIV and Syphilis Region 1—New Orleans

Syphilis is rising in Louisiana and specifically in the New Orleans region (Jefferson, Orleans, Plaquemines and St. Bernard parishes). According to the recently released Centers for Disease Control and Prevention (CDC) 2013 STD and HIV national surveillance profiles, Louisiana ranked 3<sup>rd</sup> in the nation for primary and secondary syphilis with a case rate of 9.2 per 100,000. Louisiana also ranked 3<sup>rd</sup> in the nation for HIV, with a case rate of 30.3 per 100,000.

#### **REGION 1—PRIMARY AND SECONDARY SYPHILIS**

- From 2012 to 2014, the number of primary and secondary (P&S) syphilis diagnoses in the New Orleans region has more than tripled (62 diagnoses in 2012, 188 diagnoses in 2014).
- In 2014, the New Orleans region had the highest number of P&S syphilis diagnoses in Louisiana.
- The proportion of new P&S syphilis diagnoses among men has increased from 87% in 2012 to 93% in 2014. In 2014, there were only 8 new diagnoses of P&S syphilis among women.
- In 2014, 66% of all P&S syphilis diagnoses were among blacks and an additional 29% were among whites.
- Almost 50% of all P&S syphilis diagnoses were among persons between the age of 20-29 years. A significant proportion of diagnoses (18%) were in persons 45 and older.
- From 2012 to 2014, the proportion of new P&S syphilis diagnoses among men who have sex with men (MSM) has increased from 66% of all new diagnoses to 76% of all new diagnoses.

#### **REGION 1—HIV**

- The New Orleans region also has a very high number of new HIV diagnoses. In 2013 and 2014, the region diagnosed over 430 new cases of HIV each year. The New Orleans region has the highest number of HIV diagnoses and the highest HIV diagnosis rate in the state.
- In the most recent CDC HIV Surveillance report, the New Orleans metropolitan area ranked 2<sup>nd</sup> in the nation for estimated HIV case rates in 2013 (43.4 per 100,000).
- The proportion of new HIV diagnoses among men has increased from 70% in 2012 to 81% in 2014.
- In 2014, 64% of persons diagnosed with HIV were black and an additional 25% were white. There were 39 new diagnoses among Hispanic/Latinos which accounted for 9% of all new HIV diagnoses in the region.

New P&S Syphilis and HIV Diagnoses, 2012—2014, New Orleans Region							
	2012	2013	2014				
New P&S Syphilis Diagnoses	62	93	188				
New HIV Diagnoses	340	439	460				
HIV Coinfected P&S Syphilis Diagnoses	30	49	100				



# 2014 HIV and Syphilis Region 1—New Orleans

#### REGION 1—P&S SYPHILIS AND HIV COINFECTION

- The proportion of P&S syphilis diagnoses that are coinfected with HIV is increasing in New Orleans.
- In 2012, 48% of P&S syphilis diagnoses were coinfected with HIV, and in 2014 this proportion increased to 53%. A new diagnosis of P&S syphilis in New Orleans is more likely to be coinfected with HIV than not.
- In 2014, among the P&S syphilis diagnoses that were coinfected with HIV, 72% had been diagnosed with HIV for a year or longer before they were diagnosed with syphilis.
- Of the HIV coinfected P&S syphilis diagnoses in 2014:
  - 97% were male, 3% female
  - 72% were black, 24% white, 3% Hispanic/Latino
  - 49% were between 20-29 years old, 24% were 30-39 years old, and 25% were 40 and older
  - 90% were men who have sex with men

#### REGION 1 - P&S SYPHILIS AND HIV AMONG MSM

- The majority of all new diagnoses of P&S syphilis and HIV are among MSM. New diagnoses among MSM accounted for 76% of P&S syphilis diagnoses, 80% of HIV diagnoses, and 90% of HIV coinfected P&S syphilis cases in the New Orleans region in 2014.
- The proportion of new diagnoses of P&S syphilis and HIV among MSM is on the rise nationally and that is being mirrored in New Orleans.
- The majority of new diagnoses among MSM are black and between the ages of 20-29 years old.

#### STD AND HIV SCREENING GUIDELINES

- Persons newly diagnosed with HIV or P&S syphilis should always be counseled on and tested for both diseases. Additional screening for gonorrhea and chlamydia should also be conducted.
- According to Louisiana law (Act 459), all pregnant women should be tested for HIV and syphilis
  at their first prenatal care visit, and repeat tests for both diseases should be conducted at the
  first prenatal care visit of the third trimester.
- Yearly screening for syphilis, chlamydia, gonorrhea and HIV should be conducted for all sexually active MSM.
- Repeat HIV and STD screening should be performed every three months in MSM with a diagnosis of an STD in the previous year or additional risk factors.
- Additional information on STD and HIV testing and treatment can be found online at www.cdc.gov/std/treatment.

For more information on Louisiana's STD and HIV epidemics, please visit:

www.HIV411.org or dhh.louisiana.gov/std